



**HAL**  
open science

## Plant and spider communities benefit differently from the presence of planted hedgerows in highway verges

Isabelle Le Viol, Romain Julliard, Christian Kerbiriou, Louis de Redon, Nathalie Carnino, Nathalie Machon, Emmanuelle Porcher

### ► To cite this version:

Isabelle Le Viol, Romain Julliard, Christian Kerbiriou, Louis de Redon, Nathalie Carnino, et al.. Plant and spider communities benefit differently from the presence of planted hedgerows in highway verges. *Biological Conservation*, 2008, 141 (6), pp.1581-1590. 10.1016/j.biocon.2008.04.003 . mnhn-02265400

**HAL Id: mnhn-02265400**

**<https://mnhn.hal.science/mnhn-02265400>**

Submitted on 9 Aug 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 PLANT AND SPIDER COMMUNITIES BENEFIT DIFFERENTLY FROM THE  
2 PRESENCE OF PLANTED HEDGEROWS IN HIGHWAY VERGES.

3

4 Isabelle LEVIOL\*, Romain JULLIARD, Christian KERBIRIOU, Louis de REDON,  
5 Nathalie CARNINO, Nathalie MACHON, Emmanuelle PORCHER.

6

7 National Museum of Natural History, Species conservation, restoration and monitoring  
8 of populations, CERSP-UMR5173, 55 rue Buffon, 75005 Paris, FRANCE.

9 [ileviol@mnhn.fr](mailto:ileviol@mnhn.fr); [julliard@mnhn.fr](mailto:julliard@mnhn.fr); [kerbiriou@mnhn.fr](mailto:kerbiriou@mnhn.fr); [redon@mnhn.fr](mailto:redon@mnhn.fr);

10 [ncarnino@club-internet.fr](mailto:ncarnino@club-internet.fr); [machon@mnhn.fr](mailto:machon@mnhn.fr); [porcher@mnhn.fr](mailto:porcher@mnhn.fr)

11

12 **Corresponding author:**

13 **Isabelle LEVIOL.**

14 Postal address: National Museum of Natural History, Species conservation, restoration  
15 and monitoring of populations, CERSP-UMR5173, 55 rue Buffon, 75005 Paris, France.

16 Telephone number: 0033 140 795 359 – Fax number: 0033 140 793 835 – E-mail:

17 ileviol@mnhn.fr

18

19

20 PLANT AND SPIDER COMMUNITIES BENEFIT DIFFERENTLY FROM THE  
21 PRESENCE OF PLANTED HEDGEROWS IN HIGHWAY VERGES.

22 **Abstract:** Road verges should play a crucial role as a refuge for native flora and fauna  
23 in human dominated landscapes. However, the influence of construction choices, such  
24 as plantation of woody species, on the biodiversity supported by roadsides has received  
25 little attention, although the presence of hedgerows in roadsides is likely to enhance  
26 their role as a refuge, notably for woodland species. Using standardised methods, we  
27 assessed the impact of planted hedgerows on two taxonomic groups (plants and spiders)  
28 inhabiting highway verges within an intensive agricultural landscape. We examined  
29 community richness, taxonomic and functional composition in sites with and without  
30 planted hedgerows. At the site level, the response of plant and spider communities to  
31 the presence of planted hedgerows differed markedly: hedgerows were associated with  
32 significantly higher plant richness (higher  $\alpha$ -diversity), but similar spider richness. Plant  
33 communities in sites without hedgerows appeared as a subset of communities in sites  
34 with hedgerows, whereas spider communities in non-planted sites were complementary  
35 to that of planted sites (increased  $\beta$ -diversity). The presence of planted hedgerows was  
36 also associated with increased taxonomic and functional trait diversity at the landscape  
37 level ( $\gamma$ -diversity), through an increased  $\beta$ -diversity in both plants and spiders. Our  
38 results thus suggest that a mosaic of planted hedgerows and grassland habitats is crucial  
39 for the maintenance of biodiversity at a landscape scale. By providing information for  
40 road practitioners and policy makers regarding their potential impact on biodiversity,  
41 these results have important direct implications for the management of road networks.

42 **Key words:**  $\alpha$  and  $\beta$ -diversity, Biodiversity, Functional traits, Roadside management,  
43 Revegetation, RLQ analysis.

44

## 45 **1. Introduction**

46

47 Road networks, which have expanded over large areas with human population growth  
48 (Watts et al., 2007), are known to influence landscape structure and have major negative  
49 impacts on ecosystems dynamics (Trombulak and Frissel, 2000). However, the potential  
50 biological value of road verges has also long been recognised (Way, 1977) with  
51 possible contribution to the conservation of indigenous flora (Spooner et al., 2004;  
52 O'Farrell and Milton, 2006) and fauna (Meunier et al., 2000; Ries et al., 2001). This  
53 role as a refuge depends on the surrounding landscape: in natural habitats, generally  
54 supporting a high species diversity, road verges may not serve as a refuge (O'Farrell  
55 and Milton, 2006) and even have negative effects, notably by promoting invasion by  
56 non-indigenous species (Hansen and Clevenger, 2005; Brown, 2006). In contrast, in  
57 human dominated areas such as intensive agricultural landscapes, where non-  
58 agricultural habitats (e.g. edges) are critical to the conservation of biological diversity  
59 and ecological processes (Burel, 1996), road verges should play a crucial role as a  
60 refuge and as ecological corridors (Tikka et al., 2001; Smart et al., 2006).

61 Road verges support intensively managed habitats, which are artificially created  
62 on bare soil after road construction (restoration) and subsequently experience frequent  
63 mowing. This has direct consequences on the communities they can support and there is  
64 a need to identify and promote biodiversity-friendly construction and management  
65 practices, in order to optimise the role of road verges as a refuge for wild flora and  
66 fauna. As in any habitat, community composition in road verges is controlled by an  
67 array of distinct factors acting as filters at both landscape and local scales (Bochet et al.,  
68 2007), few of which can easily be manipulated to optimise biodiversity conservation.

69 There have been extensive studies addressing the influence of management on roadside  
70 biodiversity, notably mowing frequency and timing (Schaffers, 2002; Jantunen et al.,  
71 2007). In contrast, construction choices have received much less attention, with the  
72 exception of revegetation with herbaceous species (Matesanz et al., 2006; Leps et al.,  
73 2007). In particular, the role of planted trees in the maintenance of diversity in road  
74 verges was never considered to our knowledge, even though the presence of hedgerows  
75 along roadsides is likely to enhance their role as a refuge, by increasing the diversity of  
76 microhabitats. This role of hedgerows as a refuge however depends on their ability to  
77 receive a significant proportion of the local species pool and of the functional diversity  
78 they shelter (Roy and de Blois, 2006). Outside of roadsides, hedgerows appear to be a  
79 sustainable habitat and even functional corridors (flora: Corbit et al., 1999, Roy and de  
80 Blois, 2008; fauna: Maudsley, 2000, Griffiths et al., 2007; Davies and Pullin, 2007 for a  
81 review) even though other authors consider hedgerows as sink habitats (McCollin et al.,  
82 2000; Sitzia, 2007).

83 The aim of the present study is to evaluate the effect of hedgerows on the  
84 surrounding biodiversity of highway verges at different spatial scales, and on the  
85 distribution of biodiversity within and among sites in the context of an intensive  
86 agricultural landscape. Highway verges offer ideal settings to study the influence of  
87 hedgerows, because, in contrast to other habitats, the influence of history on community  
88 composition is strongly reduced, due to common history of all study sites. To address  
89 the impact of hedgerows on biodiversity, we compared the community richness and  
90 composition in sites with and without planted hedgerows for two taxonomic groups  
91 (plants and spiders), because no single taxon can inform on the ecological behaviour of  
92 other groups (Lowell et al., 2007); In addition, in the context of the search for

93 biodiversity indicators, it is of great importance to better assess the characteristics of  
94 different taxa (Pearman and Weber, 2007). These two groups differ in their ecological  
95 requirements and dispersal capabilities and are likely differentially affected by the  
96 presence of hedgerows: plant community composition is expected to be controlled  
97 mostly by soil and light characteristics (Schaffers, 2002) whereas spider communities  
98 are known to be highly sensitive to habitat structure and microclimatic conditions (Marc  
99 et al., 1999; Entling et al., 2007; see also Beals, 2006) and may exhibit variation on  
100 much smaller time scales than plants. By comparing similar grassland habitats in sites  
101 with and without planted hedgerows, we specifically examined (1) whether the presence  
102 of planted hedgerows influenced the surrounding specific and functional trait diversity,  
103 (2) whether plant and spider communities exhibited different environmental patterns,  
104 and (3) whether artificial hedgerows resulted in a higher overall biodiversity supported  
105 by highway verges.

106

## 107 **2. Methods**

108

### 109 **2.1. Study area**

110

111 This study was conducted within a 50 km section of highway A11 (1°39'30''E –  
112 48°29'15''N to 1°03'E – 48°14'N) in the Beauce region, an intensive agricultural  
113 landscape, West of Paris, France. The Beauce covers about 6 000 km<sup>2</sup> and is  
114 characterised by a temperate climate and high base status soils. This study area is a  
115 fairly typical modern open field landscape with a complete absence of hedgerows and

116 very few semi-natural habitats, such as remnant woodland patches. Agricultural  
117 practices consist mostly of intensive cereal production.

118 A11 Highway was built thirty years ago and enlarged in 1992. At this time, the  
119 newly created verges, devoid of vegetation, were sown with the same seed mixture,  
120 including mainly *Leguminosae* and *Poaceae* species (*Festuca rubra*, *Festuca ovina*,  
121 *Lolium perenne*, *Festuca arundinacea*, *Bromus inermis*, but also *Medicago lupulina*,  
122 *Lotus corniculatus*, *Trifolium arvense* and *Achillea millefolium*), along the whole  
123 highway section studied, and were planted with trees at discrete locations (*Cofiroute*  
124 construction company, personal communication). In the following, we define  
125 hedgerows as linear patches of planted shrubs and trees longer than 150 m, following  
126 the definition of Baudry et al. (2000). Sites with and without planted hedgerows are  
127 subsequently referred to as “WT” sites and “WO” sites, respectively. Management  
128 practices of highway verges depend on distance from the carriageway edge: all verges  
129 receive a visibility cut up to 6 m from the road edge twice a year, but areas behind this  
130 safety zone are only cut once a year, whereas vegetation growing under planted trees is  
131 seldom, if ever, cut.

132

## 133 **2.2. Field sampling and data collection**

134

### 135 2.2.1. Field sampling

136 We selected 25 field sites (Fig. 1) based on the following structural and biological  
137 criteria: highway sections were “roadcuts” (resulting from excavation), with edges of  
138 the same shape (width = 16.1 m ± 0.7 SE, height = 4 m ± 0.3 SE), slope (30° ± 2.4 SE)  
139 and aspect, and were bordered by crop fields. Within the 50 km study area, we selected

140 11 sites with planted hedgerows (WT) along these criteria, using geomorphic maps,  
141 aerial photographs, and field surveys. Hedgerows consisted of mature deciduous trees in  
142 the canopy layer, and were on average above 3 meters tall and 5 meters wide ( $5.2 \pm 0.5$   
143 SE, varying between 3 and 8 meters). They were generally pluri-specific, including  
144 most frequently the following planted species: *Acer campestre*, *Acer platanoides*, *Acer*  
145 *pseudoplatanus*, *Alnus cordata*, *Betula verrucosa*, *Betula pendula*, *Carpinus betulus*,  
146 *Cornus alba*, *Cornus mas*, *Cornus sanguinea*, *Corylus avellana*, *Cotinus coggygria*,  
147 *Fraxinus excelsior*, *Prunus avium*, *Prunus mahaleb*, *Prunus padus*, *Prunus spinosa*,  
148 *Quercus sessiliflora*, *Quercus robur*, *Robinia pseudo-acacia*, *Salix caprea*, *Sambucus*  
149 *nigra*, *Sorbus aucuparia*, *Sorbus torminalis*, *Viburnum lantana*, *Viburnum opulus*.  
150 When possible, one site without hedgerows (WO) was chosen within 1.5 km of a given  
151 site with hedgerows (WT). We thus selected 7 pairs of geographically close sites with  
152 and without hedgerows and completed this sample with isolated WT ( $n = 4$ ) and WO  
153 sites ( $n = 7$ ), for a total of 25 sampled sites (11 WT - 14 WO). WO sites were always  
154 located at least 50 m from the closest tree.

155

#### 156 2.2.2. Data collection

157 All inventories were performed through a 60 m transect running parallel to the  
158 carriageway, located at least 7.5 m from the carriageway edge and at  $8.2 \pm 0.6$  SE from  
159 the neighbouring crop field. Such location corresponded to extensive management  
160 practices. In WT sites, transects were also located 0.5 to 1 m from the canopy of planted  
161 trees (vertical projection of lower branches), and started and ended at least 50 m away  
162 from the end of the hedgerow, to avoid border effects.



163 For flora, sites were surveyed from May 10th to June 10th 2006. In each site, all  
164 vascular plant species were inventoried once in five 1 m<sup>2</sup> quadrats regularly distributed  
165 over the 60 m transect. Almost all taxa were identified to species level according to the  
166 International Plant Names Index. A few species that were difficult to distinguish when  
167 not in flower or fruit (*Vicia gr tetrasperma-hirsuta* for example) were grouped.

168 Spiders were sampled using five pitfall traps per site. Traps were placed within  
169 30 cm of each vegetation quadrat on the 60 m transect and were left in place from May  
170 10 to June 10 2006. Each trap was 8.5 cm wide in diameter and 10 cm deep, and  
171 contained ethylene glycol, a non-attractive preservative. Two pitfall traps were  
172 destroyed and were omitted in the analyses (n = 123). Adult spiders were identified to  
173 species level following the nomenclature of Platnick (2007). We failed to identify three  
174 individuals to species level; nevertheless, because they were obviously different from  
175 other species, they were conserved for analyses. Spider juveniles were generally not  
176 identifiable to species level and were not considered.

177 Sites characteristics: sampling design was defined to test impact of hedgerows,  
178 so that the variation of other environmental variables was minimised. We recorded sites  
179 characteristics and checked the absence of correlation between presence/absence of  
180 hedgerows and other environmental factors using pearson correlations, to check that the  
181 effects of hedgerows were not confounded with other environmental factors. For each  
182 site, we recorded geographic coordinates, verge width (final variation: 10-20 m),  
183 adjacent land use type (crop vs. temporary fallow), slope, distance from crop edge,  
184 aspect, and soil nutrient status. The latter was characterised via determination of soil  
185 pH, as well as phosphorus (P<sub>2</sub>O<sub>5</sub>, Olsen %), salt (Na<sub>2</sub>O), and total nitrogen (%)  
186 concentration in a bulked representative soil sample, taken to a depth of 10 cm in each

187 of five plots and subsequently mixed. For each site, we also recorded the distance from  
188 the nearest woodland patch ( $411.6 \text{ m} \pm 70.4 \text{ SE}$ ), as well as the total woodland area  
189 within 250 m, 500 m and 1 000 m radius buffers, using aerial photographs and the  
190 Geographical Information System package ARCGIS 8 (ESRI, 2000). In this open-field  
191 system, woodland patches can be considered as the main natural habitat in Beauce.  
192 Consequently, we considered that they could be important reservoirs of species  
193 compared to others habitats, providing a pool of both plants and spiders from which  
194 species were able to disperse and potentially colonise highway verges.

195

### 196 **2.3. Species attributes**

197

198 To address the influence of hedgerows on community functional composition, we  
199 characterised species based on readily available functional traits. As such data were  
200 analyzed using multivariate methods that are highly sensitive to “rare” species or  
201 classes, we chose to group some classes to minimize the resulting bias (see caption of  
202 fig. 2). For plants, these functional traits include Raunkier types, maximum adult height,  
203 and seed dispersal mode, scored according to a literature review (supplementary  
204 material 1). For spiders, we retained hunting strategies, body size and aerial-dispersal  
205 behaviour (ballooning vs. non-ballooning species). Hunting strategies (diurnal cursorial  
206 hunters, nocturnal cursorial hunters, ambush hunters, and web-builders) and species-  
207 specific body sizes were derived from Roberts (1985, 1987). As body sizes usually  
208 differ between sexes, the mean value of male and female median sizes was retained.  
209 Species exhibiting aerial dispersal using silk (ballooning) are considered species with  
210 great dispersal capabilities (Bell et al., 2005 for review) and were defined using the

211 world species list of ballooning Araneomorphae established by Bell et al. (2005). All  
212 species that did not appear in this list were considered “non ballooners”, which may be  
213 incorrect because the list is not exhaustive. However, as we worked with common  
214 species frequently found in agricultural landscapes and whose biology is generally well-  
215 known, we expect that this bias should be small. For both plants and spiders, a given  
216 taxon was assigned attributes only when it was identified to species level and observed  
217 in more than four instances (quadrats or traps), which yielded a total of 46 plant species  
218 (100% of species observed in more than four quadrats) and 56 spider species (84% of  
219 species detected in more than four pitfalls) for which all traits were scored.

220

#### 221 **2.4. Data analysis**

222

223 To remove the obvious increase in plant species richness due to the simple presence of  
224 planted species and examine solely indirect effects of planted hedgerows on plant  
225 community composition, we chose to work with spontaneous plant species only. Hence,  
226 unless otherwise stated, shrub or tree species that were observed in the herbaceous strata  
227 in a site where they had been planted (as inferred from historical plantation map data,  
228 *Cofiroute – Vegetude*) were discarded in subsequent analyses, i.e. they were excluded of  
229 this particular site; not however that these species were retained in other sites. In  
230 addition, for spider species, we chose to work with presence/absence data only, so that a  
231 range of different analyses, some of which valid with presence/absence data only, could  
232 be used on the same dataset. With few exceptions, including abundance data yielded  
233 very similar results. All analyses were conducted with R (R Development Core Team,  
234 2007).

235

#### 236 2.4.1. Species richness

237 To assess the effect of hedgerows at different spatial scales, we considered species  
238 richness at the quadrat scale ( $S_{\text{local}}$ , plants only), at the site scale ( $S_{\text{site}}$ , in a transect of 60  
239 m, plants and spiders) and at the highway section scale, within each type of site ( $S_{\text{type}}$ ,  
240 within WT or WO types of sites, plants and spiders). For plants, we considered local  
241 species richness ( $S_{\text{local}}$ ) within quadrats because we assumed that all species were  
242 detected, whereas for spiders, pitfall trapping is known to be influenced by the activity-  
243 density of species (Melbourne, 1999). In addition, analysis on local spider richness  
244 showed no effect of any environmental variable. Comparing observed species richness  
245 across sites and habitats might not be valid in case of unequal species detection  
246 probabilities (Boulinier et al., 1998; Nichols et al., 1998). We therefore used statistical  
247 methods derived from capture-recapture approaches in population or community  
248 dynamics to estimate species richness and changes in community composition As in  
249 recent studies (Lekve et al., 2002; Doherty et al., 2003; Selmi and Boulinier, 2003;  
250 Devictor et al., 2007; Kerbiriou et al., 2008) addressing richness estimation and  
251 differences in community composition from species count data, we used the program  
252 COMDYN (Hines et al., 1999), based on the jack-knife estimator of Burnham and  
253 Overton (1979). To estimate species richness at the site level ( $S_{\text{site}}$ ), we used quadrat or  
254 pitfall captures as replicates. To estimate richness at type of site level ( $S_{\text{type}}$  within WT  
255 or WO type of sites), we used sites as replicates ( $n = 11$ ): analyses were based on  
256 presence-absence data at the site level, pooled across the five quadrats for plants and the  
257 five pitfall traps for spiders. As 11 replicates only were available for WT sites, we  
258 sampled randomly without replacement 11 sites out of 14 in the WO group; this

259 sampling procedure was performed 15 times. Each of 15 pairs of (1) 11 sampled WO  
260 sites plus (2) the 11 WT sites was input into program COMDYN to obtain estimated  
261 richness per type of site. The estimated richness in each type of site was then  
262 characterised using the mean and standard error of the 15 outputs.

263 We analysed the combined influence of hedgerows and environmental variables  
264 on both local species richness for plants ( $S_{\text{local}}$ ) and estimated richness per site for  
265 plants and spiders ( $S_{\text{site}}$ ), using linear models and analyses of variance, as follows: for  
266 each taxonomic group (plants and spiders) and dependent variable ( $S_{\text{local}}$  and  $S_{\text{site}}$ ), we  
267 selected the most informative model using a stepwise algorithm based on Akaike  
268 Information Criterion (StepAIC, library MASS in R) starting from the most complete  
269 model including all environmental variables likely to influence species richness (listed  
270 above : see “site characteristics”). Plant richness was included as an additional  
271 environmental variable for spiders. Data were log-transformed when necessary to meet  
272 normality assumptions; the effects of variables retained by the stepwise algorithm were  
273 tested adjusted to others and the nested structure of the data was taken into account  
274 (local richness per quadrat nested within site: lme, library nlme in R).

275

#### 276 2.4.2. Species composition

277 We investigated a possible spatial autocorrelation in community composition by  
278 performing mantel tests (mantel.randtest, library ade4 in R) to assess the correlation  
279 between matrices of geographic vs. specific distances among sites. For plants and  
280 spiders, matrices of specific distances across sites were computed with the Jaccard  
281 index on presence/absence matrices. This effect was also tested separately within both  
282 types of edges (WT and WO sites). For these analyses, all species were retained.

283           We also examined the effect of hedgerows on the composition of plant and  
284 spider communities in the herbaceous strata using two types of multivariate analyses:  
285 Constrained Analysis of Principal Coordinates (CAP) and partial Constrained  
286 Correspondence Analysis (pCCA), followed-up by ANOVA-like permutation tests ( $n =$   
287 999) to assess significance of effects. CAP (capscale, library vegan in R, see Anderson  
288 and Willis, 2003) is an ordination method similar to Redundancy Analysis (RDA), i.e.  
289 exploring the relationship between two sets of variables, but allowing non-Euclidean  
290 dissimilarity indices. It was used to examine whether species similarity among sites  
291 depended on the presence/absence of hedgerows, and on the distance to nearest  
292 woodland patch, which was used as a covariable. It was performed on presence/absence  
293 matrices across sites, using the Jaccard index. pCCA (cca, library vegan in R, ter Braak,  
294 1986) is an ordination method based on CCA, a constrained ordination method  
295 developed to relate community composition to known variation in the environment. It is  
296 considered as a good choice if the user has clear and strong *a priori* hypotheses on  
297 constraints. Here, we used pCCA to test whether the presence of some species was  
298 associated with the presence or absence of planted hedgerows. We chose pCCA instead  
299 of a simple CCA to remove the effect of the distance to the nearest woodland patch,  
300 used as a covariable, as in CAP. As both analyses are sensitive to rare species, we only  
301 retained species observed in more than four samples, i.e. 46 plant species and 67 spider  
302 species.

303           Third, we examined the complementarity of communities (Nichols et al., 1998)  
304 in each type of site (WO type and WT type) using COMDYN with sites as replicates ( $n$   
305 = 11). In addition to estimating richness within each type of site ( $S_{\text{type}}$  within WO and  
306 WT type of sites), we obtained the following estimators: complement of “extinction

307 probability” between the two types,  $\Phi$  (i.e. proportion of WT species present in WO),  
308 estimated complement of species “turnover”,  $\gamma$  (i.e. proportion of WO species present in  
309 WT), and “estimated local colonising” species,  $B$  (i.e. number of species not present in  
310 WT but present in WO). As for richness, we compared the mean and standard error  
311 (over 15 re-sampling events) of these estimators between WT and WO types.

312

### 313 2.4.3. Functional trait composition

314 We investigated whether some species attributes were more likely associated with one  
315 type of site and distance to neighbouring woody patch (environmental variables). To  
316 this end, we used a multivariate ordination analysis, RLQ, which is an extension of  
317 coinertia analysis. RLQ performs a double inertia analysis of two arrays (R, here the  
318 environmental array and Q, here the trait array), i.e. measures the adequacy between the  
319 two arrays, with a link expressed by a contingency table (L, here the species array:  
320 matrix of presence-absence of species across sites) (rlq, library ade4 in R, complete  
321 description: see Doledec et al., 1996; for an example: Ribera et al., 2001). The general  
322 significance of the relationship between environmental variables and species traits was  
323 tested using random permutations test ( $n = 999$ ). We also used only species observed in  
324 more than four samples to limit the potentially strong effects of rare species.

325

## 326 **3. Results**

327

328 Across 25 highway verge sites, we observed a total of 85 plant species and 123 different  
329 spider species. For spiders, we captured 8 300 individuals, 7 862 of which were adults  
330 that could be identified to species level. Only eight of the 85 plant species (A.

331 *campestre*, *A. pseudoplatanus*, *B. pendula*, *F. excelsior*, *P. avium*, *P. padus*, *P. spinosa*,  
332 *V. opulus*) were identical to those planted in the immediately neighbouring hedgerow in  
333 WT sites and were excluded from most analyses; note however that including them had  
334 no effect on the following results.

335

### 336 **3.1. Species richness**

337

#### 338 3.1.1. Influence of planted hedgerows on plant communities

339 We observed a significant positive effect of the presence of hedgerows (Table 1) on  
340 plant richness at the local scale (quadrat level,  $n = 125$ ,  $S_{\text{localWT}} = 10.20 \pm 0.46$  SE;  
341  $S_{\text{localWO}} = 8.20 \pm 0.47$  SE,  $F_{1,20} = 5.79$ ,  $p = 0.026$ ). We also found a negative effect of  
342 distance from crop edge ( $F_{1,20} = 5.75$ ,  $p = 0.026$ ), as well as tendencies for effects of  
343 adjacent land use (higher richness with temporary fallow,  $F_{1,20} = 4.15$ ,  $p = 0.055$ ) and  
344 phosphorous rate (negative :  $F_{1,20} = 4.00$ ,  $p = 0.059$ ) on local plant richness. In contrast,  
345 no effect of hedgerows or any environmental factor was detected on plant richness at the  
346 site level ( $n = 25$ ,  $S_{\text{siteWT}} = 25.45 \pm 1.66$  SE;  $S_{\text{siteWO}} = 23.75 \pm 3.12$  SE) and there was no  
347 difference in detection probabilities between the two types of sites. However, at the  
348 section level ( $n = 11$ ), the estimated plant richness was 139.16 over all WT sites ( $S_{\text{typeWT}}$ ,  
349  $\pm 29.20$  SE; detection probability  $dp = 0.47 \pm 12.42$  SE) vs. 87.51 over all WO sites  
350 ( $S_{\text{typeWO}}$ ,  $\pm 19.43$  SE; detection probability  $dp = 0.68 \pm SE = 0.12$ ).

351

#### 352 3.1.2. Influence of planted hedgerows on spider communities

353 We found no effect of the presence of planted hedgerows on spider richness at the site  
354 level. Spider richness was significantly affected by distance to the nearest woodland



355 patch (negative effect,  $F_{1,18} = 5.94$ ,  $p = 0.025$ ), site slope (positive effect,  $F_{1,18} = 4.65$ ,  $p$   
356  $= 0.044$ ) and plant richness (positive effect,  $F_{1,18} = 6.81$ ,  $p = 0.018$ ). There was no  
357 difference in detection probabilities between the two types of sites. At the section level,  
358 ( $n = 11$ ), estimated spider richness was 119.3 over all WT sites ( $S_{\text{typeWT}}$ ,  $\pm 19.6$  SE;  
359 detection probability  $dp = 0.80 \pm 0.10$  SE) and 142.0 over all WO sites ( $S_{\text{typeWO}}$ ,  $\pm 27.15$   
360 SE; detection probability  $dp = 0.64 \pm 0.11$  SE).

361

## 362 **3.2. Community composition**

363

### 364 3.2.1. Spatial autocorrelation

365 Using Mantel tests on the full dataset, we found significant spatial autocorrelation in  
366 plant communities ( $p = 0.001$ ), but not in spider communities ( $p = 0.066$ ). In contrast,  
367 within types of site, spatial autocorrelation was observed in plants (WT:  $p = 0.006$ ; WO:  
368  $p = 0.001$ ) and spiders (WT:  $p = 0.007$ ; WO:  $p = 0.028$ ). However, as WO and WT sites  
369 were randomly distributed in space (Fig. 1), autocorrelation should not interfere with  
370 our main purpose, i.e. to assess the influence of hedgerows on community composition.

371

### 372 3.2.2. The presence of planted hedgerows influence community composition

373 Sites with and without planted hedgerows supported significantly different plant and  
374 spider communities, as shown by CAP (ANOVA-like permutation tests: plants:  $F_{1,22} =$   
375  $2.26$ ,  $p < 0.001$ , inertia explained by hedgerows = 9% and by distance to the nearest  
376 woodland patch = 4%; spiders:  $F_{1,22} = 1.90$ ,  $p = 0.006$ , inertia explained = 7% and 6%,  
377 respectively). This effect of hedgerows on plant and spider community composition was  
378 confirmed by the pCCA analysis (ANOVA-like permutation tests: plants:  $F_{1,22} = 2.13$ ,  $p$

379 < 0.01, inertia explained = 8% and 4% respectively; spiders :  $F_{1,22} = 1.81$ ,  $p < 0.005$ ,  
380 inertia explained = 7% and 5% respectively). This method also identified species that  
381 seemed preferably associated with WT sites. For plants, these were *A. campestre*, *B.*  
382 *pendula*, *Galium mollugo*, *Centaureum erythraea*, *Trifolium repens*, and for spiders,  
383 *Panamomops sulcifrons*, *Maso sundevalli*, *Walckenaera dysderoides*, *Ceratinella*  
384 *brevis*, and *Drassodes cupreus*. In contrast, the spider species *Tegenaria picta*, *Ero*  
385 *furcata*, and *Palliduphantes ericaeus*, were preferably associated with WO sites.

386

### 387 3.2.3. Nested vs. complementary communities

388 We found that the plant community observed in WO sites was a subset of the plant  
389 community in WT sites (Table 1). WO sites did not exhibit any private species: we  
390 found 99% of WO sites species in WT sites, so that all WO species were expected in  
391 WT sites (COMDYN estimator  $B = 0$ ). The reverse was not true: WT exhibited several  
392 private species (14%), with only 86% of WT sites species also observed in WO sites. In  
393 contrast, spider communities appeared complementary. Respectively 83% (WT) and  
394 77% (WO) of spider species were present in the other type of site, so that each type of  
395 sites exhibited private species.

396

### 397 **3.3. Relationship between species attributes and presence of hedgerows**

398

399 For both plants and spiders, there was a significant relationship between species  
400 attributes and environmental variables (Fig. 2; RLQ randtest,  $p = 0.001$  and  $p = 0.001$ ).

401 For the two taxa, the presence/absence of planted hedgerows was associated with the  
402 first axis, which explained respectively 81% and 80% of the total inertia, whereas the

403 distance to the nearest woodland patch tended to be associated with axis 2 (respectively  
404 19% and 20% of inertia). In spontaneous plant species, Raunkier type was influenced by  
405 the presence of hedgerows, with Phanerophytes and Chamaephytes (Rank: P) positively  
406 associated with WT sites (Fig. 2). In contrast, adult height was not significantly  
407 associated with site type. Finally, species with no specific means of dispersal  
408 (barochory, autochory) tended to be more present in WT sites than in WO sites, whereas  
409 animal-dispersed species (zoochory) were more common in WO than in WT sites;  
410 wind-dispersed species were not affected by the presence of hedgerows. In spiders, the  
411 mode of dispersal (ballooning vs. non ballooning) was strongly affected by the presence  
412 of hedgerows: ballooning species were more likely found in WO sites, and non-  
413 ballooning species in WT sites. In addition, web-builders were more common in WT  
414 than in WO sites, and diurnal wandering hunters in WO sites. Finally, larger species  
415 tended to be more common in WT sites than in WO sites.

416

#### 417 **4. Discussion**

418

419 We addressed the effect of the presence of planted hedgerows on neighbouring plant  
420 and spider communities of highway verges. We showed that (1) planted hedgerows  
421 were associated with higher species richness, i.e. a higher  $\alpha$ -diversity ( $\alpha_{WO} < \alpha_{WT}$ ), for  
422 plants, but not for spiders. This was due to the fact that plant communities in sites  
423 without hedgerows (WO) were a subset of communities in sites with hedgerows (WT),  
424 whereas spider communities in WO sites were complementary to communities in WT  
425 sites. Below, we argue that such differences are attributable to different ecological

426 characteristics of species, which is supported by the distribution of functional traits, and  
427 derive the consequences in terms of roadside management.

428

#### 429 **4.1. The influence of propagule source vs. local conditions on plant and spider** 430 **community composition**

431

432 Following the environmental filter model (Keddy, 1992), two non-exclusive hypotheses  
433 may explain the effect of hedgerows on community composition, depending on  
434 taxonomic group. Hedgerows could either be a source of dispersers (“source”  
435 hypothesis) or could provide microclimatic conditions that affect the presence of some  
436 species due to their environmental requirements (“local conditions” hypothesis).

437

##### 438 4.1.1. “Source” hypothesis and the effect of hedgerows on plant communities

439 The dispersal of some species from neighbouring hedgerows into a suboptimal habitat  
440 (grassland) would affect community composition in the suboptimal habitat. This  
441 phenomenon, known as the edge effect, has been observed in several taxa: contrasted  
442 adjacent habitats supporting distinct species assemblages often mix over a few meters  
443 on either side of the interface (see for example Bedford and Usher, 1994). For example,  
444 Downie et al. (1996) studying spider community composition across a plantation-  
445 pasture ecotone, found that most of the community (70% of species) was formed by the  
446 overlap of both pasture and plantation species encroaching across the interface through  
447 dispersal or foraging processes. Interface-specialist species were present but contributed  
448 less to the interface community content than the other species categories (23% of  
449 individuals and 25% of species).

450 Under the “source” hypothesis, communities are therefore expected to exhibit  
451 higher species richness in WT vs. WO sites. This was the case for plant communities  
452 ( $\alpha_{wo} < \alpha_{wt}$ ), at the quadrat scale, but not at the site level. Another prediction of the  
453 source hypothesis is that WO communities should be a subset of WT communities. This  
454 was true again for plants: in WT sites, we observed ‘private’ species which were not  
455 supported by WO sites, whereas the reverse was not true. Finally, in the case of a higher  
456 effect of dispersal vs. local conditions on species distribution within similar habitats, a  
457 distance effect can be expected independently of presence/absence of hedgerows. In  
458 plant communities, spatial autocorrelation was in fact detected regardless of the  
459 presence of hedgerows. These three observations converge to suggest that dispersal is a  
460 major driver of community composition in this group.

461 The distribution of plant functional traits is consistent with this role of  
462 dispersal. We found that Phanerophytes and Chamaephytes were more represented in  
463 WT sites, corresponding mostly to the occurrence of the following species: *A.*  
464 *campestre*, *B. pendula*, *C. scoparius*, *F. excelsior* and *P. spinosa*. These species  
465 observed in the herbaceous strata most likely originated from seed-bearer trees that  
466 grow spontaneously within the otherwise planted hedgerow. Thus, hedgerows offer a  
467 sustainable habitat where particular species, and notably woody plants, can grow and  
468 reproduce, whereas such species cannot reach maturity within WO sites, due to frequent  
469 mowing. The fact that zoochory was more represented in WO sites is however not  
470 easily interpreted in this framework.

471

472 4.1.2. Local conditions hypothesis and the effect of hedgerows on spider communities

473 Microclimatic conditions (influenced, among others, by the neighbouring hedgerows)  
474 might be responsible for observed community composition, especially in spiders. For  
475 example, spider richness within sites was positively correlated with local plant richness;  
476 plant community composition and consequently the architecture of habitat are well-  
477 known predictors of spider community composition (e.g. Beals, 2006). Moreover,  
478 woody hedgerows are known to provide particular microclimate conditions in their  
479 neighbourhood (Forman and Baudry, 1984) and most authors agree that those are a  
480 major factor controlling spider distribution (see Marc et al., 1999). For example,  
481 Downie et al. (1996), examining spider community composition, proposed that close to  
482 an interface, the role of vegetation structure is low compared to other factors, including  
483 changes in microclimate. Finally, perennial non-crop habitats are known to be used by  
484 spiders as overwintering sites and refuges (Oberg et al. 2008). Grassland habitats  
485 located next to hedgerows maybe therefore provide better conditions than sites without  
486 hedgerows for overwintering, with direct consequences on spider community  
487 composition. Thus, neighbouring hedgerows may have an environmental filter effect,  
488 which leads to the selection of species with particular ecological requirements.

489 Under the 'local conditions' hypothesis, we expect that sites with and without  
490 planted hedgerows support different species. This was the case with spiders: WT and  
491 WO communities were complementary instead of nested, thus yielding similar species  
492 richness with or without hedgerows. This implies that some spider species occurring in  
493 WO sites were absent from planted sites. As the two types of sites offered the same  
494 habitat (grassland with similar percentage cover and dominance of graminoids), the  
495 observed differences are likely be explained by specific microclimatic conditions  
496 created by the neighbouring hedgerows. Finally, the role of hedgerow-induced local

497 conditions in spider community composition is also supported by patterns of spatial  
498 autocorrelation: the dominant impact of neighbouring hedgerows on spider community  
499 composition removed spatial autocorrelation patterns across all sites, whereas spatial  
500 autocorrelation remained significant within WT and WO sites.

501         Despite the significant role of local conditions, spider community composition is  
502 also likely affected by dispersal. For example, spider richness within sites was  
503 negatively correlated with the distance to the nearest woodland patch. Such effect of  
504 distance is well known from the theory of island biogeography (Mac-Arthur and  
505 Wilson, 1967) and consistent with results from other field studies: spider species  
506 richness in wheat fields has been shown to increase with the proportion of non-crop  
507 habitats in the surrounding landscape (Schmidt and Tschardtke, 2005; see also Bianchi  
508 et al., 2006 for other groups).

509         Although the observed distribution of species between site types suggests a  
510 major role of local conditions, trait analyses outlined that aerial dispersal capabilities are  
511 an important factor explaining the interaction between the presence of hedgerows and  
512 spider community composition: species exhibiting ballooning behaviour were more  
513 represented in WO sites and the reverse was true for non-ballooning species. This result  
514 is comparable to that of Bonte et al. (2004), who found that the distribution of 29 spider  
515 species inhabiting fragmented grey dunes depended on their aerial dispersal potential, as  
516 well as on the interaction between patch connectivity and area. Ballooning is thought to  
517 be more common in species from unstable and ephemeral habitats (Bonte et al., 2003;  
518 Bell et al., 2005); moreover, high dispersal capabilities are generally dominant in  
519 disturbed habitats (e.g. Nyffeler and Sunderland, 2003 for spiders; Ribera et al., 2001  
520 for beetles), allowing the maintenance of populations in ephemeral agricultural

521 ecosystems (Weyman et al., 1995). We therefore suggest that the significantly higher  
522 proportion of ballooning species in WO sites may be due to the absence of a  
523 neighbouring hedgerow, i.e. of a stable habitat. In contrast, species that do not exhibit  
524 this behaviour should be more common in WT sites because hedgerow likely provides  
525 more stable conditions, both in terms of management (no mowing) and microclimatic  
526 variables. Hedgerows would thus be overwintering refuges (see also: Pywell et al.,  
527 2005), from which species colonise the surrounding habitats in spring. Sites without  
528 planted hedgerows, which receive a cut, generally in autumn, would be poor refuges  
529 that are partly recolonised from more distant sources in spring. Hence, they support a  
530 higher proportion of species exhibiting large dispersal capabilities, i.e. “ballooners”.  
531 Dispersal and local conditions therefore appear to drive spider community composition  
532 at different temporal scales.

533         Finally, note that taxonomic differences in composition of spider communities  
534 were also associated with differences in hunting strategies: web-builder were more  
535 represented in WT sites possibly due to the greatest heterogeneity of support provided  
536 by the presence of neighbouring hedgerow and the higher richness of plant whereas  
537 diurnal wandering hunter were more abundant in WO sites.

538

#### 539 4.1.3. Conclusions

540 Planting generally increased the  $\gamma$ -diversity of highway verges, via an increase in both  
541 taxonomic and functional trait  $\beta$ -diversity between sites with and without hedgerows,  
542 but this positive effect was different in the two taxa considered. For plants, it was  
543 mostly due to an increase in taxonomic  $\alpha$ -diversity within sites with hedgerows,  
544 whereas for spiders it was related to a direct increase of taxonomic  $\beta$ -diversity between



545 the two types of sites. Note that this partition of diversity is likely to vary through time  
546 as well (e.g. due to overwintering dispersal patterns in spiders), with different temporal  
547 scales in the two taxa, but this was not the focus of the present study. From a theoretical  
548 viewpoint, the fact that the partitioning of diversity between  $\alpha$  and  $\beta$  diversity differed  
549 between the two taxa is probably due to the different spatial and temporal scales at  
550 which the heterogeneity of interactions between organisms and their environment can  
551 be found (see Loreau 2000). According to Loreau (2000), “the challenge now is to  
552 understand the relationship between  $\alpha$  and  $\beta$  diversity at multiple scales, and the  
553 processes that determine it”. In our opinion, comparing the partitioning of diversity  
554 among several taxa and functional groups at different spatial scales is therefore a  
555 promising way to explore the processes that control the distribution of biodiversity.

556

#### 557 **4.2. Consequences for the management of highway verges**

558

559 Our findings have direct consequences for the management and conservation of  
560 highway verges in agricultural landscapes, where most biodiversity is aggregated in  
561 more stable non-crop habitats and field edges (Bianchi et al., 2006). Increasing the  
562 botanical and structural heterogeneity of hedgerows by planting is likely to have  
563 significant beneficial effects on native flora and fauna (Maudsley, 2000). Obviously,  
564 plantation increases species richness both directly (introduction of new species) and  
565 indirectly, by providing particular environmental conditions within woody hedgerows  
566 and thus contributing to the maintenance of some species (e.g. woody plants) in  
567 landscapes where they are poorly represented. But our results suggest that planted

568 hedges may also affect community composition of nearby herbaceous strata, enhancing  
569 both taxonomic and functional trait  $\gamma$ -diversity along highway verges.

570         In practice, our results highlight the importance of creating a mosaic of planted  
571 sites alternating with non planted sites in highly disturbed landscapes. Of course,  
572 “natural” dynamics could yield the same results and probably a greater biodiversity as  
573 demonstrated by Boutin et al. (2002), who compared natural vs. planted hedgerows.  
574 However, in the case of highway verges, natural hedgerows would only arise by  
575 recolonisation of the newly created habitats, which involves successional dynamics with  
576 intermediate states, and the presence of pioneer species often considered as weeds. In  
577 the context of intensive agricultural landscape, where farmers wish to limit weeds in  
578 field boundaries as much as possible, planting trees along highway verges appears as a  
579 compromise between biodiversity conservation and anthropogenic pressures. The issues  
580 of the identity of planted species (which are usually exotic species), of management of  
581 hedgerows (Croxtton et al., 2004) and of their spatial arrangement along highway verges  
582 (to provide effective corridors) remain to be addressed. Given the urge to conserve  
583 biodiversity, especially in the context of climate change, road practitioners should  
584 consider plantation not only for esthetical reasons but also as a possibility to increase  
585 the role of verges as a refuge and, consequently, landscape connectivity.

586

587

---

588 **Acknowledgements:**

589

590 This work was financially supported by the Direction of Construction of Cofiroute.

591 Emmanuelle Porcher was partly funded by Agence Nationale de la Recherche (ANR

592 grant #2006-JCJC-0032). We are grateful to *Cofiroute* for providing access to Highway

593 verges and to *Vegetude* for information about revegetation. We thank Coralie

594 Dedeystere for assistance in the field, Sophie Gachet for help with plants traits,

595 Sandrine Pavoine for precious help with statistical analysis and Didier Bas and Vincent

596 Devictor for software facilities (COMDYN).

597

598

599    **REFERENCES**

600

601    Anderson, M.J., Willis, T.J., 2003. Canonical analysis of principal coordinates: A useful  
602       method of constrained ordination for ecology. *Ecology* 84, 511-525.

603    Baudry, J., Bunce, R.G.H., Burel, F., 2000. Hedgerows: An international perspective on  
604       their origin, function and management. *Journal of Environmental Management* 60,  
605       7-22.

606    Beals, M.L., 2006. Understanding community structure: a data-driven multivariate  
607       approach. *Oecologia* 150, 484-495.

608    Bedford, S.E., Usher, M.B., 1994. Distribution of arthropod species across the margins  
609       of farm woodlands. *Agriculture Ecosystems and Environment* 48, 295-305.

610    Bell, J.R., Bohan, D.A., Shaw, E.M., Weyman, G.S., 2005. Ballooning dispersal using  
611       silk: world fauna, phylogenies, genetics and models. *Bulletin of Entomological*  
612       *Research* 95, 69-114.

613    Bianchi, F.J.J.A., Booij, C.J.H., Tscharrntke, T., 2006. Sustainable pest regulation in  
614       agricultural landscapes: a review on landscape composition, biodiversity and natural  
615       pest control. *Proceedings of the Royal Society B: Biological Sciences* 273, 1715-  
616       1727.

617    Bochet, E., Garcia-Fayos, P., Tormo, J., 2007. Road slope revegetation in semiarid  
618       mediterranean environments. Part I: Seed dispersal and spontaneous colonization.  
619       *Restoration Ecology* 15, 88-96.

620    Bonte, D., Vandebroecke, N., Lens, L., Maelfait, J.P., 2003. Low propensity for aerial  
621       dispersal in specialist spiders from fragmented landscapes. *Proceedings of the Royal*  
622       *Society of London Series B-Biological Sciences* 270, 1601-1607.

623 Bonte, D., Baert, L., Lens, L., Maelfait, J.P., 2004. Effects of aerial dispersal, habitat  
624 specialisation, and landscape structure on spider distribution across fragmented grey  
625 dunes. *Ecography* 27, 343-349.

626 Boulinier, T., Nichols, J.D., Hines, J.E., Sauer, J.R., Flather, C.H., Pollock, K.H., 1998.  
627 Higher temporal variability of forest breeding bird communities in fragmented  
628 landscapes. *Proceedings of the National Academy of Sciences of the United States*  
629 *of America* 95, 7497-7501.

630 Boutin, C., Jobin, B., Belanger, L., Choiniere, L., 2002. Plant diversity in three types of  
631 hedgerows adjacent to cropfields. *Biodiversity and Conservation* 11, 1-25.

632 Brown, G.P., Phillips, B.L., Webb, J.K., Shine, R., 2006. Toad on the road: Use of roads  
633 as dispersal corridors by cane toads (*Bufo marinus*) at an invasion front in tropical  
634 Australia. *Biological Conservation* 133, 88-94.

635 Burel, F., 1996. Hedgerows and their role in agricultural landscapes. *Critical Reviews in*  
636 *Plant Sciences* 15, 169 - 190.

637 Burnham, K.P., Overton, W.S., 1979. Robust estimation of population-size when  
638 capture probabilities vary among animals. *Ecology* 60, 927-936.

639 Corbit, M., Marks, P.L., Gardescu, S., 1999. Hedgerows as habitat corridors for forest  
640 herbs in central New York, USA. *Journal of Ecology* 87, 220-232.

641 Croxton, P.J., Franssen, W., Myhill, D.G., Sparks, T.H., 2004. The restoration of  
642 neglected hedges: a comparison of management treatments. *Biological Conservation*  
643 117, 19-23.

644 Devictor, V., Julliard, R., Couvet, D., Lee, A., Jiguet, F., 2007. Functional  
645 Homogenization Effect of Urbanization on Bird Communities. *Conservation*  
646 *Biology* 21, 741-751.

647 Davies, Z., Pullin, A., 2007. Are hedgerows effective corridors between fragments of  
648 woodland habitat? An evidence-based approach. *Landscape Ecology* 22, 333-351.

649 Doherty, P.F., Sorci, G., Royle, J.A., Hines, J.E., Nichols, J.D., Boulinier, T., 2003.  
650 Sexual selection affects local extinction and turnover in bird communities.  
651 *Proceedings of the National Academy of Sciences of the United States of America*  
652 100, 5858-5862.

653 Doledec, S., Chessel, D., ter Braak, C.J.F., Champely, S., 1996. Matching species traits  
654 to environmental variables: A new three-table ordination method. *Environmental*  
655 *and Ecological Statistics* 3, 143-166.

656 Downie, I.S., Coulson, J.C., Butterfield, J.E.L., 1996. Distribution and dynamics of  
657 surface-dwelling spiders across a pasture-plantation ecotone. *Ecography* 19, 29-40.

658 Entling, W., Schmidt, M.H., Bacher, S., Brandl, R., Nentwig, W., 2007. Niche  
659 properties of Central European spiders: shading, moisture and the evolution of the  
660 habitat niche. *Global Ecology and Biogeography* 16, 440-448.

661 ESRI., 2000. ArcView 3.2, Redlands, California

662 Forman, R.T.T., Baudry, J., 1984. Hedgerows and hedgerow networks in landscape  
663 ecology. *Environmental Management* 8, 495-510.

664 Griffiths, G.J.K., Winder, L., Holland, J.M., Thomas, C.F.G., Williams, E., 2007. The  
665 representation and functional composition of carabid and staphylinid beetles in  
666 different field boundary types at a farm-scale. *Biological Conservation* 135, 145-  
667 152.

668 Hansen, M.J., Clevenger, A.P., 2005. The influence of disturbance and habitat on the  
669 presence of non-native plant species along transport corridors. *Biological*  
670 *Conservation* 125, 249-259.

671 Hines, J.E., Boulinier, T., Nichols, J.D., Sauer, J.R., Pollock, K.H., 1999. COMDYN:  
672 software to study the dynamics of animal communities using a capture-recapture  
673 approach. *Bird Study* 46, 209-217.  
674 <http://www.mbr-pwrc.usgs.gov/software/comdyn.html>

675 Jantunen, J., Saarinen, K., Valtonen, A., Saarnio, S., 2007. Flowering and seed  
676 production success along roads with different mowing regimes. *Applied Vegetation*  
677 *Science* 10, 285-292.

678 Keddy, P.A., 1992. Assembly and Response Rules - 2 Goals for Predictive Community  
679 Ecology. *Journal of Vegetation Science* 3, 157-164.

680 Kerbiriou, C., Le Viol, I., Jiguet, F., Julliard, R., 2008. The impact of human  
681 frequentation on coastal vegetation in a Biosphere Reserve. *Journal of*  
682 *Environmental Management*. In press: doi: 10.1016/j.envman.2007.03.034

683 Leps, J., Dolezal, J., Bezemer, T.M., Brown, V.K., Hedlund, K., Igual, A.M., Jorgensen,  
684 H.B., Lawson, C.S., Mortimer, S.R., Peix Geldart, A., Rodriguez Barrueco, C.,  
685 Santa Regina, I., Smilauer, P., van der Putten, W.H., 2007. Long-term effectiveness  
686 of sowing high and low diversity seed mixtures to enhance plant community  
687 development on ex-arable fields. *Applied Vegetation Science* 10, 97-110.

688 Lekve, K., Boulinier, T., Stenseth, N.C., Gjosaeter, J., Fromentin, J.M., Hines, J.E.,  
689 Nichols, J.D., 2002. Spatio-temporal dynamics of species richness in coastal fish  
690 communities. *Proceedings of the Royal Society of London Series B-Biological*  
691 *Sciences* 269, 1781-1789.

692 Loreau, M., 2000. Are communities saturated? On the relationship between alpha, beta  
693 and gamma diversity. *Ecology Letters* 3, 73-76.

694 Lovell, S., Hamer, M., Slotow, R., Herbert, D., 2007. Assessment of congruency across  
695 invertebrate taxa and taxonomic levels to identify potential surrogates. *Biological*  
696 *Conservation* 139, 113-125.

697 Marc, P., Canard, A., Ysnel, F., 1999. Spiders (Araneae) useful for pest limitation and  
698 bioindication. *Agriculture Ecosystems and Environment* 74, 229-273.

699 Matesanz, S., Valladares, F., Tena, D., Costa-Tenorio, M., Bote, D., 2006. Early  
700 dynamics of plant communities on revegetated motorway slopes from southern  
701 Spain: Is hydroseeding always needed? *Restoration Ecology* 14, 297-307.

702 Maudsley, M.J., 2000. A review of the ecology and conservation of hedgerow  
703 invertebrates in Britain. *Journal of Environmental Management* 60, 65-76.

704 MacArthur R. H., Wilson. E.O., 1967. *The Theory of Island Biogeography*. Princeton  
705 University Press, Princeton, New Jersey.

706 McCollin, D., Jackson, J.I., Bunce, R.G.H., Barr, C.J., Stuart, R., 2000. Hedgerows as  
707 habitat for woodland plants. *Journal of Environmental Management* 60, 77-90.

708 Melbourne, B.A., 1999. Bias in the effect of habitat structure on pitfall traps: An  
709 experimental evaluation. *Australian Journal of Ecology* 24, 228-239.

710 Meunier, F.D., Verheyden, C., Jouventin, P., 2000. Use of roadsides by diurnal raptors  
711 in agricultural landscapes. *Biological Conservation* 92, 291-298.

712 Nichols, J.D., Boulinier, T., Hines, J.E., Pollock, K.H., Sauer, J.R., 1998. Inference  
713 methods for spatial variation in species richness and community composition when  
714 not all species are detected. *Conservation Biology* 12, 1390-1398.

715 Nyffeler, M., Sunderland, K.D., 2003. Composition, abundance and pest control  
716 potential of spider communities in agroecosystems: a comparison of European and  
717 US studies. *Agriculture Ecosystems and Environment* 95, 579-612.



718 Oberg, S., Mayr, S., Dauber, J., 2008. Landscape effects on recolonisation patterns of  
719 spiders in arable fields. *Agriculture Ecosystems and Environment* 123, 211-218.

720 O'Farrell, P.J., Milton, S.J., 2006. Road verge and rangeland plant communities in the  
721 southern Karoo: exploring what influences diversity, dominance and cover.  
722 *Biodiversity and Conservation* 15, 921-938.

723 Pearman, P.B., Weber, D., 2007. Common species determine richness patterns in  
724 biodiversity indicator taxa. *Biological Conservation* 138, 109-119.

725 Platnick, N.I., 2007. *The World Spider catalog, Version 8*. American Museum Of  
726 Natural History, New York, N.Y.  
727 <http://research.amnh.org/entomology/spiders/catalog/index.html>.

728 Pywell, R.F., James, K.L., Herbert, I., Meek, W.R., Carvell, C., Bell, D., Sparks, T.H.,  
729 2005. Determinants of overwintering habitat quality for beetles and spiders on  
730 arable farmland. *Biological Conservation* 123, 79-90.

731 R Development Core Team, 2007. *R: a language and environment for statistical*  
732 *computing*. R Foundation for Statistical Computing, Vienna, Austria.  
733 <http://www.r-project.org/>

734 Ribera, I., Doledec, S., Downie, I.S., Foster, G.N., 2001. Effect of land disturbance and  
735 stress on species traits of ground beetle assemblages. *Ecology* 82, 1112-1129.

736 Ries, L., Debinski, D.M., Wieland, M.L., 2001. Conservation Value of Roadside Prairie  
737 Restoration to Butterfly Communities. *Conservation Biology* 15, 401-411.

738 Roberts, M., 1985. *The Spiders of Great Britain and Ireland, Vol. 1. Atypidae to*  
739 *Theridosomatidae*. Harley Books, Colchester, Essex.

740 Roberts, M., 1987. *The Spiders of Great Britain and Ireland, Vol. 2. Linyphiidae*.  
741 Harley Books, Colchester, Essex.

742 Roy, V., Blois, S.d., 2006. Using functional traits to assess the role of hedgerow  
743 corridors as environmental filters for forest herbs. *Biological Conservation* 130,  
744 592-603.

745 Roy, V., de Blois, S., 2008. Evaluating hedgerow corridors for the conservation of  
746 native forest herb diversity. *Biological Conservation* 141, 298-307.

747 Schaffers, A.P., 2002. Soil, biomass, and management of semi-natural vegetation – Part  
748 II. Factors controlling species diversity. *Plant Ecology* 158, 247-268.

749 Schmidt, M.H., Tschardtke, T., 2005. The role of perennial habitats for Central  
750 European farmland spiders. *Agriculture, Ecosystems and Environment* 105, 235-  
751 242.

752 Selmi, S., Boulinier, T., 2003. Does time of season influence bird species number  
753 determined from point-count data? A capture-recapture approach. *Journal of Field*  
754 *Ornithology* 74, 349-356.

755 Sitzia, T., 2007. Hedgerows as corridors for woodland plants: a test on the Po Plain,  
756 northern Italy. *Plant Ecology* 188, 235-252.

757 Smart, S.M., Marrs, R.H., Le Duc, M.G., Thompson, K.E.N., Bunce, R.G.H., Firbank,  
758 L.G., Rossall, M.J., 2006. Spatial relationships between intensive land cover and  
759 residual plant species diversity in temperate farmed landscapes. *Journal of Applied*  
760 *Ecology* 43, 1128-1137.

761 Spooner, P.G., Lunt, I.D., Briggs, S.V., Freudenberger, D., 2004. Effects of soil  
762 disturbance from roadworks on roadside shrubs in a fragmented agricultural  
763 landscape. *Biological Conservation* 117, 393-406.

764 ter Braak, C.J.F., 1986. Canonical Correspondence-Analysis - a New Eigenvector  
765 Technique for Multivariate Direct Gradient Analysis. *Ecology* 67, 1167-1179.

766 Tikka, P.M., Högmander, H., Koski, P.S., 2001. Road and railway verges serve as  
767 dispersal corridors for grassland plants. *Landscape Ecology* 16, 659-666.

768 Trombulak, S.C., Frissell, C.A., 2000. Review of ecological effects of roads on  
769 terrestrial and aquatic communities. *Conservation Biology* 14, 18-30.

770 Watts, R.D., Compton, R.W., McCammon, J.H., Rich, C.L., Wright, S.M., Owens, T.,  
771 Ouren, D.S., 2007. Roadless Space of the Conterminous United States. *Science* 316,  
772 736-738.

773 Way, J.M., 1977. Roadside verges and conservation in Britain: A review. *Biological*  
774 *Conservation* 12, 65-74.

775 Weyman, G.S., Jepson, P.C., Sunderland, K.D., 1995. Do seasonal changes in numbers  
776 of aerially dispersing spiders reflect population density on the ground or variation in  
777 ballooning motivation? *Oecologia* 101, 487-493.

778

779 **Captions**

780

781 **Table 1** - Changes in species composition in the herbaceous strata of highway verges in  
782 sites without planted hedgerows (WO) vs. sites with planted hedgerows (WT).

783

784 **Fig. 1**- Location of the 25 study sites.

785

786 **Fig. 2** - RLQ analysis ordination biplot illustrating relationships among environmental  
787 parameters and plant species attributes (A) and spider species attributes (B).

788

789 **Table 1** - Change in species composition in the herbaceous strata of highway verges in  
790 sites without planted hedgerows (WO) vs. sites with planted hedgerows (WT).


791

	$\Phi$	$\gamma$
Plants	$0.86 \pm 0.11$	$0.99 \pm 0.10$
Spiders	$0.77 \pm 0.09$	$0.83 \pm 0.08$

792  $\Phi$  is the proportion of WT sites species still present in WO sites and  $\gamma$  is the proportion  
793 of WO sites species still present in WT sites.

794

795 **Fig. 1-** Location of the 25 study sites.




809

810 ▲: sites without hedgerows (WO), ○: sites with planted hedgerows (WT); line: A11  
811 highway; grey patches: woodland patches within a 2 000m radius buffer.

812

813 **Fig. 2** - RLQ analysis ordination biplot illustrating relationships between environmental  
 814 parameters and plant species attributes (A) and spider species attributes (B).



831 The analysis is based on presence/absence of species with known attributes that were  
 832 observed in more than four samples (46 plant species and 56 spider species).

833 1) **Environmental parameters.** WT: sites with planted hedgerows, WO: sites without  
 834 hedgerows; dist: distance to the distance to the nearest woodland patch. 2) **Plant traits**  
 835 **(A).** rank: raunkier type: H: Hemicryptophytes and Geophytes (28 species), P:  
 836 Phanerophytes and Chamaephytes (6), T: Therophytes (12); heigh: maximum adult  
 837 height: S: small (0 to 45 cm: 9), M: medium (46 to 100 cm: 19), H: high (100 and more:  
 838 18); seed: Seed dispersal: A : anemochory (30), B: barochory and autochory (9), Z:  
 839 zoochory (7). 2) **Spider traits (B).** ballo: ballooning behaviour: Y: Yes (30), N: No  
 840 (26); size; hunt: hunting strategies: W: diurnal wandering (18); N: nocturnal wandering  
 841 (10); W: web-builders (21); A: ambush hunters (7).