# **Supplementary Information:**

Direct evidence of the composition of a Neandertal social group: the hominin footprints at Le Rozel (Normandy, France)

# Jérémy Duveau<sup>1,\*</sup>, Gilles Berillon<sup>1</sup>, Christine Verna<sup>1</sup>, Gilles Laisné<sup>2</sup>, and Dominique Cliquet<sup>2,3,4</sup>

<sup>1</sup>UMR 7194 HNHP, Centre national de la recherche scientifique, Muséum national d'Histoire naturelle, Université Perpignan Via Domitia, 75013 Paris, France.

<sup>2</sup>Projet Collectif de Recherche « Les premiers Hommes en Normandie », Ministère de la Culture, France

<sup>3</sup>Service Régional de l'Archéologie, Direction Régionale des Affaires Culturelles Normandie, Ministère de la Culture, 14052 Caen Cedex 4, France.

<sup>4</sup> UMR 6566 CReAAH, Centre national de la recherche scientifique, Université de Rennes 1, 35042 Rennes, France

<sup>\*</sup>Email: <u>jeremy.duveau@edu.mnhn.fr</u> Adress: Musée de l'Homme, 17 Place du Trocadéro, 75116 Paris, France.

## The hominin footprints sites:



Fig. S1. Geographical distribution of pre-Holocene sites with potential hominin footprints

Legends: 1-Trachilos (Greece), 2-Laetoli (Tanzania), 3-Ileret (Kenya), 4-Koobi Fora (Kenya), 5-Happisburgh (Great-Britain), 6-Gombore II-2 (Ethiopia), 7-Terra Amata (France), 8-Roccamonfina (Italia), 9-Biache-Saint-Vaast (France), 10- Still Bay (South Africa), 11-Theopetra (Greece), 12-Nahoon (South Africa), 13-Langebaan (South Africa), 14-Vârtop (Romania), 15-Brenton-on-Sea (South Africa), 16-Valsequillo (Mexico), 17-Chauvet (France), 18-Ciur-Ibzuc (Romania), 19-Catalan Bay (Gibraltar), 20-Cussac (France), 21-Jeju Island (South Korea), 22-Pech-Merle (France), 23-Willandra Lakes (Australia), 24-Tibetan plateau (China), 25-Engare Sero (Tanzania), 26-Lascaux (France), 27-Ojo Guareña (Spain), 28-White Sands National Monument (USA), 29-Monte Verde (Chile), 30-Tuc d'Adoubert (France), 31-Calvert Island (Canada), 32-Niaux (France), 33-Tana della Basura (Italia), 34-Fontanet (France), 35-Pehuen-Co (Argentina), 36-Lake Bogoria (Kenya).

	Site	Ages	Outside/Cave	Substrate	Associated taxon	Summary description	References (e.g.)
1	Trachilos (Greece)	5.7 Ma Upper Miocene	Outside	Sand	Unknown	About 50 tracks including 2 trackways. They have a triangular shape and are relatively short. The hallux seems adducted. The identification as footprints and the association with hominins are discussed.	1-3
2	Laetoli (Tanzania)	3.7 Ma Pliocene	Outside	Tuff	Au. afarensis	4 trackways and an isolated footprint made by juveniles and adults with different statures. The footprints are broad with an abducted hallux. They reflect the first direct evidence of human bipedalism.	4-7
3	lleret (Kenya)	1.5 Ma Lower Pleistocene	Outside	Sand / Silt	H. erectus / P. boisei	About one hundred footprints including isolated footprints and more than 20 trackways. The footprint morphology differs from the Laetoli footprints. They reflect a human foot function.	8-12
4	Koobi Fora (Kenya)	1.4 Ma Lower Pleistocene	Outside	Sand / Silt	H. erectus	A trackway of 7 weakly preserved footprints. Footprint morphology differs from the Laetoli tracks and is closer to anatomically modern footprints.	8, 13-14
5	Happisburgh (Great-Britain)	1-0.8 Ma Lower Pleistocene	Outside	Sand / Silt	H. antecessor	152 footprints. Most of them are isolated. Presence of possible trackways. Only one footprint presents clear toe impressions.	15
6	Gombore II-2 (Ethiopia)	700 ka Middle Pleistocene	Outside	Sand / Silt	H. heidelbergensis	11 footprints including a trackway of 2 footprints. Their morphology is consistent with the other footprints attributed to <i>Homo</i> genus. These footprints were made by adults and juveniles. Associated with animal tracks.	16
7	Terra Amata (France)	380 ka Middle Pleistocene	Outside	Sand	H. heidelbergensis	Only one footprint reflecting a slide of the foot. The impression of the hallux is visible. Ages are discussed.	17-18
8	Roccamonfina (Italia)	349 ka Middle Pleistocene	Outside	Tuff	H. heidelbergensis	3 trackways made going down a slope. The footprints are short and wide. They reflect few anatomical details.	19-22
9	Biache-Saint-Vaast (France)	236 ka Middle Pleistocene	Outside	Silt	H. neanderthalensis	1 potential footprint surrounded by animal tracks. Not studied precisely.	23-25
10	Still Bay (South Africa)	140-91 ka Middle / Upper Pleistocene	Outside	Aeolianite	H. sapiens	A trackway of 4 potential footprints on a slab. The second and the third footprints are the most visible. Toe impressions are not visible. This trackway could have been made by a shod individual. The identification as hominin footprints is not consensual (they could be animal tracks).	26-27
11	Theopetra (Greece)	130 ka Middle Pleistocene	Cave	Clay	H. neanderthalensis	4 footprints made by young individuals (2-4 years old). One of the footprints is interpreted as made by a shod foot.	28-29
12	Nahoon (South Africa)	124 ka Upper Pleistocene	Outside	Aeolianite	H. sapiens	A trackway of 3 anatomically modern footprints. Toe impressions are visible.	27, 30-31
13	Langebaan (South Africa)	117 ka Upper Pleistocene	Outside	Aeolianite	H. sapiens	A trackway of 3 anatomically modern footprints. They are less preserved than at Nahoon.	27, 30, 32
14	Vârtop (Romania)	97-62 ka Upper Pleistocene	Cave	Calcareous mud	H. neanderthalensis	3 isolated footprints probably made by a single individual. The most complete footprint shows a "large space between the hallux and the second toe".	33-35
15	Brenton-on-Sea (South Africa)	90 ka Upper Pleistocene	Cave	Aeolianite	H. sapiens	40 footprints from 2 different layers made by going down a dune. Some footprints are part of trackways.	27, 36
16	Valsequillo (Mexico)	40 ka (?) Upper Pleistocene	Outside	Tuff	H. sapiens	Several possible trackways and isolated footprints. Initially described as one of the first human occurrences in America. The identification as footprints and their dating are highly contested. They would more likely be metal tool tracks made during quarrying.	37-41
17	Chauvet (France)	37-28 ka (?) Upper Pleistocene	Cave	Clay	H. sapiens	A trackway of about 20 footprints made by a juvenile. Dating is discussed.	42-46

# Table S1 (2/2). Pre-Holocene sites with potential hominin footprints

	Site	Ages	Outside/Cave	Substrate	Associated taxon	Summary description	References (e.g.)
18	Ciur-Izbuc (Romania)	36.5-29 ka Upper Pleistocene	Cave	Clay	H. sapiens	Several hundred footprints made by different individuals. Their taxonomic attribution and their ages are discussed.	35, 47-48
19	Catalan Bay (Gibraltar)	28 ka Upper Pleistocene	Outside	Aeolianite	H. neanderthalensis	A single potential footprint representing the descent of a dune. It is poorly preserved but some toe impressions may be deduced. Associated with several animal tracks.	49
20	Cussac (France)	29 ka Upper Pleistocene	Cave	Clay	H. sapiens	Several footprints including 3 trackways. Associated with handprints.	50-51
21	Jeju Island (South Korea)	25-3.7 ka (?) Upper Pleistocene / Holocene	Outside	Tuff	H. sapiens	505 footprints, including at least 9 trackways, made by juveniles and adults. Dating is discussed.	52-56
22	Pech-Merle (France)	25-15 ka (?) Upper Pleistocene.	Cave	Clay	H. sapiens	12 footprints made by juveniles and adults. Some tracks have been identified as those of a walking stick. Dating is uncertain.	57-59
23	Willandra Lakes (Australia)	23-19 ka Upper Pleistocene	Outside	Silt / Clay	H. sapiens	563 footprints including 23 trackways. Speed estimates indicate that some individuals were very fast.	60-62
24	Tibetan plateau (China)	21 ka Upper Pleistocene	Outside	Travertine	H. sapiens	11 footprints associated with handprints. They represent one of the first human occurrences in Tibet.	63-64
25	Engare Sero (Tanzania)	19.1-5.8 ka Upper Pleistocene / Holocene	Outside	Tuff	H. sapiens	More than 400 footprints including 24 trackways and isolated footprints.	65-68
26	Lascaux (France)	17 ka Upper Pleistocene	Cave	Clay	H. sapiens	Footprints made by adolescents. Ages are discussed.	69-70
27	Ojo Guareña (Spain)	15.6 ka Upper Pleistocene	Cave	Clay	H. sapiens	More than 1 000 footprints including trackways and isolated footprints made by at least 8 individuals. Discussed as an exploration of the cave by prehistoric groups.	71
28	White Sands National Monument (USA)	15.6-10.0 ka Upper Pleistocene	Outside	Sand / Mud	H. sapiens	Nearly 100 footprints including trackways and isolated footprints associated with sloth tracks. They could reflect hunting behaviors.	72
29	Monte Verde (Chile)	14.6 ka Upper Pleistocene	Outside	Clay	H. sapiens	3 footprints made by an adolescent or a young individual.	73-75
30	Tuc d'Audoubert (France)	13.9 ka Upper Pleistocene	Cave	Clay	H. sapiens	Several trackways and isolated footprints showing toe impressions. Some footprints are attributed to juveniles.	58, 76-77
31	Calvert island (Canada)	13.3-12.6 ka Upper Pleistocene	Outside	Clay	H. sapiens	29 footprints made by at least 3 individuals including 1 juvenile. Toe impressions are still visible.	78
32	Niaux (France)	12.9-12.4 ka Upper Pleistocene	Cave	Clay	H. sapiens	About 40 footprints. Some tracks could have been made during children's game.	79-82
33	Tana della Basura (Italia)	12.3 ka Upper Pleistocene	Cave	Clay	H. sapiens	More than 30 footprints made by juveniles and adults. Initially attributed to Neandertals before new dating associated them with <i>Homo sapiens</i>	83-87
34	Fontanet (France)	12.0 ka Upper Pleistocene	Cave	Clay	H. sapiens	Several isolated footprints. One of them was interpreted as made by a shod foot. Some footprints were attributed to a child following an animal.	42, 88-90
35	Pehuen-Co (Argentina)	12.0 ka Upper Pleistocene	Outside	Sand / Silt	H. sapiens	15 footprints including a trackway and 2 isolated footprints. Toe impressions are visible.	91-92
36	Bogoria (Kenya)	Upper Pleistocene	Outside	Silt	H. sapiens	1 isolated and poorly preserved footprint.	93

#### Text S1: The footprints attributed to Neandertals:

The footprints attributed to Neandertals are very scarce in the fossil record. As we underline in the article, only 9 footprints attributed to this taxon had been reported from 4 different sites (Fig. S2). Before describing these sites and the footprints that they yielded, it is necessary to discuss discoveries made at two other sites: the caves from Ciur-Ibzuc (Romania) and Tana della Basura (Italy).

The researches that were undertaken in the cave from Ciur-Ibzuc during the 1960s initially yielded 400 human footprints before three quarters of them were destroyed (47-48). Initially dated between 15,000 and 10,000 years based on the presence of cave bears, new studies realized from 2012 enabled to date them by radiocarbon between 36,500 and 29,000 years BP (35, 48). These dates place them during a transition phase in Europe characterized by the first evidences of *Homo sapiens* and the very last occurrences of Neandertals (48). Furthermore, the absence of cultural material associated with the footprints makes their taxonomic attribution even more complex. Yet, based on the discovery of *Homo sapiens* remains close to the cave from Ciur-Ibzuc covering contemporary periods, but also on ages for the transition between the two taxa, it is more likely that these footprints were made by *Homo sapiens* (48).

The footprints discovered in the Italian site of Tana della Basura were for a long time attributed to Neandertals (84-85). Indeed, L. Pales used the presence of a Mousterian industry in a nearby cave and remains of cave bears that he considered as contemporary of Neandertals. However, radiometric dating at around 12,000 years BP makes impossible this taxonomic association (86).

#### Biache-Saint-Vaast (France):

The site with the oldest potential Neandertal footprint is Biache-Saint-Vaast. Discovered in 1976, it presents several occupation layers with lithic industry and remains of two human skulls dated to 236,000 years (23-25). A footprint described as poorly preserved and surrounded by ungulate tracks was found in the same layer as the human remains (23-24). It seems that this footprint has also been trampled by ungulates, making difficult its identification and its analysis (24). The presence of a Mousterian industry and the clear attribution of the human remains to Neandertals (94) allow to associate this footprint to a Neandertal individual. A cast of the track surface was realized and is kept in Arkéos Archaeological Museum (Douai, France).

#### Theopetra cave (Greece):

In 1996, four human footprints were discovered in a Mousterian layer of the cave from Theopetra (Greece). This level was firstly radiocarbon dated to 46,300 years BP (95). However, more recent dating using thermoluminescence gave an earlier date: about 130,000 years (29). These footprints were attributed to Neandertals based on these dates and on Mousterian industries found in the same layer (28-29). They would have been made by different individuals (28). The second and third footprints are complete and are 13.8 and 15 cm long which corresponds to statures of 0.86 and 1.00 m and ages of 2 and 4 years (28). The third footprint could have been made by a shod foot representing the oldest occurrence of "footwear" (28). In addition to photographs, the footprints were cast, and the two most complete footprints were digitized in 3D (28).

#### Vârtop cave (Romania):

In 1974, three human footprints were found in a calcareous mud layer in the Romanian cave at Vârtop. U-Th dating has shown that these footprints were made between 97,000 and 62,000 years (34). No archaeological or paleoanthropological remains were discovered in this cave, the association with Neandertals is only based on the U-Th dating. These three footprints appear to have been made by a single individual (35). Two of the three footprints are partial and represent heel or forefoot impressions. The third is relatively complete (22 cm long and 10.6 cm wide) and corresponds to a stature of 1.46 m (33). According to Onac *et al.* (34), a distinctive feature of this complete footprints is a large space (1.6 cm) between the hallux and the second toe. Its morphology would be consistent with our knowledge of Neandertal foot anatomy (34). The most complete footprint has been cast and is kept at the Institute of Speleology in Cluj (Romania).

#### Catalan Bay (Gibraltar):

More recently, a potential Neandertal footprint was discovered in the dune complex from Catalan Bay at Gibraltar. It was made in aeolianite dating by OSL of 28,000 years (49). No archeological or paleoanthropological remains are associated with this footprint. The authors attribute it with Neandertals based on geochronological correlation using archeological and osteological findings at Gibraltar (49). This footprint, described as poorly preserved, was made by descending a slope by an individual whose height was estimated between 1.06 and 1.26 m (49). Given its late chronological age, its attribution to Neandertal remains to be demonstrated.



Fig. S2. Geographical distribution of sites with footprints attributed to Neandertals

# Cherbourg-Octeville Cle Rozel Saint-Lô Channel 20 km

# The archaeological site at Le Rozel (Normandy, France):



Photo: D. Cliquet

Photo: D. Cliquet

Fig. S3. Geographical location of the archaeological site at Le Rozel



The riprap at the base of the dune was built in order to limit the tidal sapping action

Fig. S4. View of Le Rozel site: during the first excavations in 1969 (left) and in 2014 (right).

Satellite view: Geoportail



Fig. S5. Aerial representation of the archeological site from Le Rozel



**Fig. S6.** Geological cross-section of the dune system from Le Rozel and locations of the Paleolithic occupations







Fig. S8. Footprint areas. A-D3b-1, B-D3b-3, C-D3b-2, D-D3b-4, E-D3b-5



**Fig. S9.** Archeological material associated with the footprints: A-Levallois flakes, B-blades, C-knapping spot, D-hearth, E-Burnt log, F-butchery area, G-deer's lower jaw, H-deer's vertebrae

## The tracks from Le Rozel:



**Fig. S10.** Relative frequencies of the Le Rozel tracks (n = 271) from stratigraphic subunits (A), types of tracks (B) and laterality of hominin footprints (n = 257) (C)

**Table S2 (1/5).** Inventory of the tracks discovered at Le Rozel. The maximal length ( $L_{max}$ ) is measured along the longitudinal axis. It will be equal to the total length (L) if the track is longitudinally complete. The distal width (w) corresponds to the largest breadth of the forefoot impression and is measured along the mediolateral axis. The measurements preceded by the sign " $\approx$ " were realized in situ. The measurements made from 3D models are accurate to the millimeter.

	Tavan	Anatomical	Latarality			Recording			Measurement	s
Inventory number	Taxon	part	Laterality	Photos	Cast	Original extraction	3D capture	L <sub>max</sub>	L	w
LREI2012-01	Ното	foot	left	✓				≈13 cm	≈13 cm	≈6 cm
LREI2012-02	Ното	foot	left	$\checkmark$				≈20 cm	unknown	≈9-10 cm
LREI2012-03	Ното	foot	left	$\checkmark$				≈20 cm	unknown	≈11 cm
LREI2013-01	Ното	foot	left	$\checkmark$	$\checkmark$		$\checkmark$	21.9 cm	unknown	6.9 cm
LREI2013-02	Ното	foot	left	✓				≈28 cm	≈28 cm	≈10 cm
LREI2013-03	Ното	foot	unknown	$\checkmark$	$\checkmark$		$\checkmark$	16.9 cm	unknown	unknown
LREI2013-04	Ното	foot	left (?)	$\checkmark$	$\checkmark$		$\checkmark$	23.6 cm	unknown	10.6 cm
LREI2013-05	Ното	foot	left	✓	✓		$\checkmark$	21.4 cm	21.4 cm	9.4 cm
LREI2013-06	Ното	hand	right	✓	✓			≈16 cm	unknown	unknown
LREI2013-07	Ното	foot	right	✓				≈23 cm	unknown	≈8 cm
LREI2013-08	Ното	foot	right	✓				≈26 cm	unknown	≈11 cm
LREI2014-01	Ното	foot	unknown	✓	✓			≈24 cm	unknown	≈11 cm
LREI2014-02	Ното	hand	right	✓	✓			≈19 cm	unknown	unknown
LREI2014-03	Ното	foot	unknown	✓	✓		$\checkmark$	26.0 cm	unknown	unknown
LREI2014-04	Ното	foot	right	✓	✓		✓	29.3 cm	28.7 cm	unknown
LREI2014-05	Ното	foot	right	✓	✓		✓	26.9 cm	26.9 cm	12.3 cm
LREI2014-06	Ното	foot	right	✓			✓	25.6 cm	25.6 cm	9.8 cm
LREI2014-07	Ното	foot	right	✓				≈14 cm	≈14 cm	≈7 cm
LREI2014-08	Ното	foot	left	✓	✓		✓	22.6 cm	22.6 cm	8.9 cm
LREI2014-09	Ното	foot	left	✓				≈11 cm	unknown	≈7 cm
LREI2014-10	Ното	foot	unknown	$\checkmark$				≈6 cm	unknown	≈5 cm
LREI2014-11	Ното	foot	right	$\checkmark$				≈18 cm	unknown	unknown
LREI2014-12	Ното	foot	left	✓	~		✓	24.0 cm	24.0 cm	10.6 cm
LREI2014-13	Ното	hand	right	$\checkmark$	$\checkmark$			≈18 cm	unknown	unknown
LREI2015-01	Ното	foot	left	✓				≈23 cm	unknown	≈9 cm
LREI2015-02	Ното	foot	right (?)	✓				≈8 cm	unknown	unknown
LREI2015-03	Ното	foot	right	✓				≈13 cm	unknown	unknown
LREI2015-04	Ното	foot	right	✓				11.4 cm	11.4 cm	4.5 cm
LREI2015-05	Ното	foot	left	<b>√</b>				≈10 cm	≈10 cm	≈5 cm
LREI2015-06	Ното	foot	unknown	✓				≈7 cm	unknown	unknown
LREI2015-07	Ното	foot	right	<b>√</b>	~		✓	11.5 cm	11.5 cm	4.8 cm
LREI2015-08	Ното	foot	right	✓				≈29 cm	≈29 cm	unknown
LREI2015-09	Ното	foot	unknown	✓	1		1	≈8 cm	unknown	unknown
LREI2015-10	ното	foot	right	•	•		*	7.1 cm	unknown	3.8 cm
LREI2015-11	Homo	foot	right	•	v		<b>▼</b>	12.3 cm	12.3 cm	3.9 cm
LREI2015-12	ното	foot	right	•			•	16.2 cm	unknown	6.1 cm
LREI2015-13	Homo	foot	right	<b>v</b>			<b>▼</b>	17.2 cm	17.2 cm	5.4 cm
LREI2015-14	Homo	foot	left	•			•	19.5 cm	unknown	8.5 CM
LREI2015-15	ното	foot	left	v ./			Y	10.4 cm	unknown	
LREI2015-10	Homo	foot	left	• •			1	≈20 cm	unknown	≈8 cm
LREI2015-17	Homo	foot	right	• •			·	25.5 cm	unknown	7.9 cm
LREI2015-10	Ното	foot	right	•			•	19.5 CIII	unknown	8.0 cm
LREI2015-19	Ното	foot	right	<b>√</b>			1	~24 CIII	unknown	≈9 cm
LREI2015-20	Homo	foot	right	· √	1			13.4 cm	unknown	0.5 cm
LREI2015-21	Ното	foot	unknown	<b>√</b>	•		•	~22.5 CIII	unknown	9.0 cm
LREI2015-22	Homo	foot	right	✓				~20 cm	unknown	~8 cm
LRFI2015-24	Homo	foot	right	✓				≈18 cm	unknown	unknown
LREI2015-24	Homo	foot	unknown	1				unknown	unknown	unknown
I RFI2015-26	Homo	foot	left	✓	✓		✓	21.2 cm	21.2 cm	8.9 cm
I RFI2015-27	Homo	foot	right	✓			✓	≈17 cm	unknown	≈8 cm
I RFI2015-27	Homo	foot	left	✓				~17 Cm ≈21 cm	≈21 cm	unknown
I RFI2015-20	Homo	foot	unknown	<ul> <li>✓</li> </ul>	~		✓	18.5 cm	unknown	unknown
LRFI2015-30	Ното	foot	right	√				≈14 cm	≈14 cm	≈7 cm
LREI2015-30	Homo	foot	right	1				≈12 cm	unknown	unknown
LIVE12013-31	nomo	1001	ngin	•				~12 (11)	UTIKITUWIT	UNKIUWII

**Table S2 (2/5).** Inventory of the tracks discovered at Le Rozel. The maximal length ( $L_{max}$ ) is measured along the longitudinal axis. It will be equal to the total length (L) if the track is longitudinally complete. The distal width (w) corresponds to the largest breadth of the forefoot impression and is measured along the mediolateral axis. The measurements preceded by the sign " $\approx$ " were realized in situ. The measurements made from 3D models are accurate to the millimeter.

	Toyon	Anatomical	Latarality			Recording			Measurement	s
Inventory number	Taxon	part	Laterality	Photos	Cast	Original extraction	3D capture	L <sub>max</sub>	L	w
LREI2015-32	Ното	foot	left	✓				≈18 cm	unknown	≈9 cm
LREI2015-33	Ното	foot	right	✓				≈20 cm	≈20 cm	≈8 cm
LREI2015-34	Ното	foot	left (?)	✓				≈21 cm	unknown	≈6 cm
LREI2015-35	Ното	foot	left	$\checkmark$				≈17 cm	≈17 cm	≈8 cm
LREI2015-36	Ното	foot	right	✓			✓	18.8 cm	18.8 cm	7.7 cm
LREI2015-37	Ното	foot	right	$\checkmark$			✓	13.3 cm	13.3 cm	5.9 cm
LREI2015-38	Ното	foot	right	✓	✓		✓	14.3 cm	unknown	unknown
LREI2015-39	Ното	foot	unknown	$\checkmark$				≈6 cm	unknown	≈5 cm
LREI2015-40	Ното	foot	left	✓				≈21 cm	≈21 cm	≈7 cm
LREI2015-41	Ното	foot	left	$\checkmark$				≈18 cm	≈18 cm	≈8 cm
LREI2015-42	Ното	foot	unknown	✓	~		~	21.5 cm	unknown	7.5 cm
LREI2015-43	Ното	foot	left	$\checkmark$	✓		$\checkmark$	24.9 cm	24.9 cm	unknown
LREI2015-44	Ното	foot	left	✓			✓	25.2 cm	unknown	10.1 cm
LREI2015-45	Ното	foot	left	$\checkmark$	✓		$\checkmark$	21.2 cm	unknown	8.9 cm
LREI2015-46	Ното	foot	right	✓	~		~	18.5 cm	18.5 cm	unknown
LREI2015-47	Ното	foot	left	$\checkmark$	✓		$\checkmark$	18.3 cm	18.3 cm	6.9 cm
LREI2015-48	Ното	foot	left	✓				22.8 cm	unknown	8.9 cm
LREI2015-49	Ното	foot	right	✓	✓		$\checkmark$	16.2 cm	16.2 cm	6.9 cm
LREI2015-50	Ното	foot	right	✓				≈24 cm	≈24 cm	≈9 cm
LREI2015-51	Ното	foot	right	$\checkmark$				≈14 cm	unknown	unknown
LREI2015-52	Ното	foot	left	~				≈18 cm	unknown	≈7 cm
LREI2015-53	Ното	foot	right	✓	✓		√	18.0 cm	18.0 cm	9.6 cm
LREI2016-01	Ното	foot	right	<b>√</b>	~		$\checkmark$	16.5 cm	unknown	6.6 cm
LREI2016-02	Ното	foot	left	✓				≈21 cm	unknown	≈8 cm
LREI2016-03	Ното	foot	left	<b>√</b>	✓		<b>√</b>	13.8 cm	13.8 cm	5.4 cm
LREI2016-04	Ното	foot	right	✓	✓		<b>√</b>	21.3 cm	unknown	9.9 cm
LREI2016-05	Ното	foot	left (?)	✓	✓		<b>√</b>	18.2 cm	unknown	8.4 cm
LREI2016-06	Ното	foot	right	✓	✓		<b>√</b>	22.5 cm	unknown	10.0 cm
LREI2016-07	Ното	foot	left	✓			~	16.4 cm	16.4 cm	8.3 cm
LREI2016-08	Ното	foot	right	✓				≈15 cm	unknown	≈5 cm
LREI2016-09	Ното	hand	right	<b>√</b>	✓		<b>√</b>	11.4 cm	11.4 cm	unknown
LREI2016-10	Ното	foot	left	<b>√</b>	✓		<b>√</b>	17.0 cm	unknown	8.0 cm
LREI2016-11	Ното	foot	left	✓	<b>√</b>		<b>√</b>	12.1 cm	12.1 cm	5.0 cm
LREI2016-12	Ното	foot	right	✓	✓		✓	12.5 cm	12.5 cm	4.9 cm
LREI2016-13	Ното	foot	left	✓	✓		✓	10.1 cm	unknown	unknown
LREI2016-14	Ното	foot	left	✓	✓		~	13.5 cm	unknown	5.3 cm
LREI2016-15	Homo	foot	left	<b>√</b>	✓		<b>√</b>	10.9 cm	unknown	5.9 cm
LREI2016-16	ното	foot	UNKNOWN	•	•		•	19.6 cm	unknown	8.8 cm
LREI2016-17	Ното	foot	right (?)	✓	✓		<b>v</b>	14.5 cm	unknown	6.7 cm
LREI2016-18	Ното	foot	left (?)	✓	•		•	17.5 cm	17.5 cm	7.2 cm
LREI2016-19	Homo	foot	right	<b>v</b>	•		•	26.8 cm	26.8 cm	10.9 cm
LREI2016-20	Homo	foot	left	•	•		•	17.4 cm	unknown	6.9 cm
LREI2010-21	Homo	foot		•	•		•	9.7 cm	unknown	7.5 cm
LREI2010-22	нотто	foot	iert (?)	•	•		•	17.0 cm	unknown	7.3 Cm
LREI2016-23	Homo	foot	right	v .(	•		•	16.5 cm	unknown	6.8 cm
LREI2010-24	Homo	foot	unknown loft	• •	v		v ./	14.3 cm		5.5 Cm
LREI2010-25	Homo	hand	right		1			16.1 cm	16.1 cm	
LREI2010-20	Homo	foot	right					~12 cm		~5 cm
LILIZO10-27	Homo	foot	loft	✓			1	22.0 cm	unknown	~5 cm
LICE12010-20	Homo	foot	right				✓	18.2 cm	unknown	6.7 cm
LREI2010-23	Homo	foot	loft					18.0 cm	unknown	
LREI2010-50	Homo	hand	right				✓	20.9 CIII	unknown	unknown
LILIZO10-31	Homo	foot	right				✓	24.6 cm	unknown	10.5 cm
LREI2010-32	Homo	foot	left	✓			✓	16.1 cm	16.1 cm	7.9 cm
LILIZO10-33	Homo	foot	loft	1				23.6 cm		7.5 cm
LILI2010-34	Homo	1001	leit	v			v	25.0 CIII	unknown	7.7 Cm

**Table S2 (3/5).** Inventory of the tracks discovered at Le Rozel. The maximal length ( $L_{max}$ ) is measured along the longitudinal axis. It will be equal to the total length (L) if the track is longitudinally complete. The distal width (w) corresponds to the largest breadth of the forefoot impression and is measured along the mediolateral axis. The measurements preceded by the sign " $\approx$ " were realized in situ. The measurements made from 3D models are accurate to the millimeter.

Inventory number	Taxan	Anatomical	Latavality	Recording			Measurements			
inventory number	Taxon	part	Laterality	Photos	Cast	<b>Original extraction</b>	3D capture	L <sub>max</sub>	L	w
LREI2016-35	Ното	foot	left (?)	✓			✓	21.5 cm	21.5 cm	8.7 cm
LREI2016-36	Ното	foot	right (?)	$\checkmark$			$\checkmark$	13.1 cm	13.1 cm	unknown
LREI2016-37	Ното	foot	right (?)	✓			✓	17.2 cm	unknown	10.5 cm
LREI2016-38	Ното	foot	unknown	$\checkmark$			$\checkmark$	19.1 cm	unknown	unknown
LREI2016-39	Ното	foot	left	$\checkmark$			✓	16.9 cm	16.9 cm	6.2 cm
LREI2016-40	Ното	foot	right	$\checkmark$			$\checkmark$	11.7 cm	unknown	4.5 cm
LREI2016-41	Ното	foot	left	✓			✓	13.3 cm	unknown	5.3 cm
LREI2016-42	Ното	foot	left	$\checkmark$			$\checkmark$	12.1 cm	unknown	5.9 cm
LREI2016-43	Ното	foot	right	$\checkmark$			$\checkmark$	12.0 cm	unknown	4.6 cm
LREI2016-44	Ното	foot	left	$\checkmark$	$\checkmark$		$\checkmark$	25.8 cm	unknown	10.2 cm
LREI2016-45	Ното	foot	unknown	$\checkmark$			$\checkmark$	17.0 cm	unknown	8.0 cm
LREI2016-46	Ното	foot	right	$\checkmark$				≈23 cm	unknown	unknown
LREI2016-47	Ното	foot	right (?)	$\checkmark$			$\checkmark$	12.2 cm	unknown	7.5 cm
LREI2016-48	Ното	foot	right	$\checkmark$				≈20 cm	≈20 cm	≈8 cm
LREI2016-49	Ното	foot	right	$\checkmark$				≈14 cm	unknown	≈5.5 cm
LREI2016-50	Ното	foot	right	$\checkmark$	$\checkmark$			≈14 cm	unknown	≈6 cm
LREI2016-51	Ното	foot	right	$\checkmark$	$\checkmark$		$\checkmark$	25.1 cm	unknown	9.4 cm
LREI2016-52	Ното	foot	left	$\checkmark$			$\checkmark$	23.5 cm	unknown	8.3 cm
LREI2016-53	Ното	foot	right	$\checkmark$				≈18 cm	unknown	≈7 cm
LREI2016-54	Ното	foot	right	$\checkmark$				≈18 cm	unknown	≈9 cm
LREI2016-55	Ното	foot	left	$\checkmark$	$\checkmark$		$\checkmark$	14.1 cm	unknown	unknown
LREI2016-56	Ното	foot	right	$\checkmark$	$\checkmark$		$\checkmark$	16.4 cm	16.4 cm	6.1 cm
LREI2016-57	Ното	foot	right	$\checkmark$	$\checkmark$		$\checkmark$	21.3 cm	unknown	8.7 cm
LREI2016-58	Ното	foot	left	$\checkmark$				≈25 cm	unknown	≈9 cm
LREI2016-59	Ното	foot	left	✓			$\checkmark$	20.1 cm	unknown	7.3 cm
LREI2016-60	Ното	foot	left	✓	$\checkmark$		$\checkmark$	14.8 cm	14.8 cm	5.6 cm
LREI2016-61	Ното	foot	unknown	✓	$\checkmark$		$\checkmark$	17.5 cm	17.5 cm	5.9 cm
LREI2016-62	Ното	foot	left	$\checkmark$				≈21 cm	unknown	≈7 cm
LREI2016-63	Ното	foot	left	✓				≈20 cm	unknown	≈9 cm
LREI2016-64	Ното	foot	left	$\checkmark$	$\checkmark$			≈16 cm	≈16 cm	unknown
LREI2016-65	Ното	foot	left	~				≈15 cm	unknown	≈8 cm
LREI2016-66	Ното	foot	left	✓	✓		✓	19.5 cm	19.5 cm	8.8 cm
LREI2016-67	Ното	foot	right	~	~		~	15.0 cm	15.0 cm	6.5 cm
LREI2016-68	Ното	foot	right	✓	✓		$\checkmark$	21.7 cm	21.7 cm	6.9 cm
LREI2016-69	Ното	foot	unknown	<b>√</b>				unknown	unknown	unknown
LREI2016-70	Ното	foot	left	√				≈16.5 cm	unknown	unknown
LREI2016-71	Ното	foot	left	<b>√</b>			<b>√</b>	23.6 cm	23.6 cm	8.7 cm
LREI2016-72	Ното	foot	left	<b>√</b>			✓	17.5 cm	17.5 cm	7.8 cm
LREI2016-73	Ното	foot	left	<b>√</b>			✓	22.8 cm	22.8 cm	8.1 cm
LREI2016-74	Homo	foot	left	V				≈14 cm	unknown	≈7 cm
LREI2016-75	Homo	toot	right	<b>√</b>			<b>√</b>	6.5 cm	unknown	4.5 cm
LREI2016-76	Ното	toot	left	V			~	9.7 cm	unknown	4.6 cm
LKE12016-77	ното	foot	left	v				≈14 cm	≈14 cm	≈/cm
LREI2016-78	Ното	foot	right (?)	✓			<b>√</b>	10.3 cm	unknown	5.4 cm
LREI2016-79	Ното	foot	right	<b>v</b>		1	~	22.1 cm	22.1 cm	9.6 cm
LREI2017-01	Ното	foot	left	✓		✓		≈19 cm	unknown	unknown
LREI2017-02	ното	foot	right (?)	<b>v</b>				≈19 cm	unknown	≈7.5 cm
LKEI2017-03	ното	1000	left (?)	v				≈12.5 cm	unknown	≈5 cm
LKEI2017-04	ното	1000	iert (?)	v		<b>v</b>	•	17.4 cm	unknown	7.6 CM
LKEI2017-05	ното	1000	unknown	v		V		≈19 cm	unknown	≈10 cm
LKEI2017-06	ното	1000	right (?)	v		1		≈18 cm	unknown	≈7.5 cm
LKEI2017-07	ното	1000	right	v		V	•	15.4 CM	15.4 CM	7.3 CM
LKEI2017-08	ното	1000	right	v		<b>v</b>	<b>v</b>	17.5 cm	unknown	5.9 CM
LKEI2017-09	Homo	foot	unknown	· · ·		V	v	17.8 cm	unknown	7.8 cm
LREI2017-10	Hom -	foot	right (2)	v				≈21 CM	unknown	≈o cm
LKEI2017-11	ното	root	right (?)	~		✓	✓	20.3 cm	unknown	7.2 cm

**Table S2 (4/5).** Inventory of the tracks discovered at Le Rozel. The maximal length ( $L_{max}$ ) is measured along the longitudinal axis. It will be equal to the total length (L) if the track is longitudinally complete. The distal width (w) corresponds to the largest breadth of the forefoot impression and is measured along the mediolateral axis. The measurements preceded by the sign " $\approx$ " were realized in situ. The measurements made from 3D models are accurate to the millimeter.

Inventory number	Tayon	Anatomical	Pr	Recording				Measurements		
Inventory number	Taxon	part	Laterality	Photos	Cast	Original extraction	3D capture	L <sub>max</sub>	L	w
LREI2017-12	Ното	foot	left	$\checkmark$		$\checkmark$		≈15 cm	≈15 cm	≈7 cm
LREI2017-13	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	17.9 cm	17.9 cm	unknown
LREI2017-14	Ното	foot	unknown	$\checkmark$		$\checkmark$	$\checkmark$	15.0 cm	unknown	6.3 cm
LREI2017-15	Ното	foot	right (?)	$\checkmark$				≈19 cm	unknown	≈5 cm
LREI2017-16	Ното	foot	right (?)	$\checkmark$				≈18 cm	unknown	≈7 cm
LREI2017-17	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	24.3 cm	unknown	9.8 cm
LREI2017-18	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	19.2 cm	19.2 cm	8.6 cm
LREI2017-19	Ното	foot	right	$\checkmark$			$\checkmark$	21.7 cm	unknown	9.3 cm
LREI2017-20	Ното	foot	left (?)	$\checkmark$		$\checkmark$	$\checkmark$	20.1 cm	unknown	10.6 cm
LREI2017-21	Ното	foot	unknown	$\checkmark$			$\checkmark$	12.2 cm	unknown	unknown
LREI2017-22	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	20.5 cm	20.5 cm	8.1 cm
LREI2017-23	Ното	foot	left	$\checkmark$				≈17 cm	unknown	≈9 cm
LREI2017-24	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	20.7 cm	unknown	9.5 cm
LREI2017-25	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	22.0 cm	unknown	9.8 cm
LREI2017-26	Ното	foot	right	$\checkmark$			$\checkmark$	19.5 cm	unknown	unknown
LREI2017-27	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	25.9 cm	unknown	11.2 cm
LREI2017-28	Ното	hand	right	$\checkmark$		$\checkmark$	$\checkmark$	15.7 cm	15.7 cm	9.1 cm
LREI2017-29	Ното	foot	left	$\checkmark$			$\checkmark$	22.9 cm	unknown	9.0 cm
LREI2017-30	Ното	foot	right	$\checkmark$		$\checkmark$	$\checkmark$	26.4 cm	unknown	12.8 cm
LREI2017-31	Ното	foot	right	$\checkmark$			$\checkmark$	18.4 cm	unknown	7.8 cm
LREI2017-32	Ното	foot	right (?)	$\checkmark$		$\checkmark$	$\checkmark$	21.3 cm	21.3 cm	10.3 cm
LREI2017-33	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	18.7 cm	unknown	7.8 cm
LREI2017-34	Ното	foot	right (?)	$\checkmark$				≈17 cm	unknown	≈9 cm
LREI2017-35	Ното	foot	right (?)	$\checkmark$				≈24.5 cm	unknown	≈9 cm
LREI2017-36	Ното	foot	unknown	$\checkmark$		$\checkmark$	$\checkmark$	24.0 cm	unknown	9.5 cm
LREI2017-37	Ното	foot	right	$\checkmark$			$\checkmark$	20.8 cm	unknown	9.2 cm
LREI2017-38	Ното	foot	right	✓				≈23 cm	unknown	≈9 cm
LREI2017-39	Ното	foot	right	$\checkmark$				≈23 cm	unknown	≈12.5 cm
LREI2017-40	Ното	foot	unknown	✓				≈15 cm	≈15 cm	≈7 cm
LREI2017-41	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	18.1 cm	18.1 cm	7.6 cm
LREI2017-42	Ното	foot	right	$\checkmark$				22.0 cm	unknown	9.9 cm
LREI2017-43	Ното	foot	right	✓				≈16 cm	unknown	≈8 cm
LREI2017-44	Animalia (Carnivora)	paw	unknown	$\checkmark$		$\checkmark$	$\checkmark$	7.6 cm	unknown	4.8 cm
LREI2017-45	Homo	foot	right	✓		$\checkmark$	$\checkmark$	22.6 cm	unknown	9.4 cm
	Animalia			,		,	,			
LREI2017-46	(Carnivora)	paw	unknown	v		V	v	11.2 cm	unknown	9.0 cm
LREI2017-47	Homo	foot	right	$\checkmark$		$\checkmark$	$\checkmark$	24.7 cm	24.7 cm	12.3 cm
LREI2017-48	Ното	foot	left	$\checkmark$		✓	✓	17.2 cm	unknown	9.2 cm
LREI2017-49	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	21.4 cm	21.4 cm	9.2 cm
LREI2017-50	Animalia (Ruminantia . Cervidae)	paw	unknown	$\checkmark$		$\checkmark$	✓	8.1 cm	unknown	3.3 cm
LREI2017-51	Homo	foot	right	$\checkmark$		✓	$\checkmark$	29.1 cm	unknown	10.5 cm
LREI2017-52	Ното	foot	right	~		✓		23.2 cm	23.2 cm	9.4 cm
LREI2017-53	Ното	foot	right (?)	$\checkmark$			✓	22.1 cm	unknown	10.1 cm
LREI2017-54	Ното	foot	unknown	✓		✓	✓	15.2 cm	unknown	unknown
LREI2017-55	Ното	foot	unknown	$\checkmark$		✓	$\checkmark$	13.4 cm	unknown	unknown
LREI2017-56	Ното	foot	right	~				22.9 cm	22.9 cm	8.7 cm
LREI2017-57	Ното	foot	left	$\checkmark$			$\checkmark$	19.4 cm	19.4 cm	7.8 cm
LREI2017-58	Ното	foot	left	$\checkmark$				≈18 cm	unknown	≈7 cm
LREI2017-59	Ното	foot	left (?)	$\checkmark$				≈16 cm	unknown	≈7 cm
LREI2017-60	Ното	foot	unknown	~				≈14 cm	unknown	≈8 cm
LREI2017-61	Ното	foot	right	$\checkmark$		$\checkmark$	$\checkmark$	18.3 cm	unknown	6.4 cm
LREI2017-62	Ното	foot	left (?)	~		✓	~	17.7 cm	unknown	6.5 cm
LREI2017-63	Ното	foot	right	$\checkmark$		$\checkmark$	$\checkmark$	23.8 cm	unknown	9.2 cm

**Table S2 (5/5).** Inventory of the tracks discovered at Le Rozel. The maximal length ( $L_{max}$ ) is measured along the longitudinal axis. It will be equal to the total length (L) if the track is longitudinally complete. The distal width (w) corresponds to the largest breadth of the forefoot impression and is measured along the mediolateral axis. The measurements preceded by the sign " $\approx$ " were realized in situ. The measurements made from 3D models are accurate to the millimeter.

	-	Anatomical				Recording		Measurements		
Inventory number	Taxon	part	Laterality	Photos	Cast	Original extraction	3D capture	L <sub>max</sub>	L	w
LREI2017-64	Ното	foot	left	✓			✓	23.4 cm	23.4 cm	8.9 cm
LREI2017-65	Animalia	waq	unknown	✓		✓	$\checkmark$	3.8 cm	unknown	6.6 cm
	(Carnivora)	( ).	1.0	1						
LREI2017-66	Homo	foot	left	V			<b>√</b>	21.6 cm	unknown	11.2 cm
LREI2017-67	Animalia (Carnivora.	paw	unknown	✓		~	✓	7.4 cm	7.4 cm	5.7 cm
	Canidae)									
LREI2017-68	Homo	foot	left	✓		✓	✓	23.8 cm	unknown	11.2 cm
LREI2017-69	Ното	hand	right	✓		$\checkmark$	$\checkmark$	14.1 cm	14.1 cm	13.7 cm
LREI2017-70	Ното	foot	left (?)	<b>√</b>			,	≈20 cm	unknown	≈10 cm
LREI2017-71	Ното	foot	right	<b>√</b>		1	✓	23.4 cm	23.4 cm	11.1 cm
LREI2017-72	Ното	foot	left	<b>√</b>		✓	<b>√</b>	20.4 cm	unknown	9.5 cm
LREI2017-73	Ното	foot	left	V		✓	✓	22.1 cm	22.1 cm	10.2 cm
LREI2017-74	Homo	foot	right	<ul> <li>✓</li> </ul>		v		≈15.5 cm	unknown	≈9 cm
LREI2017-75	Homo	foot	right	• •		v ./	•	17.3 cm	17.3 cm	8.6 cm
LREI2017-70	Homo	foot	right	• ./		v	v	20.9 cm	unknown	9.4 cm
LREI2017-77	Homo	foot	right (2)	•		1	1	≈14 cm	unknown	≈0 cm
LREI2017-70	Ното	foot	loft (2)	• •		v	¥	~10 cm		~10 cm
LREI2017-75	Δnimalia	1001	iert (:)	•				~19 Cill	~19 CIII	~10 cm
LREI2017-80	(Carnivora)	paw	unknown	$\checkmark$		$\checkmark$	$\checkmark$	7.3 cm	unknown	8.9 cm
LREI2017-81	Homo	foot	left (?)	$\checkmark$			$\checkmark$	19.7 cm	unknown	8.0 cm
LREI2017-82	Ното	foot	left	✓		✓		≈23.5 cm	unknown	≈10 cm
LREI2017-83	Ното	foot	right	$\checkmark$		$\checkmark$		≈22 cm	unknown	≈9 cm
LREI2017-84	Ното	foot	right	✓		✓		≈28 cm	≈28 cm	≈12 cm
LREI2017-85	Ното	foot	right	$\checkmark$				≈23 cm	unknown	≈9 cm
LREI2017-86	Ното	foot	right	$\checkmark$		$\checkmark$	$\checkmark$	28.4 cm	28.4 cm	12.2 cm
LREI2017-87	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	27.5 cm	27.5 cm	9.9 cm
LREI2017-88	Ното	foot	left (?)	✓				≈16 cm	unknown	≈5 cm
LREI2017-89	Ното	foot	left	$\checkmark$		$\checkmark$	$\checkmark$	20.2 cm	unknown	unknown
LREI2017-90	Ното	foot	left	✓			✓	13.8 cm	unknown	8.9 cm
LREI2017-91	Ното	foot	right	✓		$\checkmark$	√	16.2 cm	unknown	8.8 cm
LREI2017-92	Ното	foot	left	✓			✓	22.1 cm	unknown	10.1 cm
LREI2017-93	Ното	foot	right (?)	✓				≈20 cm	unknown	≈7 cm
LREI2017-94	Ното	foot	right	<b>√</b>		,		≈18 cm	unknown	≈9 cm
LREI2017-95	Ното	foot	right	<b>√</b>		<b>√</b>	✓	18.8 cm	18.8 cm	7.6 cm
LREI2017-96	Ното	foot	left	<b>√</b>		✓	<b>√</b>	15.1 cm	unknown	7.0 cm
LREI2017-97	Ното	foot	left	<b>v</b>			•	19.9 cm	unknown	9.3 cm
LREI2017-98	Homo	foot	right	V		V	<b>v</b>	24.9 cm	unknown	11.4 cm
LREI2017-99	Homo	foot	right	•		•	v	17.4 Cm	unknown	7.9 Cm
LREI2017-100	Ното	foot	right	• •		<b>v</b>	1	≈13 cm	15.9 cm	~o cm
LREI2017-101	Homo	foot	right	· ✓		•		15.8 cm	15.8 cm	7.0 cm
LREI2017-102	Ното	foot	right (?)	√			√	22.1 cm	22.1 cm	12.5 cm
LREI2017-104	Ното	foot	right	√		✓	√	17.0 cm	17.0 cm	9.2 cm
LREI2017-105	Ното	foot	left	✓		✓	✓	21.7 cm	21.7 cm	8.8 cm
LREI2017-106	Ното	foot	left	✓		✓		≈26 cm	unknown	≈9 cm
LREI2017-107	Ното	foot	left	$\checkmark$		✓	✓	16.1 cm	unknown	9.2 cm
LREI2017-108	Ното	foot	right (?)	~		$\checkmark$	$\checkmark$	16.5 cm	unknown	8.5 cm
LREI2017-109	Ното	foot	left (?)	$\checkmark$			$\checkmark$	19.5 cm	unknown	8.9 cm
LREI2017-110	Ното	foot	right	$\checkmark$			$\checkmark$	19.3 cm	19.3 cm	10.0 cm
LREI2017-111	Ното	foot	unknown	$\checkmark$			✓	16.9 cm	16.9 cm	unknown
LREI2017-112	Ното	foot	left	~		✓	✓	15.6 cm	15.6 cm	7.1 cm
LREI2017-113	Ното	foot	unknown	✓		$\checkmark$	$\checkmark$	14.6 cm	unknown	8.2 cm
LREI2017-114	Ното	foot	right (?)	$\checkmark$		$\checkmark$	$\checkmark$	34.0 cm	unknown	12.2 cm
LREI2017-115	Ното	foot	right	$\checkmark$		$\checkmark$	$\checkmark$	16.1 cm	unknown	≈9 cm

# The hominin footprints from Le Rozel:



Fig. S11 (1/6). Views and associated outlines of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)



Fig. S11 (2/6). Views and associated outlines of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)



Fig. S11 (3/6). Views and associated outlines of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)



Fig. S11 (4/6). Views and associated outlines of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)







LREI2017-86

LREI2017-87

Fig. S11 (5/6). Views and associated outlines of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)



Fig. S11 (6/6). Views and associated outlines of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)

## Text S2: Depth distribution of the footprints from Le Rozel:

The "depth maps" below were obtained thanks to the *Foot Processor* software. This software, running in *Mathworks Runtime Compiler 7.11*, was written by M. Budka (Bournemouth University). It is available for download (<u>http://footprints.bournemouth.ac.uk/</u>).



Fig. S12 (1/4). Views and shaded 3D elevation models of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)



Fig. S12 (2/4). Views and shaded 3D elevation models of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)



LREI2017-71

Fig. S12 (3/4). Views and shaded 3D elevation models of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)





LREI2017-87





LREI2017-105

Fig. S12 (4/4). Views and shaded 3D elevation models of the footprints from Le Rozel (scale bar: 2 cm) (Photos: D. Cliquet)

## **Text S3: Experimental footprints:**

An experimental study was carried out in 2017. 192 footprints were (Fig. S13) made by 21 individuals, from 1 to 36 years old, under deposition conditions similar to the fossil footprints.

## • Experimental area:

An experimental trackway (Fig. S13) was built from sediments, a fine and medium brown to black sand, extracted from the footprint areas during the excavations. This trackway is composed of a flat part, 6 meters long and 2 meters wide, where the experimental sequences described below took place. In order to respect the non-uniform moisture conditions in which the fossil footprints were made, the experimental sequences were carried out under different weather conditions, ranging from dry periods to important rainfalls that brought mud flows comparable to those observed on the archeological site.

## Participants:

The participants were born in France and lived near the site during the excavations in 2017. They were recruited after volunteering and signed a charter to give their consent after being informed of the purpose and the conduct of the study. Their date of birth was informed as well as their stature and the length of their feet which were measured on the same day as the experimental sequence (Tables S3-S4).

## Experimental sequence:

Each participant was asked to move barefoot and in a straight line by walking and slow running. The length and width of each footprint were measured *in situ* by several observers (Tables S3-S4). For each sequence, a right and a left footprint were also 3D digitized in *Agisoft Photoscan* (v.1.4.0) using a Canon EOS 1300D. These 3D models allowed to control the measurements realized *in situ* and they were studied in the morphometric analyses (Texts S5-S6).



Fig. S13. Experimental footprints (A) and trackway (B)

	Number of individuals	Min.	Mean	Max.
Age (years)	21	1	21	36
Stature (cm)	21	87.0	162.2	182.3
Average foot length (cm)	20	20.4	23.4	26.6
Number of measurable footprints	21	2	9.1	19
Footprint length (cm)	21	13.1	23.7	29.2

Table S4.	Experimental	data
-----------	--------------	------

	1.00	Ctatura	Average feet	Number of		Fc	otprint	length (cm)
Individuals	Age (years)	(cm)	length (cm)	measurable footprints	Min.	Mean	Max.	Maximum deviation from the mean (m.d.)
#1	11	153.9	22.4	11	22.7	25.3	28.5	12.8%
#2	22	177.1	24.5	9	23.6	25.7	28.2	9.7%
#3	22	166.9	21.8	15	21.6	23.1	25.1	8.6%
#4	13	153.8	20.5	8	21.5	23.0	25.2	9.8%
#5	31	171.2	23.7	15	23.2	25.4	27.9	9.7%
#6	22	175.8	24.1	19	23.6	25.5	28.2	10.4%
#7	34	160.0	23.2	10	22.1	23.3	24.5	5.3%
#8	36	171.8	25.5	7	24.5	25.0	25.3	2.1%
#9	23	182.3	23.3	11	22.2	24.7	27.1	10.2%
#10	23	170.8	25.3	15	24.6	25.9	27.5	6.1%
#11	1	87.0		2	13.1	14.1	15.0	6.8%
#12	28	172.9	26.6	7	24.2	25.3	26.5	4.6%
#13	19	173.5	26.2	9	24.1	26.9	29.2	10.3%
#14	10	146.0	21.5	4	20.5	21.7	23.5	8.3%
#15	21	169.0	22.7	9	22.6	23.7	25.8	8.8%
#16	21	178.5	25.1	10	24.3	25.5	26.8	5.1%
#17	34	172.0	24.0	6	22.3	23.4	24.5	4.8%
#18	24	157.0	20.7	6	21.2	22.3	23.5	5.6%
#19	25	162.0	22.6	4	22.2	22.9	23.5	2.8%
#20	21	156.0	20.4	5	20.1	21.3	22.5	5.7%
#21	22	171.8	24.3	10	23.1	24.4	25.5	5.2%

### **Text S4: Comparative materials**

For both the identification of the Le Rozel tracks as human footprints and for morphometric analyses, we used several 3D models of footprints from different sites and ages made by barefoot individuals. Most of them were made by *Homo sapiens* in soft ground comparable to the sedimentary deposit conditions at Le Rozel. Furthermore, in order to consider the impact of the habitual wearing of footwear on the pedal morphology, particularly on the longitudinal arch (*e.g.* 98-99), footprints made by both habitually shod (experimental footprints) and presumably unshod individuals (Holocene archeological footprints) were used.

This comparative material includes 52 footprints from our experimental study (see Text S3) as well as 3D models available online and in particular the database of Professor M. Bennett (<u>http://footprints.bournemouth.ac.uk/</u>) that we would like to acknowledge. These comparative samples are:

- 21 Holocene footprints from the Formby Point site (Great-Britain) made between 3.6 and 3.2 Ka BP in a soft and wet ground composed of silt to fine sand in a dune context (100-101). <a href="http://footprints.bournemouth.ac.uk">http://footprints.bournemouth.ac.uk</a>
- 22 Pleistocene footprints from the White Sands National Monument site (USA) made probably between 15.6 and 10.0 Ka BP in muds and sands located in a paleolake context by several individuals that presumably hunted sloth (72).

http://footprints.bournemouth.ac.uk

 18 Pliocene footprints from the Laetoli site, which were made in 3.66 Ma old volcanic ash and are attributed to *Australopithecus*. 11 of them compose the G1 trackway (5). <a href="http://footprints.bournemouth.ac.uk">http://footprints.bournemouth.ac.uk</a>

They are completed by 6 footprints from the S1 trackway and a footprint from the S2 trackway (7) whose 3D models are available at:

<u>https://www.morphosource.org/Detail/ProjectDetail/Show/project\_id/298</u> (morphometric data acquired by S. Menconero, access provided by M. Cherin, Università di Perugia, Italy).

#### Text S5: Identification as human footprints:

Identification criteria of the footprints from Le Rozel as hominin footprints are the following: a rounded heel, a longitudinal arch, relatively short toes, a mediolateral decrease of the length of the toes, an adducted hallux and maximum depth beneath the heel and the forefoot (41, 102-105).

We also tested the diagnosis through the morphometric method of Morse et al. (41, 106). Five geometric landmarks (Fig. S14) were placed using *Geomagic Studio 2013* and their 2D coordinates were subjected to a Generalized Procrustes Analysis using *PAST* (v.3.0; refs. 107-108). These landmarks were placed on 93 footprints from Le Rozel that meet all the following criteria: the footprint must (a) be digitized in 3D, (b) be complete enough to place the landmarks, (c) be made on a flat ground and (d) show no evidence of slide.

This method focuses on geometry rather than on the conservation of anatomical details, which is useful for footprints whose conservation is variable, as it is the case in dune context. For Le Rozel, we applied it to 36 footprints showing detailed anatomical traits, such as clear toe impressions, and to 57 less well-preserved footprints but that showed sufficient details to allow placing the 5 landmarks. Statistical comparisons were carried out between these two sub-samples for post-Procrustes lengths (interlandmark distance along the longitudinal axis) and widths (interlandmark distance along the longitudinal axis). No statistical difference was observed (ANOVA: P >> 0.05).

The positions of the landmarks were then compared with those of other footprints from our comparative materials (for more details see Texts S3-S4). The spatial distribution of the landmarks differs little from the other hominin footprints (Fig. S14). The maximum depth points are located either at the heel or at the forefoot, which is a characteristic for hominin footprints (41, 106). However, one footprint from Le Rozel (LREI2016-23) has a maximum depth located at the midfoot. This is however more probably linked to an irregularity of the ground rather than to anatomical differences or an erroneous identification. Such a midfoot position is also found in a few instances in the *H. sapiens* comparative samples (Fig. S14).



**Fig. S14.** Post-Procrustes landmark positions of the footprints from Le Rozel compared with other hominin footprints. The first four landmarks are placed on the axes of an ellipse enclosing the footprint. 1, 2: ends of the longitudinal axis; 3-4: ends of the footprints along the small axis of the ellipse; 5: position where the depth of the footprint is maximum (41, 106)

Both post-Procrustes lengths and widths of Le Rozel footprints fall within the distribution range of the other hominin footprints (Fig. S15). On average, they are longer than other hominin footprints except those from the Pleistocene White Sands National Monument footprints. They are less wide than the footprints from Laetoli and White Sands National Monument. They are close to the widths of the Holocene footprints from Formby Point. Finally, we grouped the footprints by sites and by attributed taxa and we applied cluster analysis (UPGMA), using PAST to the mean coordinates of each footprint group. The footprints from Le Rozel are included in the same group as the other hominin footprints. They are grouped with the footprints attributed to *Homo sapiens* and differ more from the Laetoli footprints. They are closer to the Pleistocene footprints from White Sands National Monument, their sister-group, than to the Holocene archaeological footprints and the experimental footprints (Fig. S16). Therefore, the spatial distribution of the landmarks is consistent with their identification as hominin footprints.



Fig. S15. Post-Procrustes length and width



**Fig. S16.** UPGMA-tree obtained from the mean coordinates of footprints grouped by sites (A) and attributed taxon (B)

#### Text S6: Geometric morphometrics:

We studied the locations of 11 landmarks based on those used by Bennett *et al.* (Fig. S17; ref. 8). Nine of them informed on footprint outline. The first two (#1-2) are located at the ends of the longitudinal axis. The following six are placed two by two along mediolateral axes and represent the maximum heel impression width (#3-4), the minimum midfoot impression width (#5-6) and the maximum forefoot impression width (#7-8). The ninth landmark is located at the hallux tip impression (#9). The last two landmarks represent the deepest points of the heel (#10) and the forefoot (#11) that inform on weight transfer during human walking (ref. 8).



**Fig. S17.** Landmarks used during the geometric morphometric analysis.

The landmarks were placed on 3D models of the footprints that comply with all of the following conditions: each footprint must be longer than 18 cm to avoid ontogenetic biases; its outline must be complete without discontinuity; there must be no apparent evidence of sedimentary reworking or alteration that occurred after track formation; it must have been made on a flat ground and does not show any evidence of slide.

Each landmark was placed in *Geomagic studio 2013* on 14 footprints from Le Rozel (1 from the D3b-3 stratigraphic subunit, 12 from the D3b-4, 1 from the D3b-5), as well as on 48 of our experimental footprints (Text S3). Additionally, they were placed on 10 Holocene archaeological footprints, made by individuals presumed to be habitually unshod and on 17 Laetoli footprints as an outgroup (Text S4 and Table S5). Once the landmarks were placed, their 2D coordinates were subjected to a Generalized Procrustes Analysis in PAST (v.3.0; refs. 107-108) removing the effects of size and orientation. The mean 2D coordinates of the footprints made by a same individual were calculated and subsequently used to avoid biases caused by statistical replication. The 2D coordinates were then used within a Principal Component Analysis (PCA) in PAST.

**Table S5.** 3D models used during the geometric morphometric analysis.<sup>1</sup>3D surfaces digitized in situ.<sup>2</sup>3D surfaces from freely accessible databases(http://footprints.bournemouth.ac.uk/; https://www.morphosource.org/)

Site	Period	Attributed taxon	Ground properties	Number of individuals	Number of footprints
Exp	erimental <sup>1</sup>	H. sapiens	Nearby the site in a similar sediment	18	46
Formby Point <sup>2</sup>	Holocene	H. sapiens	Silt and sand	10	10
Le Rozel <sup>2</sup>	Upper Pleistocene	H. neanderthalensis	Dune sand/Sandy mud	Unknown	14
Laetoli <sup>2</sup>	Pliocene	Au. afarensis	Volcanic tuff	3 (G1, S1, S2)	17

# Biometry of the footprints from the D3b-4 stratigraphic subunit

		Longitudinal	Width (cm) -	Estimated stature (cm)			
Footprints	Length (cm)	completeness		from footprint length	from footprint width		
LREI2015-04	11.4	complete	4.5	73.8	65.8		
LREI2015-12	16.2	uncertain	6.1		90.2		
LREI2015-45	21.2	uncertain	8.9		130.9		
LREI2015-46	18.5	complete		120.0			
LREI2015-47	18.3	complete	6.9	119.1	102.0		
LREI2015-48	22.8	uncertain	8.9		131.6		
LREI2015-49	16.2	complete	6.9	105.4	101.3		
LREI2015-53	18.0	complete	9.6	117.1	141.2		
LREI2016-01	16.5	uncertain	6.6		97.6		
LREI2016-03	13.8	complete	5.4	89.8	79.8		
LREI2016-14	13.5	uncertain	5.3		78.4		
LREI2016-15	10.9	uncertain	5.9		87.2		
LREI2016-16	19.6	uncertain	8.8		130.1		
LREI2016-23	16.5	uncertain	6.8		100.6		
LREI2016-24	14.3	uncertain	5.5		81.3		
LREI2016-25	16.2	complete	7.2	105.4	106.5		
LREI2016-28	22.0	uncertain	10.4		153.8		
LREI2016-32	24.6	uncertain	10.5		155.3		
LREI2016-33	16.1	complete	7.9	104.7	116.8		
LREI2016-34	23.6	uncertain	7.7		113.9		
LREI2016-35	21.5	complete	8.7	139.9	128.6		
LREI2016-36	13.1	complete		85.2			
LREI2016-37	17.2	uncertain	10.5		154.5		
LREI2016-39	16.9	complete	6.2	110.0	91.7		
LREI2016-40	12.5	uncertain	4.9		72.5		
LREI2016-42	12.1	uncertain	5.9		86.5		
LREI2016-43	12.0	uncertain	4.6		68.0		
LREI2016-44	25.8	uncertain	10.2		150.8		
LREI2016-45	17.0	uncertain	8.0		117.6		
LREI2016-47	12.2	uncertain	7.5		110.2		
LREI2016-51	25.1	uncertain	9.4		139.0		
LREI2016-52	23.5	uncertain	8.3		122.7		
LREI2016-56	16.4	complete	6.1	106.7	90.2		
LREI2016-59	20.1	uncertain	7.3		107.9		
LREI2016-60	14.8	complete	5.6	96.0	82.8		
LREI2016-75	8.1	uncertain	4.6		67.3		
LREI2016-78	10.3	uncertain	5.4		79.8		
LREI2017-04	17.4	uncertain	7.6		112.4		
LREI2017-07	15.4	complete	7.3	100.2	107.9		
LREI2017-08	17.5	uncertain	5.9		87.2		
LREI2017-09	17.8	uncertain	7.8		115.3		
LREI2017-11	20.3	uncertain	7.2		106.5		
LREI2017-13	17.9	complete		116.5			
LREI2017-14	15.0	uncertain	6.3		93.2		
LREI2017-17	24.3	uncertain	9.8		144.9		
LREI2017-18	19.2	complete	8.6	124.9	127.2		
LREI2017-19	21.7	uncertain	9.3		137.5		
LREI2017-20	20.1	uncertain	10.6		156.7		
LREI2017-22	20.5	complete	8.1	133.4	119.8		
LREI2017-24	20.7	uncertain	9.5		140.5		
LREI2017-25	22.0	uncertain	9.8		144.9		

Table S6 (1/2). Dimensions and estimated statures for 104 footprints from the D3b-4 subunit

		Longitudinal		Estimated stature (cm)			
Footprints	Length (cm)	completeness	Width (cm)	from footprint length	from footprint width		
LREI2017-27	25.9	uncertain	11.2		165.6		
LREI2017-29	22.9	uncertain	9.0		133.1		
LREI2017-30	26.4	uncertain	12.8		189.3		
LREI2017-31	18.4	uncertain	8.9		131.6		
LREI2017-32	21.3	complete	10.3	138.6	152.3		
LREI2017-33	18.7	uncertain	7.8		115.3		
LREI2017-36	24.0	uncertain	9.5		140.5		
LREI2017-37	20.8	uncertain	9.2		136.0		
LREI2017-41	18.1	complete	7.6	117.4	112.4		
LREI2017-42	22.0	uncertain	9.9		146.4		
LREI2017-45	22.6	uncertain	9.4		139.0		
LREI2017-47	24.7	complete	12.3	160.7	181.9		
LREI2017-48	17.2	uncertain	9.2		136.0		
LREI2017-49	21.4	complete	9.2	138.9	135.3		
LREI2017-51	29.1	slide	10.5		155.3		
LREI2017-52	23.2	complete	9.4	150.9	139.0		
LREI2017-53	22.1	uncertain	10.1		149.3		
LREI2017-56	22.9	complete	8.7	149.0	128.6		
LREI2017-57	19.4	complete	7.8	126.2	115.3		
LREI2017-63	23.8	uncertain	9.2		136.0		
LREI2017-64	23.4	complete	8.9	151.9	131.6		
LREI2017-66	21.6	uncertain	11.2		165.6		
LREI2017-68	23.8	uncertain	11.2		164.9		
LREI2017-71	23.4	complete	11.1	151.9	164.1		
LREI2017-73	22.1	complete	10.2	143.8	150.8		
LREI2017-75	17.3	complete	8.6	112.6	127.2		
LREI2017-76	20.9	uncertain	9.4		139.0		
LREI2017-78	22.3	uncertain	8.7		128.6		
LREI2017-81	19.7	uncertain	8.0		118.3		
LREI2017-86	28.4	complete	12.2	184.8	180.4		
LREI2017-87	27.5	complete	9.9	178.9	146.4		
LREI2017-90	13.8	uncertain	8.9		131.6		
LREI2017-91	16.2	uncertain	8.8		130.1		
LREI2017-92	22.1	uncertain	10.1		149.3		
LREI2017-95	18.8	complete	7.6	122.3	112.4		
LREI2017-96	15.1	uncertain	7.0		103.5		
LREI2017-97	19.9	uncertain	9.3		137.5		
LREI2017-98	24.9	uncertain	11.4		168.6		
LREI2017-99	17.4	uncertain	7.9		116.8		
LREI2017-101	15.8	complete	7.0	102.5	102.8		
LREI2017-102	16.4	complete	7.2	106.4	106.5		
LREI2017-103	22.1	complete	12.5	143.5	184.1		
LREI2017-104	17.0	complete	9.2	110.3	135.3		
LREI2017-105	21.7	complete	8.8	141.2	130.1		
LREI2017-107	16.1	uncertain	9.2		135.3		
LREI2017-108	16.5	uncertain	8.5		125.7		
LREI2017-109	19.5	uncertain	8.9	105 -	131.6		
LREI2017-110	19.3	complete	10	125.6	147.9		
LREI2017-111	16.9	complete	7.05	110.0	101.2		
LKEI2017-112	15.6	complete	7.05	101.2	104.2		
LKEI2017-113	14.6	uncertain	8.2		121.3		
LKEI2017-114	34.0	slide	12.2		180.4		
LKEIZU1/-115	10.1	uncertain	ð.3		122.7		

Table S6 (2/2). Dimensions and estimated statures for 104 footprints from the D3b-4 subunit

	Frequency	Min.	Mean	Max.	s.d.
Length (cm)	39	11.4	19.0	28.4	3.8
Width (cm)	100	4.5	8.5	12.8	1.9
Width / Length	35	0.36	0.44	0.56	0.05

**Table S7.** Statistical parameters of the dimensionsof the Le Rozel footprints (n = 104) from the D3b-4 stratigraphic subunit



Fig. S18. Sizes of the footprints from the D3b-4 stratigraphic subunit at Le Rozel



Fig. S19. Length vs width of 35 footprints from the D3b-4 stratigraphic subunit

#### Text S7: Minimum Number of Individuals:

For the D3b-4 subunit, 39 footprints allowed for the measurement of length and 100 allowed for the measurement of width. In order to determine a Minimum Number of Individuals the 12.8% largest maximum intra-individual deviation obtained for the experimental footprint length. We considered that footprints falling within the interval  $[L\times(1-m.d.); L\times(1+m.d.)]$  of each other have been made by the same individual. The 39 footprints are associated with a minimum of 4 individuals according to 4 configurations (Fig. S20).

The experimental intraindividual maximum deviation in widths reaches 38,5% (which gave a MNI of 2), which is too high to allow any reliable estimation of the number of individuals using this method. However, in order to study their size distribution, based on the results obtained on the lengths (MNI = 4), the 100 width measurements were divided into 4 metric classes using the quartiles of their dispersion (4.5-12.8 cm).



Fig. S20. MNI estimate for the footprints from the D3b-4 stratigraphic subunit

#### Text S8: Estimation of statures:

For each footprint, stature was estimated from the length by achieving two methodological steps (Fig. S21). First of all, the variation between foot length and stature was quantified thanks to published osteometric data for *H. sapiens* (109-141). Then, footprint length was estimated from foot length using the mean ratio calculated in our experimental study (Footprint length / Foot length = 103.6%, r = 0.74, Table S4). A relationship between footprint length and stature is thus obtained.

The osteometric database (109-141, Table S8) encompasses different populations around the world, which enable us to take into account the differences in body proportions (*e.g.* 110, 142). These data also represent different age groups, from new born to adult (*e.g.* 112-113). The average values of foot length and stature were obtained either directly when they were available, or were estimated from published regressions when the foot length range is known and the correlation coefficient is significant enough (r > 0.60). An average foot length to stature ratio was calculated for each population sample and was used in order to calculate the global mean (14.8%) for all samples. The maximum (17.0%) and minimum (12.8%) ratios were also calculated.

Prehistoric footprints, such as those made at Le Rozel, were made by individuals whose feet are presumed to be morphologically different from those of current habitually shod populations (*e.g.* 98-99). Our database includes both shod and unshod individuals. In the literature, studies of body segment proportions between habitually shod and unshod groups usually involve samples from different regions (*e.g.* 137, 139-140). It is therefore difficult to know whether potential differences are related to footwear behavior or geographical differences. In our database, the habitually shod populations have an average foot length to stature ratio of about 15%, which does not differ from that of the habitually shod populations (*e.g.* 137, 139-140).

The application of the footprint length to foot length ratio enabled us to obtain an average linear relationship (Fig. S22) between stature (S) and footprint length (L):  $S = 6.51 \times L$ .

Furthermore, the maximum (17.6%) and minimum (13.2%) footprint length to stature ratios were determined in order to define the range of this ratio. The footprint length and stature data from our experimental study fall within this range (Fig. S22).



Fig. S21. Methods used in order to estimate statures from footprint lengths



**Fig. S22.** Relationship between footprint length and stature. Number of population samples: 38. Total number of individuals: 37,328. Using the data from refs. 109-141.

Number of			Habitually	Mean foot length to	
individuals	Age (year)	Geographical origin	Shod/Unshod	stature ratio	Reference
168	Mean: 22 y	India	Unshod	15.2%	137
38	[12-18 y]	Kenya	Unshod	14.6%	139
385	[6-18 y]	South Africa	Unshod	15.5%	140
338	[6-16 y]	USA and Northern Europe	Shod	15.5%	109
271	[28-50 y]	Northern America	Shod	14.8%	110
23	0 у	USA	Shod	15.2%	112
100	[17-24 y]	Africa	Shod	15.3%	112
100	[16-24 y]	India	Shod	15.0%	112
20	[1-18 y]	USA	Shod	14.9%	113
93	[0->55 y]	Alaska	Shod	15.1%	114
7119	[6-11 y]	USA	Shod	15.2%	115
250	[18-23 y]	Canada	Shod	14.8%	116
94	0 у	England	Shod	15.5%	117
8012	Mean: 23 y	USA	Shod	15.3%	118
270	[17-55 y]	India	Shod	14.8%	119
311	[19-71 y]	Turkey	Shod	14.3%	121
155	[17-23 y]	Turkey	Shod	14.8%	122
250	[18-30 y]	Mauritius	Shod	14.8%	123
246	[17-20 y]	India	Shod	14.6%	124
516	[18-83 y]	Turkey	Shod	15.0%	125
5093	[5-20 y]	Greece	Shod	15.0%	126
350	[18-50 y]	India	Shod	14.8%	129
256	0 у	Taiwan	Shod	14.5%	130
300	[18-30 y]	Nigeria	Shod	14.9%	131
7788	[1-13 y]	Germany	Shod	15.6%	133
61	Mean: 40 y	Caucasia	Shod	15.0%	134
160	[25-30 y]	Sudan	Shod	15.2%	135
190	Mean: 24 y	China	Shod	14.4%	137
250	[18-24 y]	Slovakia	Shod	14.5%	138
38	[12-18 y]	Kenya	Shod	14.9%	139
425	[6-18 y]	Germany	Shod	15.1%	140
478	[5-45 y]	Australia (indigeneous)	Unknown	15.4%	111
476	[6-18 y]	Japan	Unknown	14.6%	120
200	[18-80 y]	India	Unknown	14.9%	127
1040	[18-30 y]	India	Unknown	14.9%	128
154	[13-18 y]	India	Unknown	15.2%	132
1020	[19-42 y]	India	Unknown	14.1%	136
200	[18-30 y]	India	Unknown	14.7%	141

Table S8. Features of the published osteometric data used in order to estimate stature



Fig. S23. Foot length from 0 to 18 years old for different modern populations and position of the 39 complete footprints from the D3b-4 stratigraphic subunit after applying the experimental footprint length/foot length ratio (103.6%). Total number of individuals: 19,275.
 Using the data from refs. 109, 112-113, 115, 133, 143-144.

**Table S9.** Frequencies of the footprints from the D3b-4 stratigraphic subunit per age class determined using the curve representing variation between foot length and age (Fig. S23) for different modern populations. The frequencies for footprint width were obtained by estimating footprint length from the footprint width to length ratio (0.44)

		Children	Adolescents	Adults
		(0-10 years old)	(10-10 years old)	(> 10 years olu)
longth	Number of footprints	30	7	2
length	Relative frequency	76.9%	17.9%	5.1%
width	Number of footprints	72	20	9
wiath	Relative frequency	72.0%	20.0%	9.0%
Mean	Relative frequency	74.5%	19.0%	7.1%

Sito	Individual	Age		S	Stature (cm)		Boforoncoc	
Site	maiviauai	Estimates (year)	Methods	Estimates (cm)	Methods	Sex	References	
Amud	Amud 1	Adult	Osteological	177 cm	Regressions from femur	malo	145 140	
(Israel)	Amud 1	Adult	development	[174-180 cm]	measurements	male	145-149	
Cova Negra	Cours Norra	<b>F</b>	Osteological	95.5 cm	Regressions from femur		150 151	
(Spain)	Cova Negra	5 Y	development	[93-98 cm]	measurements		150-151	
	Dedaxingh 4	1.9 y	Tooth	84 cm	Regressions from long			
Dederiyeh	Dederiyen 1	[1.25-2.5 y]	development	[80-88 cm]	bones measurements		151 155	
(Syria)		2.2 y	Tooth	76 cm	Regressions from long		151-155	
	Dederiyen 2	[1.8-2.5 y]	development	[75-77 cm]	bones measurements			
	A shale for some la	A	Osteological	160	Regressions from	6		
	Adult female	Adult	development	160 cm	humerus measurements	remaie		
El Sidròn			Osteological		Regressions from ulna			
(Spain)	Adult male	Adult	development	167 cm	measurements	male	148, 156	
		7.5 y	Tooth	112.5 cm	Regressions from long			
	Juvenile 1	[7-8 y]	development	[105-120 cm]	bones measurements			
Fond-de-Forêt			Osteological	161.5 cm	Regressions from femur			
(Belgium)	Fond-de-Forët 1	Adult	development	[159-164 cm]	measurements	male	146-147, 149, 157	
Kebara			Osteological	171 cm	Regressions from long			
(Israel)	Kebara 2	Adult	development	[165-177 cm]	bones measurements	male	147-148, 158	
	Kiik-Koba 1		Osteological	teological 162 cm Regressions from tibia				
Kiik-Koba		Adult	development	[161-163 cm]	measurements	male		
(Ukraine)		0.5 y	Osteological	61 cm	Regressions from femur		147-148, 151, 159-162	
	Kiik-Koba 2	[0.4-0.6 y]	development	[58-64 cm]	measurements			
La Chapelle-aux-Saints			Osteological	164 cm	Regressions from			
(France)	La Chapelle-aux-Saints 1	Adult	development	[161-167 cm]	humerus measurements	male	146-148	
			Osteological	170.5 cm	Regressions from ulna			
	La Ferrassie 1	Adult	development	[168-173 cm]	measurements	male		
	La Ferrassie 2		Osteological	151.5 cm	Regressions from long			
		Adult	development	[146-157 cm]	bones measurements	female		
La Ferrassie			Osteological				146-148, 151, 161-163	
(France)	La Ferrassie 4	0.1 y	and tooth	52 cm	Regressions from femur			
		[0-0.2 y]	development	[51-53 cm]	measurements			
		4 v	Osteological	84 cm	Regressions from femur			
	La Ferrassie 6	(3-5 v)	development	[82-86 cm]	measurements			
La Quina			Osteological	157.5 cm	Regressions from ulna			
(France)	La Quina 5	Adult	development	[156-159 cm]	measurements	uncertain	147-148, 158, 164	
Le Moustier		0.2 y	Osteological	50.5 cm	Regressions from long			
(France)	Le Moustier 2	[0-0.3 y]	development	[49-52 cm]	bones measurements		151, 165-167	
Lezetxiki			Osteological		Regressions from			
(Spain)	Lezetxiki 1	Adult	development	166 cm	humerus measurements	uncertain	158	
		0 у	Osteological	52	Demostant f			
Mezmaïskaya	Mezmaïskaya	zmaïskaya [Fetus (0.6 v) -		52 cm	Regressions from femur		151, 168-169	
(Russia)	,-		Infant (0.2 y)]	development	[51-53 cm]	measurements		

# Table S10 (1/2). Estimations of stature, age and sex based on Neandertal osteological remains

C ito	Individual	Age			Stature		Poforoncoc
Site		Estimates (year)	Methods	Estimates (cm)	Methods	Sex	References
Neanderthal (Germany)	Neanderthal 1	Adult	Osteological development	165.5 cm [162-169 cm]	Regressions from long bones measurements	male	146-149
Regourdou (France)	Regourdou 1	Adult	Osteological development	164.5 cm [162-167 cm]	Regressions from humerus and radius measurements	uncertain	147-148, 170
Roc de Marsal (France)	Roc de Marsal 1	3.2 y [2.5-4 y]	Osteological development	83 cm [81-85 cm]	Regressions from femur measurements		151, 163, 171-172
Saint-Césaire (France)	Saint-Césaire 1	Adult	Osteological development	165 cm	Regressions from femur measurements	male	158
	Shanidar 1	Adult	Osteological development	168.5 cm [164-173 cm]	Regressions from long bones measurements	male	
	Shanidar 2	Adult	Osteological and tooth development	157 cm [154-160 cm]	Regressions from femur and tibia measurements	male	
	Shanidar 3	Adult	Osteological and tooth development	168 cm [166-170 cm]	Regressions from femur measurements	male	
Shanidar (Irak)	Shanidar 4	Adult	Osteological and tooth development	161.5 cm [158-165 cm]	Regressions from femur measurements	male	146-149, 151, 173-174
	Shanidar 5	Adult	Osteological and tooth development	165.5 cm [161-170 cm]	Regressions from femur measurements	male	
	Shanidar 6	Adult	Osteological and tooth development	147.5 cm [144-151 cm]	Regressions from femur measurements	female	
	Shanidar 10	1.5 y [1-2 y]	Osteological development	79 cm [78-80 cm]	Regressions from tibia measurements		
Sima de las Palomas	Palomas 92	Adult	Osteological development	152 cm	Regressions from femur measurements	uncertain	140
(Spain)	Palomas 96	Adult	Osteological development	154 cm	Regressions from femur measurements	uncertain	145
Ѕру	Spy 1	Adult	Osteological development	161 cm	Regressions from femur measurements	female	148 157-158
(Belgium)	Spy 2	Adult	Osteological development	158.5 cm [156-161 cm]	Regressions from femur and tibia measurements	male	140, 137-130
Tabun (Israel)	Tabun C1	Adult	Osteological development	155.5 cm [151-160 cm]	Regressions from long bones measurements	female	146, 148, 158

# Table S10 (2/2). Estimations of stature, age and sex from Neandertal osteological remains

 Table S11. Length of Neandertal femora and second metatarsals.

() Some femurs or second metatarsals are incomplete, their length was estimated from regressions.
 \*The femoral length of Kiik koba 1 was estimated from the tibial length and the mean Neandertal crural index (0.79, ref. 175)

C:to	ابه مانه داما د	Latavality	Maximal length	NAT/F	Defense	
Site	individu	Laterality	Second Metatarsal (MT)	Femur (F)		References
Amud	Amud 1	left	(8.0)	(48.4)	16.5%	176
Kiik-Koba	Kiik-Koba 1	left	7.6	(43.8)*	17.4%	159
	La Ferrassie 1	left	8.0	(46.5)	17.2%	177
La rell'assie	La Ferassie 2	left	(6.9)	41.1	16.7%	1//
Shanidar	Shanidar 1	right	(8.0)	(46.1)	17.2%	173
Sima de las Palomas	Palomas 92	left	7.2	39.7	18.1%	178
Tabun	Tabun C1	right	7.4	41.6	17.8%	179
	Mean		7.6	43.9	17.3%	

## Text S9: Three-Dimensional captures of Le Rozel footprints

A total of 180 tracks, including 169 hominin footprints, were digitized in 3D (Fig. S24). 133 were digitized by a scanner (Noomeo Optinum) that generates points clouds thanks to its high definition CCD cameras, and 70 by photogrammetry with Agisoft Photoscan (v.1.4.0) and a Canon EOS 1300D camera. 129 were digitized *in situ* and 55 were modeled from their casts (Fig. S25).





Fig. S25. Cast of a footprint

Fig. S24. 3D models of the footprint LREI2017-95 obtained by photogrammetry (A) and a scanner (B)

We tested potential differences between digitization techniques (scanner/photogrammetry) or the types of material (original footprints/casts) on their measurements. For both length and width, the average differences are always less than or equal to 0.1 cm (Fig. S26). Furthermore, ANOVA, realized after testing the normality of residuals and the homoscedasticity of variances, show no differences between acquisition techniques (Fig. S26, ANOVA: P >> 0.05). Data were then analyzed independently of the acquisition techniques. For measurements on the same footprint obtained for different acquisition techniques, the value that we considered is the average of the measurements on all 3D models of this footprint.



Fig. S26. Biometrical differences between digitization techniques (A) and the type of digitized material (B)

#### Supporting Information references:

1. R.H. Crompton, Making the case for possible hominin footprints from the Late Miocene (c. 5.7 Ma) of Crete? *Proc Geol Assoc* **128**, 692–693 (2017).

2. G.D. Gierliński *et al.*, Possible hominin footprints from the late Miocene (c. 5.7 Ma) of Crete? *Proc Geol Assoc* **128**, 697–710 (2017).

3. J. Meldrum, E. Sarmiento, Comments on possible Miocene hominin footprints. Proc Geol Assoc 129, 577-580 (2018).

4. R.J. Clarke, Early hominid footprints from Tanzania. S. Afr. J. Sci. 75, 148-149 (1979).

5. M.D. Leakey, R.L. Hay, Pliocene footprints in the Laetolil Beds at Laetoli, northern Tanzania. Nature 278, 317-323 (1979).

6. R.H. Crompton *et al.*, Human-like external function of the foot, and fully upright gait, confirmed in the 3.66 million year old Laetoli hominin footprints by topographic statistics, experimental footprint-formation and computer simulation. *J R Soc Interface* **9**, 707–719 (2011).

7. F.T. Masao *et al.*, New footprints from Laetoli (Tanzania) provide evidence for marked body size variation in early hominins. *Elife* **5**, e19568 (2016).

8. M.R. Bennett *et al.*, Early hominin foot morphology based on 1.5-million-year-old footprints from Ileret, Kenya. *Science* **323**, 1197–1201 (2009).

9. H.L. Dingwall, K.G. Hatala, R.E. Wunderlich, B.G. Richmond, Hominin stature, body mass, and walking speed estimates based on 1.5 million-year-old fossil footprints at Ileret, Kenya. J. Hum. Evol. 64, 556–568 (2013).

10. K.G. Hatala *et al.*, Footprints reveal direct evidence of group behavior and locomotion in Homo erectus. *Sci Rep* **6**, 28766 (2016).

11. N.T. Roach *et al.*, Pleistocene footprints show intensive use of lake margin habitats by Homo erectus groups. *Sci Rep* **6**, 26374 (2016).

12. K.G. Hatala *et al.*, Hominin track assemblages from Okote Member deposits near Ileret, Kenya, and their implications for understanding fossil hominin paleobiology at 1.5 Ma. *J. Hum. Evol.* **112**, 93–104 (2017).

13. A.K. Behrensmeyer, L.F. Laporte, Footprints of a Pleistocene hominid in northern Kenya. Nature 289,167-169 (1981).

14. F.H. Brown, B. Haileab, I. McDougall, Sequence of tuffs between the KBS Tuff and the Chari Tuff in the Turkana Basin, Kenya and Ethiopia. *J Geol Soc London* **163**, 185–204 (2006).

15. N. Ashton et al., Hominin footprints from early Pleistocene deposits at Happisburgh, UK. PLoS ONE 9, e88329 (2014).

16. F. Altamura *et al.*, Archaeology and ichnology at Gombore II-2, Melka Kunture, Ethiopia: everyday life of a mixed-age hominin group 700,000 years ago. *Sci Rep* **8**, 2815 (2018).

17. H. de Lumley, Les fouilles de Terra Amata à Nice. Premiers résultats. *Bulletin du Musée d'Anthropologie Préhistorique de Monaco* **13**, 29–51 (1966).

18. M.-A. de Lumley, P. Lamy, B. Mafart, "Une empreinte de pied humain acheuléen dans la dune littorale du site de Terra Amata. Ensemble stratigraphique C16" In *Terra Amata: Nice, Alpes-Maritimes, France. Tome II, Palynologie, anthracologie, faunes, mollusques, écologie et biogéomorphologie, paléoanthropologie, empreinte de pied humain, coprolithes,* H. de Lumley, Ed. (CNRS, Paris, 2011), pp. 483-507.

19. P. Mietto, M. Avanzini, G. Rolandi, Palaeontology: Human footprints in Pleistocene volcanic ash. Nature 422, 133 (2003).

20. M. Avanzini, P. Mietto, A. Panarello, M. De Angelis, G. Rolandi, The devil's trails: Middle Pleistocene human footprints preserved in a volcanoclastic deposit of southern Italy. *Ichnos : an international journal for plant and animal traces* **15**, 179–189 (2008).

21. S. Scaillet, G. Vita-Scaillet, H. Guillou, Oldest human footprints dated by Ar/Ar. Earth Planet. Sci. Lett. 275, 320–325 (2008).

22. A. Panarello, L. Santello, G. Farinaro, M.R. Bennett, P. Mietto, Walking along the oldest human fossil pathway (Roccamonfina volcano, Central Italy)? *J Archaeol Sci Rep* **13**, 476–490 (2017).

23. A. Tuffreau, Les fouilles du gisement paléolithique de Biache-Saint-Vaast (Pas-de-Calais) : années 1976 et 1977-premiers résultats. *Bulletin de l'Association française pour l'étude du quaternaire* **15**, 46–55 (1978).

24. A. Tuffreau, Les habitats du Paléolithique inférieur et moyen dans le Nord de la France (Nord, Pas-de-Calais, Somme). *Revue archéologique de Picardie* **1**, 91–104 (1988).

25. J.-J. Bahain *et al.*, ESR/U-series dating of faunal remains from the paleoanthropological site of Biache-Saint-Vaast (Pas-de-Calais, France). *Quat Geochronol* **30**, 541-546 (2015).

26. C.W. Helm, R. McCrea, D. Helm, A South African Pleistocene avian and mammal track site with purported prints of a shod hominid. *The Digging Stick* **29**, 17-20 (2012).

27. C.W. Helm, M.G. Lockley, K. Cole, T.D. Noakes, R.T. McCrea, Hominin tracks in southern Africa: A review and an approach to identification. *Palaeontologica Africana* 53:81-96 (2019).

28. S. Manolis, C.L. Aiello, R. Henessy, N. Kyparissi-Apostolika, "The Middle Palaeolithic Footprints from Theopetra Cave (Thessaly, Greece)" in *Theopetra Cave. Twelve years of excavation and research 1987–1998*, N. Kyparissi-Apostolika, Ed. (Greek Ministry of Culture and Institute for Aegean Prehistory, Athens, 2000), pp. 87-93.

29. H. Valladas *et al.*, TL age-estimates for the Middle Palaeolithic layers at Theopetra cave (Greece). *Quat Geochronol* **2**, 303–308 (2007).

30. D.L. Roberts, Last interglacial hominid and associated vertebrate fossil trackways in coastal eolianites, South Africa. *Ichnos* : an international journal for plant and animal traces **15**, 190–207 (2008).

31. Z. Jacobs, D.L. Roberts, Last Interglacial Age for aeolian and marine deposits and the Nahoon fossil human footprints, Southeast Coast of South Africa. *Quat Geochronol* **4**, 160–169 (2009).

32. D.L. Roberts, R.L. Berger, Last interglacial (c. 117 kyr) human footprints from South Africa. S. Afr. J. Sci. 93, 349–350 (1997).

33. I. Viehmann, Prehistoric Human Footprints in Romania's Caves. Theoretical and Applied Karstology 3, 229-234 (1987).

34. B.P. Onac *et al.*, U–Th ages constraining the Neanderthal footprint at Vârtop Cave, Romania. *Quat Sci Rev* **24**, 1151–1157 (2005).

35. K. Harvati, M. Roksandic, "The Human Fossil Record from Romania: Early Upper Paleolithic European Mandibles and Neanderthal Admixture" in *Paleoanthropology of the Balkans and Anatolia*, K. Harvati, M. Roksandic, Eds. (Springer, Netherlands, 2016), pp. 51–68.

36. C.W. Helm et al., A New Pleistocene Hominin Tracksite from the Cape South Coast, South Africa. Sci Rep 8, 3772 (2018).

37. P.R. Renne et al., Geochronology: Age of Mexican ash with alleged 'footprints'. Nature 438, E7 (2005).

38. S. González, D. Huddart, M.R. Bennett, A. González-Huesca, Human footprints in Central Mexico older than 40,000 years. *Quat Sci Rev* 25, 201–222 (2006).

39. D. Huddart, M.R. Bennett, S. González, X. Velay, Analysis and preservation of Pleistocene human and animal footprints: An example from Toluquilla, Valsequillo basin (Central Mexico). *Ichnos : an international journal for plant and animal traces* **15**, 232–245 (2008).

40. D.F. Mark, S. Gonzalez, D. Huddart, H. Böhnel, Dating of the Valsequillo volcanic deposits: Resolution of an ongoing archaeological controversy in Central Mexico. J. Hum. Evol. **58**, 441–445 (2010).

41. S.A. Morse, M.R. Bennett, S. Gonzalez, D. Huddart, Techniques for verifying human footprints: reappraisal of pre-Clovis footprints in Central Mexico. *Quat Sci Rev* **29**, 2571–2578 (2010).

42. P.G. Bahn, J. Vertut, Images of the Ice Age (Oxford University Press, Oxford, 1988).

43. S.P.M. Harrington, Human footprints at Chauvet cave. Archaeology 52, 18 (1999).

44. M.-A. Garcia, "Les empreintes et les traces humaines et animales" in *La Grotte Chauvet: l'art des Origines,* J. Clottes, Ed. (Seuil, Paris, 2001), pp. 34–43.

45. M.-A. Garcia, Ichnologie générale de la grotte Chauvet. Bulletin de la société préhistorique française 102, 103–108 (2005).

46. Quiles A *et al.*, A high-precision chronological model for the decorated Upper Paleolithic cave of Chauvet-Pont d'Arc, Ardèche, France. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 4670–4675 (2016).

47. T. Rusu, I. Viehmann, G. Racovita, V. Craciun, Primele urme de pasi ale omului preistoric din pesterile României (First footprints of prehistoric humans from Romanian caves, in Romanian in original). *Ocrotirea Nat* **13**, 191–200 (1969).

48. D. Webb, Ancient human footprints in Ciur-Izbuc Cave, Romania. Am. J. Phys. Anthropol. 155, 128–135 (2014).

49. F. Muñiz et al., Following the last Neanderthals: Mammal tracks in Late Pleistocene coastal dunes of Gibraltar (S Iberian Peninsula). Quat Sci Rev 217, 297-309 (2019).

50. J. Jaubert *et al.*, The chronology of human and animal presence in the decorated and sepulchral cave of Cussac (France). *Quat Int* **432**, 5–24 (2017).

51. L. Ledoux *et al.*, Traces of human and animal activity (TrAcs) in Cussac Cave (Le Buisson-de-Cadouin, Dordogne, France): Preliminary results and perspectives. *Quat Int* **430**, 141–154 (2017).

52. J.Y. Kim, K.S. Kim, C.Z. Lee, J.D. Lim, "Occurrence of hominid and other vertebrate footprints of Jeju Island, Korea" in *Proceedings of International Symposium on the Quaternary Footprints of Hominids and Other Vertebrates*, J.Y. Kim, K.S. Kim, S.I. Park, M.-K. Shin, Eds. (Namjejugun, Jeju, 2004), pp. 9–10.

53. D.-L. Cho, K.-H. Park, J.-H. Jin, W. Hong, Age Constraints on Human Footmarks in Hamori Formation, Jeiu Island, Korea. *The Journal of the Petrological Society of Korea* **14**, 149–156 (2005).

54. K.S. Kim, J.Y. Kim, S.H. Kim, C.Z. Lee, J.D. Lim, Preliminary report on hominid and other vertebrate footprints from the late Quaternary strata of Jeju Island, Korea. *Ichnos : an international journal for plant and animal traces* **16**, 1–11 (2009).

55. C.B. Kim, J.Y. Kim, K.S. Kim, H.S. Lim, New age constraints for hominid footprints found on Jeju Island, South Korea. J. Archaeol. Sci. **37**, 3338–3343 (2010).

56. Y.K. Sohn *et al.*, Stratigraphy and age of the human footprints-bearing strata in Jeju Island, Korea: Controversies and new findings. *J Archaeol Sci Rep* **4**, 264–275 (2015).

57. H. Duday, M.A. Garcia, Les empreintes de l'Homme préhistorique. La grotte du Pech-Merle à Cabrerets (Lot) : une relecture significative des traces de pieds humains. *Bulletin de la Société Préhistorique Française* **80**, 208–215 (1983).

58. A. Pastoors *et al.*, Tracking in caves: experience based reading of Pleistocene human footprints in French caves. *Cambridge Archaeological Journal* **25**, 551–564 (2015).

59. A. Pastoors et al., Experience based reading of Pleistocene human footprints in Pech-Merle. Quat Int 430, 155–162 (2017).

60. S. Webb, M.L. Cupper, R. Robins, Pleistocene human footprints from the Willandra Lakes, southeastern Australia. J. Hum. Evol. **50**, 405–413 (2006).

61. S. Webb, Further research of the Willandra Lakes fossil footprint site, southeastern Australia. *J Hum Evol* **52**, 711–715 (2007).

62. J. Ruiz, A. Torices, Humans running at stadiums and beaches and the accuracy of speed estimations from fossil trackways. *Ichnos : an international journal for plant and animal traces* **20**, 31–35 (2013).

63. D.D. Zhang, S.H. Li, Optical dating of Tibetan human hand-and footprints: An implication for the palaeoenvironment of the last glaciation of the Tibetan Plateau. *Geophys Res Lett* **29**, 16–1 (2002).

64. D.D. Zhang, S.H. Li, Y.Q. He, B.S. Li, Human settlement of the last glaciation on the Tibetan plateau. *Current Science* 84, 701–704 (2003).

65. K.G. Hatala et al., Early modern human footprints from Engare Sero, Tanzania. Am. J. Phys. Anthropol. 552, 158 (2011).

66. K.G. Hatala, B.G. Richmond, W.H. Harcourt-Smith, C.M. Liutkus, B. Zimmer, A snapshot of the anatomy, locomotion and social behavior of early modern humans as evidenced by fossil footprints at Engare Sero, Tanzania. *J Vertebr Paleontol* **32**, 107 (2012).

67. B. Zimmer, C. Liutkus, S. Carmichael, B. Richmond, S. Hewitt, A snapshot in time: Determining the age, environment, and social structures of early Homo sapiens using trace fossils in volcaniclastic rocks at the Engare Sero footprint site, Lake Natron, Tanzania. *Geological Soc of America Abstracts of 108th Annual Meeting* **44**, 13 (2012).

68. C.M. Liutkus-Pierce *et al.*, Radioisotopic age, formation, and preservation of Late Pleistocene human footprints at Engare Sero, Tanzania. *Palaeogeogr Palaeoclimatol Palaeoecol* **463**, 68–82 (2016).

69. C. Barriere, A. Sahly, Les empreintes humaines de Lascaux. Miscelanea en homenaje a Abate H Breuil 1, 173–180 (1964).

70. P.G. Bahn, The impact of direct dating on Palaeolithic cave art: Lascaux revisited. Anthropologie 33, 191–199 (1995).

71. E.L. Marcos, El descubrimiento de las Huellas Prehistoricas. Ojo Guarena. *Cubía Boletín Grupo Espeleológico Edelweiss* **3**, 34–35 (2001).

72. D. Bustos *et al.*, Footprints preserve terminal Pleistocene hunt? Human-sloth interactions in North America. *Sci Adv* **4**, eaar7621 (2018).

73. T.D. Dillehay, Monte Verde. Science 245, 1436 (1989).

74. D.J. Meltzer, On the Pleistocene antiquity of Monte Verde, southern Chile. Am Antiq 62, 659–663 (1997).

75. T.D. Dillehay *et al.*, Monte Verde: seaweed, food, medicine, and the peopling of South America. *Science* **320**, 784–786 (2008).

76. H. Bégouën, H. Vallois, "Étude des empreintes de pieds humains du Tuc d'Audoubert, de Cabrerets et de Ganties" in *Congrès International d'Anthropologie et d'Archéologie Préhistoriques* (Amsterdam, 1927), pp. 323–37.

77. R. Bégouën *et al.*, *Le sanctuaire secret des bisons: il y a 14 000 ans, dans la caverne du Tuc d'Audoubert* (Somogy Editions d'Art, Paris, 2009).

78. D. McLaren *et al.*, Terminal Pleistocene epoch human footprints from the Pacific coast of Canada. *PLoS ONE* **13**, e0193522 (2018).

79. L. Pales, M.T. De Saint-Péreuse, Les empreintes de pieds humains dans les cavernes: Les empreintes du réseau nord de la caverne de Niaux (Ariège) (Masson, Paris, 1976).

80. J. Clottes, H. Valladas, H. Cachier, M. Arnold, Des dates pour Niaux et Gargas. *Bulletin de la Société Préhistorique Française* 89, 270–274 (1992).

81. H. Valladas *et al.*, Direct radiocarbon dates for prehistoric paintings at the Altamira, El Castillo and Niaux caves. *Nature* **357**, 68 (1992).

82. M. Lockley, C. Meyer, Dinosaur tracks and other fossil footprints of Europe (Columbia University Press, New York, 2000).

83. V.G. Chiapella, Orsi e nomini preistorici nella grotta della 'Strega', (Genova). Revista del Comune A 29, 22-29 (1952).

84. L. Pales, Les empreintes de pieds humains de la Tana della Bàsura" (Toirano). Rivista di studi liguri 20, 5–12 (1954).

85. L. Pales, Les empreintes de pieds humains de la" grotta della Basura". Rivista di studi liguri 26, 25-90 (1960).

86. H. de Lumley, G. Vicino, New data concerning the dating and interpretation of human footprints present in the "Grotta della Basura" at Toirano (Savona, Northern Italy). Results of an international round table. *J. Hum. Evol.* **13**, 537–540 (1984).

87. P. Citton, M. Romano, I. Salvador, M. Avanzini, Reviewing the Upper Pleistocene human footprints from the 'Sala dei Misteri'in the Grotta della Bàsura (Toirano, northern Italy) cave: An integrated morphometric and morpho-classificatory approach. *Quat Sci Rev* **169**, 50–64 (2017).

88. J. Delteil, P. Durbas, L. Wahl, Présentation de la galerie ornée de Fontanet (Ornolac-Ussat-les-Bains, Ariège). Bulletin de la Société Préhistorique de l'Ariège 27, 01–10 (1971).

89. J. Clottes, R. Simonnet, Une datation radiocarbone dans la grotte ornée de Fontanet. Bulletin de la Société Préhistorique Française **71**, 106–107 (1974).

90. J. Clottes, Midi-Pyrénées. Gallia préhistoire 18, 613-650 (1975).

91. S.A. Aramayo, T.M. de Bianco, Late Quaternary palaeoichnological sites from the southern Atlantic coast of Buenos Aires Province, Argentina: mammal, bird and hominid evidence. *Ichnos : an international journal for plant and animal traces* **16**, 25–32 (2009).

92. C. Bayón, T. Manera, G. Politis, S. Aramayo, Following the tracks of the first South Americans. *Evolution: Education and Outreach* **4**, 205–217 (2011).

93. J.J. Scott, R.W. Renaut, R.B. Owen, Preservation and paleoenvironmental significance of a footprinted surface on the Sandai Plain, Lake Bogoria, Kenya Rift Valley. *Ichnos : an international journal for plant and animal traces* **15**, 208–231 (2008).

94. G. Guipert, M.-A. de Lumley, A. Tuffreau, B. Mafart, A late Middle Pleistocene hominid: Biache-Saint-Vaast 2, North France. *C R Palevol* **10**, 21–33 (2011).

95. Y. Facorellis, N. Kyparissi-Apostolika, Y. Maniatis, The cave of Theopetra, Kalambaka: radiocarbon evidence for 50,000 years of human presence. *Radiocarbon* **43**, 1029–1048 (2001).

96. B. van Vliet-Lanoë *et al.*, (2006) L'abri sous-roche du Rozel (France, Manche) : un habitat de la phase récente du Paléolithique moyen dans son contexte géomorphologique. *Quaternaire* **17**, 207–258.

97. N. Mercier, L. Martin, S. Kreutzer, V. Moineau, D. Cliquet, Dating the palaeolithic footprints of 'Le Rozel' (Normandy, France). *Quat Geochronol* **49**, 271-277 (2019).

98. K. Ashizawa, C. Kumakura, A. Kusumoto, S. Narasaki, Relative foot size and shape to general body size in Javanese, Filipinas and Japanese with special reference to habitual footwear types. *Ann. Hum. Biol.* **24**, 117-129 (1997).

99. K. D'Août, T.C. Pataky, D. De Clercq, P. Aerts, The effects of habitual footwear use: foot shape and function in native barefoot walkers. *Footwear Science* 1:81-94 (2009).

100. G. Roberts, S. Gonzalez, D. Huddart, Intertidal Holocene footprints and their archaeological significance. *Antiquity* **70**, 647-651 (1996).

101. D. Huddart, G. Roberts, S. Gonzalez, Holocene human and animal footprints and their relationships with coastal environmental change, Formby Point, NW England. *Quat Int* **55**, 29–41 (1999).

102. M.R. Bennett, S.A. Morse, Human footprints: fossilised locomotion? (Springer, Heidelberg, 2014).

103. L. Aiello, C. Dean, An introduction to human evolutionary anatomy (Academic Press, New York, 1990).

104. L. Klenerman, B. Wood, The human foot: a companion to clinical studies (Springer, Heidelberg, 2006).

105. R.H. Tuttle, Footprint clues in hominid evolution and forensics: lessons and limitations. *Ichnos : an international journal for plant and animal traces* **15**, 158-165 (2008).

106. M.R. Bennett, S. Gonzalez, D. Huddart, J. Kirby, E. Toole, Probable Neolithic footprints preserved in inter-tidal peat at Kenfig, South Wales (UK). *Proc Geol Assoc* **121**, 66–76 (2010).

107. Ø. Hammer, D.A.T. Harper, P.D. Ryan, PAST-palaeontological statistics, ver. 1.89. Palaeontol electron 4, 1–9 (2001).

108. J.C. Gower, Generalized procrustes analysis. Psychometrika 40, 33–51 (1975).

109. C.B. Davenport, The growth of the human foot. Am. J. Phys. Anthropol. 17, 167–211 (1932).

110. A. Hrdlička, The Pueblos. With comparative data on the bulk of the tribes of the Southwest and northern Mexico. *Am. J. Phys. Anthropol.* **20**, 235–460 (1935).

111. T.D. Campbell, J.H. Gray, C.J. Hackett, Physical anthropology of the Aborigines of Central Australia. *Oceania* **7**, 106–139 (1936).

112. H.V. Meredith, Human foot length from embryo to adult. Hum. Biol. 16, 207-282 (1944).

113. M. Anderson, M. Blais, W.T. Green, Growth of the normal foot during childhood and adolescence. Length of the foot and interrelations of foot, stature, and lower extremity as seen in serial records of children between 1–18 years of age. *Am. J. Phys. Anthropol.* **14**, 287–308 (1956).

114. P.L. Jamison, S.L. Zegura, An anthropometric study of the Eskimos of Wainwright, Alaska. *Arctic Anthropol* **7**, 125-143 (1970).

115. R.M. Malina, P.V. Hamill, S. Lemeshow, *Selected body measurements of children 6-11 years, United States* (US Government Printing Office, Washington, 1973).

116. H. Helmuth, Body height, foot size and the secular trend in growth. Z Morphol Anthropol 66, 31–42 (1974).

117. D.K. James, E.H. Dryburgh, M.L. Chiswick, Foot length–a new and potentially useful measurement in the neonate. *Arch. Dis. Child.* **54**, 226–230 (1979).

118. E. Giles, P.H. Vallandigham, Height estimation from foot and shoeprint length. J. Forensic Sci. 36, 1134–1151 (1991).

119. T.S. Singh, M.N. Phookan, Stature and footsize in four Thai communities of Assam, India. *Anthropologischer Anzeiger* **51**, 349-355 (1993).

120. K. Liu, K. Shinoda, T. Akiyoshi, H. Watanabe, Longitudinal analysis of adolescent growth of foot length and stature of children living in Ogi area of Japan: a 12 years data. *Z Morphol Anthropol* **82**, 87–101 (1998).

121. A. Özaslan, M.Y. İşcan, I. Özaslan, H. Tuğcu, S. Koç, Estimation of stature from body parts. *Forensic Sci. Int.* **132**, 40-45 (2003).

122. S.G. Sanli et al., Stature estimation based on hand length and foot length. Clin Anat 18, 589-596 (2005).

123. A.K. Agnihotri, B. Purwar, K. Googoolye, S. Agnihotri, N. Jeebun, Estimation of stature by foot length. *J Forensic Leg Med* 14, 279–283 (2007).

124. K. Krishan, A. Sharma, Estimation of stature from dimensions of hands and feet in a North Indian population. *J Forensic Leg Med* **14**, 327–332 (2007).

125. D. Atamturk, I. Duyar, Age-related factors in the relationship between foot measurements and living stature and body weight. *J. Forensic Sci.* **53**, 1296–1300 (2008).

126. T.B. Grivas, C. Mihas, A. Arapaki, E. Vasiliadis, Correlation of foot length and weight in school age children. *J Forensic Leg Med* **15**, 89-95 (2008).

127. T. Kanchan et al., Stature estimation from foot dimensions. Forensic Sci. Int. 179, 241-e1 (2008).

128. K. Krishan, Estimation of stature from footprint and foot outline dimensions in Gujjars of North India. *Forensic Sci. Int.* **175**, 93-101 (2008).

129. J. Sen, S. Ghosh, Estimation of stature from foot length and foot breadth among the Rajbanshi: an indigenous population of North Bengal. *Forensic Sci. Int.* **181**, 55-e1 (2008).

130. T.-Y. Ho *et al.*, Assessment of growth from foot length in Taiwanese neonates. *Pediatrics & Neonatology* **50**, 287–290 (2009).

131. E. Oa *et al.*, Stature estimation from Foot Dimensions of an Adult Nigerian Population. *Anatomica kamataka* **6**, 8-12 (2012).

132. K. Krishan, T. Kanchan, N. Passi, J.A. DiMaggio, Stature estimation from the lengths of the growing foot—A study on North Indian adolescents. *The Foot* **22**, 287–293 (2012).

133. S. Müller, A. Carlsohn, J. Müller, H. Baur, F. Mayer, Static and dynamic foot characteristics in children aged 1–13 years: a cross-sectional study. *Gait Posture* **35**, 389–394 (2012).

134. S. Reel, S. Rouse, W.V. Obe, P. Doherty, Estimation of stature from static and dynamic footprints. *Forensic Sci. Int.* **219**, 283-e1 (2012).

135. A.A. Ahmed, Estimation of stature using lower limb measurements in Sudanese Arabs. *J Forensic Leg Med* **20**, 483–488 (2013).

136. T.N. Moorthy, A.M.B. Mostapa, R. Boominathan, N. Raman, Stature estimation from footprint measurements in Indian Tamils by regression analysis. *Egyptian Journal of Forensic Sciences* **4**, 7-16 (2014).

137. Y. Shu, Q. Mei, J. Fernandez, Z. Li, N. Feng, Y. Gu, Foot morphological difference between habitually shod and unshod runners. *PLoS ONE* **10**, e0131385 (2015).

138. P. Uhrová et al., Estimation of stature using hand and foot dimensions in Slovak adults. Leg Med 17, 92-97 (2015).

139. H. Aibast *et al.*, Foot structure and function in habitually barefoot and shod adolescents in Kenya. *Curr Sports Med Rep* **16**, 448–458 (2017).

140. K. Hollander *et al.*, Growing-up (habitually) barefoot influences the development of foot and arch morphology in children and adolescents. *Sci Rep* **7**, 8079 (2017).

141. R.K. Shukla, A.S. Lodha, S. Das, Stature estimation from footprint: a study on central Indian population. *European Journal* of Forensic Sciences 4, 1-9 (2017).

142. P. Topinard, L'Anthropologie (C. Reinwald, Paris, 1876).

143. P.H. Dangerfield, C.J. Taylor, Anthropometric standards for term neonates. Early human development 8, 225–233 (1983).

144. J.R. Gohil, M. Sosi, S.N. Vani, A.B. Desai, Foot length measurement in the neonate. Indian J Pediatr 58, 675–677 (1991).

145. E. Hovers, Y. Rak, R. Lavi, W.H. Kimbel, Hominid remains from Amud Cave in the context of the Levantine Middle Paleolithic. *Paléorient* **21**, 47–61 (1995).

146. A. Thoma, The average stature of Neandertals. Z Morphol Anthropol 80, 195–198 (1995).

147. H. Helmuth, Body height, body mass and surface area of the Neandertals. Z Morphol Anthropol 82, 1–12 (1998).

148. J.-M. Carretero *et al.*, Stature estimation from complete long bones in the Middle Pleistocene humans from the Sima de los Huesos, Sierra de Atapuerca (Spain). *J. Hum. Evol.* **62**, 242–255 (2012).

149. S.E. Churchill, Thin on the ground: Neandertal biology, archeology and ecology (John Wiley & Sons, Hoboken, 2014).

150. J.L. Arsuaga et al., New Neandertal remains from Cova Negra (Valencia, Spain). J. Hum. Evol. 52, 31–58 (2007).

151. J.A. Martín-González, A. Mateos, I. Goikoetxea, W.R. Leonard, J. Rodríguez, Differences between Neandertal and modern human infant and child growth models. *J. Hum. Evol.* **63**, 140–149 (2012).

152. T. Akazawa et al., Neanderthal infant burial from the Dederiyeh Cave in Syria. Paléorient 21, 77-86 (1995).

153. T. Akazawa, S. Muhesen, H. Ishida, O. Kondo, C. Griggo, New Discovery of a Neanderthal Child Burial from the Dederiyeh Cave in Syria. *Paléorient* **25**, 129–142 (1999).

154. O. Kondo, Y. Dodo, T. Akazawa, S. Muhesen, Estimation of stature from the skeletal reconstruction of an immature Neandertal from Dederiyeh Cave, Syria. J. Hum. Evol. **38**, 457–473 (2000).

155. T. Akazawa, S. Muhesen, Neanderthal Burials: Excavations of the Dederiyeh Cave, Afrin, Syria: Studies in Honour of Hisashi Suzuki (KW publications, Auckland, 2003).

156. Rosas A *et al.*, The growth pattern of Neandertals, reconstructed from a juvenile skeleton from El Sidrón (Spain). *Science* **357**, 1282–1287 (2017).

157. E. Trinkaus, Sexual differences in Neanderthal limb bones. J. Hum. Evol. 9, 377–397 (1980).

158. C.B. Ruff, E. Trinkaus, T.W. Holliday, Body mass and encephalization in Pleistocene Homo. Nature 387, 173 (1997).

159. G.A. Bonch-Osmolovski, Grot Kiik-Koba (Paleolit Kryma, Moscow, 1940).

160. E. Vlček, Postcranial skeleton of a Neandertal child from Kiik-Koba, USSR. J. Hum. Evol. 2, 537–544 (1973).

161. J.-L. Heim, Les enfants néandertaliens de La Ferrassie. Étude anthropologique et analyse ontogénique des hommes de Néandertal (Masson, Paris, 1982).

162. R.L. Tompkins, E. Trinkaus, La Ferrassie 6 and the development of Neandertal pubic morphology. *Am. J. Phys. Anthropol.* **73**, 233–239 (1987).

163. J. Granat, J.-L. Heim, Nouvelle méthode d'estimation de l'âge dentaire des Néandertaliens. *L'anthropologie* **107**, 171–202 (2003).

164. E. Trinkaus, The sexual attribution of the La Quina 5 Neandertal. La diagnose sexuelle du Néandertalien La Quina 5. *Bull Mem Soc Anthropol Paris* 28, 111–117 (2016).

165. P. Sellier, A.-M. Tillier, J. Brůžek, A la recherche d'une référence pour l'estimation de l'âge des fœtus, nouveau-nés et nourrissons des populations archéologiques européennes. *Anthropologie et préhistoire* **108**, 75–87 (1997).

166. B. Maureille, La redécouverte du nouveau-né néandertalien Le Moustier 2. *PALEO Revue d'archéologie préhistorique* **14**, 221–238 (2002).

167. B. Maureille, Anthropology: A lost Neanderthal neonate found. Nature 419, 33 (2002).

168. L.V. Golovanova, J.F. Hoffecker, V.M. Kharitonov, G.P. Romanova, Mezmaiskaya cave: a Neanderthal occupation in the northern Caucasus. *Curr. Anthropol.* **40**, 77–86 (1999).

169. V. Barriel, A.-M. Tillier, L'enfant de Mezmaiskaya (Caucase) examiné dans une double perspective paléogénétique et paléoanthropologique. *Bull Mem Soc Anthropol Paris* **14**, 1–2 (2002).

170. V. Meyer, J. Bruzek, C. Couture, S. Madelaine, B. Maureille, Un nouveau bassin néandertalien. Description morphologique des restes pelviens de Regourdou 1 (Montignac, Dordogne, France). *PALEO Revue d'archéologie préhistorique* **22**, 207–222 (2011).

171. P. Legoux, Détermination de l'âge dentaire de l'enfant néandertalien du Roc-Marsal (Maloine, Paris, 1966).

172. M. Skinner, Dental wear in immature Late Pleistocene European hominines. J. Archaeol. Sci. 24, 677–700 (1997).

173. E. Trinkaus, The Shanidar Neandertals (Academic Press, New York, 1983).

174. L.W. Cowgill, E. Trinkaus, M.A. Zeder, Shanidar 10: A middle paleolithic immature distal lower limb from Shanidar cave, Iraqi Kurdistan. *J. Hum. Evol.* **53**, 213–223 (2007).

175. T.W. Holliday, Postcranial evidence of cold adaptation in European Neandertals. *Am. J. Phys. Anthropol.* **104**, 245-258 (1997).

176. J.R. Endo, T. Kimura, "Postcranial skeleton on the Amud Man" in *The Amud Man and his cave site*, H. Suzuki, F. Takai, Eds. (University of Tokyo Press, Tokyo, 1970).

177. J.-L. Heim, Les hommes fossiles de la Ferrassie. Vol. 2. Le squelettes adultes (squelette des membres) (Masson, Paris, 1982).

178. M.J. Walker, J. Ortega, K. Parmová, M.V. López, E. Trinkaus, Morphology, body proportions, and postcranial hypertrophy of a female Neandertal from the Sima de las Palomas, southeastern Spain. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 10087-10091 (2011).

179. T.D. McCown, A. Keith, *The Stone Age of Mount Carmel. Vol. 2. The Fossil Human Remains from the Levalloiso-Mousterian* (Clarendon Press, Oxford, 1939).