

Fossil record of the Icacinaceae and its paleogeographic implications

Cédric del Rio, Dario de Franceschi

▶ To cite this version:

Cédric del Rio, Dario de Franceschi. Fossil record of the Icacinaceae and its paleogeographic implications. Review of Palaeobotany and Palynology, 2020, 273, pp.104135. 10.1016/j.revpalbo.2019.104135. mnhn-02559005

HAL Id: mnhn-02559005 https://mnhn.hal.science/mnhn-02559005

Submitted on 21 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0034666719302258 Manuscript_9504a0477eb9a1d7ef6c0608763eab25

1	Fossil record of the Icacinaceae and its paleogeographic implications
2	
3	Cédric Del Rio ^{1,2,3*} , Dario De Franceschi ¹
4	
5	¹ CR2P - Centre de Recherche en Paléontologie – Paris. MNHN - Sorbonne Université –
6	CNRS, CP38, 57 rue Cuvier, 75231 Paris Cedex 05, France
7	
8	² ISYEB - Institut de Systématique, Évolution, Biodiversité, CNRS, MNHN, Sorbonne-
9	Université, EPHE, CP39, 57 rue Cuvier, 75231 Paris Cedex 05, France
10	
11	³ CAS, Key Laboratory of Tropical Forest Ecology, Chinese Academy of Sciences, Mengla,
12	Yunnan 666303, China
13	
14	*Corresponding author: cedric.del-rio@edu.mnhn.fr
15	

16 Abstract

17

In the past decades, the concept of Icacinaceae has been refined greatly, as morphological and 18 19 molecular data have led to a narrower circumscription of a monophyletic Icacinaceae family 20 with only 23 genera (vs 58 sensu Sleumer, 1942). This family possesses an extensive fossil record, important to the biogeographic history of the Northern Hemisphere, but the reported 21 fossils need to be carefully evaluated in the current phylogenetic framework. We evaluated 22 23 183 fossil reports of Icacinaceae from the literature but considered only 92 as reliably belonging to this family. Most of the accepted records are from endocarp remains. With this 24 25 sampling, we show an increase of the species richness during the Paleocene. A great increase of diversity in terms of genera, species, and morphological range is shown through the 26 Paleocene-Eocene interval and during the Early Eocene (Ypresian). Exchanges occurred 27 28 between North America and Europe near the PETM in both directions. During the middle and late Eocene, several of the modern genera appear first in the fossil record such as *Natsiatum*, 29 30 Phytocrene, and Pyrenacantha. Decreased diversity of post-Eocene records might be 31 explained by cooling during and subsequent to the Oligocene, which was less favourable to climbers. We observe the same pattern in other megathermal families showing the global 32 dynamic of megathermal groups of the North Hemisphere forest (boreotropical sensu. Wolfe, 33 1975) during the Paleogene. 34

35

36 Keywords: Icacinaceae, endocarp, leaves, pollen, biogeography, palaeobotany

39

60

In the past decades, the concept of Icacinaceae has been revised considerably with the 40 results of new molecular and morphological data that show that the former Icacinaceae family 41 concept with about 400 species and 58 genera (from Sleumer, 1942) was polyphyletic 42 (Kårehed, 2001). That phylogenetic study defined the Icacinaceae s.s. as belonging to the 43 44 Garryales order and placed some genera in the Aquifoliales and Apiales orders. In particular, 45 Kårehed described a new family, Stemonuraceae, and added some species to the Cardiopteridaceae family. The resulting circumscription of Icacinaceae included only 36 46 47 genera divided in four informal groups: Icacina, Cassinopsis, Emmotum, and Apodytes. Later, another study showed that this new circumscription of Icacinaceae corresponds to a 48 basal lamiid group (Lens et al., 2008). According to Lens et al. (2008), the Icacinaceae have 49 50 simple perforation plates in vessels (except for Cassinopsis), which is not the case for other 51 excluded former Icacinaceae. However, the Icacinaceae (sensu Lens et al., 2008) was still 52 considered as a polyphyletic group (Byng et al., 2014). More recently, a new Icacinaceae s.s. 53 family was delimited with only the Icacina group sensu Kårehed (2001) accepted, using a new molecular data set (Stull et al., 2015). Icacinaceae s.s. (sensu Stull et al. 2015) contains only 54 55 23 genera, with about 160 species, and is a basal lamiid group close to the Oncothecaceae 56 family. Both families are in the newly defined Icacinales order. Those taxonomic assignments 57 are accepted and used in the APG IV (APG, 2016). Thus, the Icacinaceae family s.l. belongs now to four indirectly related orders in Asterids group (Fig.1). 58 59 This family possesses an extensive fossil record that extends geographically and

Manchester, 1994; Collinson et al., 2012) but there are also reports of wood (e.g., Greguss,

geologically. The record is mainly composed of endocarps (e.g., Reid and Chandler, 1933;

62 1969), pollen (e.g., Kedves, 1970; Krutzsch & Vanhoorne, 1977; Cavagnetto, 2000), a flower

(Del Rio et al., 2017) and leaf remains (e.g., Wolfe, 1977; Tanai, 1990). However, most of the
paleobotanical discoveries and descriptions on Icacinaceae family were made before the
twenty first century. Therefore, assignments to Icacinaceae were done before the recent
modifications of the phylogenetic framework and some characters used to define Icacinaceae
can be in fact been simply apomorphic for Asterids (Fig. 1).

68 Some fossil species (e.g., Icacinoxylon alternipuncata, Icacinicarya papillaris, and 69 *Iodes germanica*), were used to calibrate an angiosperm phylogeny (Magallón et al., 2015). In 70 addition, the Icacinaceae fossil remains have been used as a paleoclimatic clue and contributed to the characterization of the Boreotropical forest definition, a vegetation type that 71 72 was widespread across middle latitudes in Northern Hemisphere during the Eocene (Wolfe, 1975). The Icacinaceae fossils also played an important role in the understanding of the 73 dispersion between Europe and North America during the early-middle Eocene (Manchester, 74 75 1994; Stull et al., 2011; Allen et al., 2015). Thus, it is crucial to evaluate the Icacinaceae fossil 76 record within a phylogenetic framework. Two reviews of the fossil record including the 77 asteridae were made (Martínez-Millán, 2010; Manchester, 2015), highlighting a Maastrichtian 78 minimum age for the Icacinaceae family. However, neither of these reviews embraces all the Icacinaceae fossils described until now. 79

In this study, we reviewed previously published reports of fossils attributed to the Icacinaceae. Our main questions are: (1) what is the fossil diversity of Icacinaceae s.s. following the new circumscription of the family and (2) what is the paleogeography and biogeographic history of this family based on this new circumscription?

84

85

2. Material and methods

We compiled all the occurrences of Icacinaceae in the literature to the extent of our 87 88 knowledge (Table 1.). The fossil record depends on available outcrops, search efforts, and findings. Here we took into consideration a majority of species from North America and 89 90 Europe. This distribution is probably due to a more thorough search effort in this area than others in palaeobotanical history (Morley & Dick, 2003). Indeed, Icacinaceae fossils were 91 recently found from Paleocene-Miocene range in South America (Stull et al., 2012). The 92 93 palaeobotanical studies of South-America have been increased during the last ten years 94 (Herrera et al., 2011, 2014; Jud et al., 2016; Pérez-Consuegra et al., 2017, 2018), and reveal a new diversity and new biogeographical hypotheses, including for Icacinaceae. We tabulated 95 96 183 fossil occurrences according to the age, locality, country and the concerned organs, among them 105 (81 species) were represented by fossil of endocarps divided in 12 fossil 97 genera: Comicilabium, Croomiocarpum, Faboidea, Hosiea, Icacinicarya, Icacinicaryites, 98 99 Iodicarpa, Palaeohosiea, Palaeophytocrene, Perforatocarpum, Sphaeriodes and Stizocarya 100 and four modern genera: Iodes, Natsiatum, Phytocrene and Pyrenacantha. Only one flower 101 has been described for this family, the extinct genus Icacinanthium. We also compiled 13 leaf 102 occurrences (10 species), five of them being from two extinct genera: Goweria and Huziokaeae and four of them from two extant genera: Merrilliodendron and Phytocrene. The 103 104 pollen record includes 26 occurrences (seven species) in two fossil genera, 105 Compositoipollenites and Echiperiporites, and only an extant ones Iodes. Finally, we 106 recorded 38 occurrences (14 species) of fossil wood records, with two fossil genera, *Icacinoxylon* and *Apodytoxylon*. The dominance in the fossil record of climbing genera, well 107 108 represented close to rivers, could be a taphonomic bias (Burnham, 1989; Burnham et al., 1992). However, because the climbing genera are well distributed in the family, we consider 109 110 our sample as representative of the entire family.

We decided to accept or reject occurrences based on the initial description and the associated figures in the literature. We checked all occurrences in accordance with new taxonomic and phylogenetic data (Stull et al. 2015). The Icacinaceae s.l. now belongs to four order in Asterids (Fig. 1), and thus some Icacinaceae characters used as criteria of fossil recognition (e.g., wood anatomy, leaves characters) cannot be longer used. We resumed the genera taxonomic changes and the associated fossil record in the Appendix 1.

We propose a reconstruction of the paleogeography of the Icacinaceae s.s. (Fig. 2) using paleomap made by Boucot et al. (2013). In this reconstruction (Fig. 2), the rejected occurrences are excluded (considered as definitively not Icacinaceae or without certainty in age of the outcrops) whereas the dubious occurrences are presented in grey (considered as

121 potential Icacinaceae, but without certainty). Accepted occurrences are in red (Fig. 2).

122 We used paleoclimatic reconstruction based on lithologic indicators in order to give a 123 brief and global idea of the link between past climate and Icacinaceae distribution (Boucot et al., 2013). This climatic ancestral reconstruction was based on the presence of lithological 124 125 remains: Coal, Bauxite and Laterite indicate a warm and wet climate (assumed as tropical), 126 the Coal and Tillites together indicate a cold and wet climate (cool temperate), the presence of Evaporite and Calcrete indicates a warm and dry climate (arid), The tillite associated with 127 128 Dropstone and Glendonite indicate a dry and cool climate (cold) and the presence of 129 Kaolinite, Crocodiles remains, palms and mangroves indicate a warm temperate climate. For more clarity, the position of the different rocks samples was removed and we only kept the 130 climatic area reconstructions. The complete lithological data can be found in the following 131 132 website: scotese.com/climate.htm. Despite more recent alternative data on reconstruction of biomes for parts of the world (e.g., South America, Jaramillo and Cardenas, 2013), we 133 134 decided to keep this global view as a good proxy for our study.

136 **3. Results**

137 **3.1 Species and occurrences rejected or dubious**

138 The endocarp record is the most meaningful for the Icacinaceae family, and probably139 the most reliable. However, some occurrences need to be revisited.

Comicilabium atkinsii was considered as belonging to Icacinaceae by its unilocular 140 endocarp, the vascular bundle on only one side and the cellular structure of the endocarp wall 141 (Manchester, 1994). The endocarp apical structure is a bulge with a lip. The size of the 142 endocarp and the wall thickness (2-4 mm) are surprising for a supposed Icacinaceae 143 144 compared with modern species endocarps (Del Rio, 2018). Despite the fact that the wall 145 anatomy, including interlocking digitate sclereids composing the endocarp and papillate 146 locule lining, is a convincing argument for the proposed affinity, we decided to consider this 147 species as dubious in order to consider the extreme differences in endocarp morphology. Icacinicarya mucronata, Icacinicarya bognorensis, Icacinicarya rotundata, and 148 Icacinicarya sp.11 are considered as dubious in a recent revision of the London Clay fossils 149 (Stull et al., 2016), because of the lack of some characteristic features and the bad 150 151 preservation of the specimens. We also followed this opinion here. Icacinicarya foveolata was 152 considered as a non-member of the Icacinaceae family in the same study because of the 153 presence of large hexagonal secretory cells lining the outside part of the seed and the thicker 154 endocarp wall. In the absence of alternative identification and another argument, we consider 155 that at least the characters given by Reid and Chandler (1933); i.e. endocarp unilocular and wall composed with partial digitate cells, are sufficient to not totally reject the potential 156 157 affinity with Icacinaceae family, so we consider it as dubious.

Icacinicarya amygdaloidea was considered as a member of Icacinaceae because of the shape, the position of the vascular bundle and the stylar canal (Chandler, 1961b). However, the specimens are smooth, which is uncommon within the Icacinaceae family (Del Rio, 2018). This trait is occasionally found in *Casimirella, Mappia* and especially in *Sarcostigma*. Additionally, the wall is composed of non-digitate cells, 10-12 μm in diameter. This kind of cells are never found in extant Icacinaceae (Del Rio 2018). Consequently, we rejected this species in our study.

Icacinicarya budvarensis from the Santonian of Germany (Knobloch & Mai, 1986) has a nodulose ornamentation, not common in the modern Icacinaceae family, and a shape also not found in extant species of Icacinaceae (Del Rio, 2018). Moreover, the description lacks the inner endocarp and the wall are partially not described. Therefore, we choose to consider this species as rejected.

Icacinicarya papillaris (Knobloch & Mai, 1986) resembles much more to an
Icacinaceae especially belonging to *Iodes*, but the preservation seems to be insufficient to
assign this species to a particular genus. We consider this species as dubious because of the
rough ornamentation and the absence of the papillae on the inner endocarp surface.

According to its original description, *Iodes germanica*, from the Maastrichtian of 174 175 Germany, have all the diagnostic characters of *Iodes* (reticulate ornamentation, papillae on the 176 inner surface, vascular bundle in one side on the endocarp wall; Knobloch & Mai, 1986). 177 Despite the lack of revision and a good illustration (Manchester, 2015), we decided to accept this last species as Icacinaceae member, based on the initial quite complete description. 178 179 Icacinicarya sp. (Chester, 1955) is rejected for this paleogeographical reconstruction because of the doubtful age of the Nigeria outcrops. Chester writes "Maastrichtian?" for this 180 181 specimen. Considering the importance of species from Cretaceous in paleogeographical

reconstructions, we decided to not take the risk to include this specimen, not formerlydescribed as a species by the author.

Icacinicarytes corruga (Pigg et al., 2008) lacks anatomical details and is based on a single unusually large specimen (38 mm x 28 mm). In addition, the morphology of this specimen seems to be difficult to interpret, in particular the ornamentation different from extant Icacinaceae. We consider it as dubious.

188 Icacinicaryites israelii from the Early Maastrichtian of Israel was a controversial species. Some authors disputed the attribution of this specimen to *Iodes* (Stull et al., 2016; 189 Del Rio et al., 2019a). In fact, this specimen lacks the anatomical detail needed to assign it. 190 191 The potential inner cotyledon discussed in the original article (Soudry & Gregor, 1997; figures 4, 8, 9) seems to be an artefact. The specimen on which this species is based is 192 193 bilateral, slightly compressed laterally, asymmetrical at the apex and reticulate on the surface. 194 All these characters could support an Icacinaceae assignment. However, the reticulation 195 pattern seems to be very orderly compared to Icacinaceae that have a more randomly arranged 196 reticulation (Del Rio, 2018). The lack of anatomical detail on the wall and locules 197 (Unilocular? Anatropous?) led us to consider this species as very questionable. Phytocrene microcarpa of Scott & Barghoorn 1957 is rejected by some authors (Stull 198 199 et al., 2012; Manchester et al., 2015). This species shows an anatomy that cannot match with 200 the Icacinaceae family (Manchester et al., 2015). Also, the shape and the size of the endocarp 201 do not match with this assignment. For this reason, we consider it as rejected.

cf. Natsiatum wilkoxiana from the Eocene of the Tallahatta Formation (Blanchard et
al., 2016) shows very few morphological clues and no anatomical characters. We cannot
exclude that this specimen corresponds to Icacinaceae, because of the ridge structures
described by the author. However, without enough diagnostic characters, we consider it as
dubious.

207 Some occurrences have been excluded from this study: an *Iodes* from the Miocene of 208 Yunnan (China) was mentioned by Stull et al. (2014) but not formerly published; one of us 209 (DDF) could observe another *Iodes* species from the Miocene of Turkey, but could not study 210 it in detail and the specimen is no longer accessible for study.

211 The leaves of modern Icacinaceae s.s. are in general simple, entire, mainly pinnatifid 212 with brochidodromous secondary venation and percurrent tertiary venation (Sleumer, 1971; Villiers, 1973; Utteridge et al., 2007; Allen et al., 2015; and personal observation). There are 213 214 some notable exceptions: leaves from Hosiea and Natsiatum are toothed with craspedodromous secondary veins; some Phytocrene and Pyrenacantha species have palmate 215 216 primary veins (Phytocrene borneensis, P. bracteata, P. palmata, P. hirsuta, Pyrenacantha malvifolia). Tertiary veins are occasionally more reticulate as some species in Casimirella, 217 Desmostachys, Miquelia, Pyrenacantha, and Stachyanthus in general. These examples show 218 219 the great variation in leaf architecture in the family Icacinaceae but also the limited variation 220 in the majority of species. Nevertheless, the classical type of leaves for Icacinaceae family is 221 common in tropical flowering plants, at least for the entire leaves' trait (Bailey & Sinnott, 222 1916; Wolfe, 1985). Thus, the leaves of this family lack diagnostic characters. While fossil leaves might appear identical in venation to modern leaves in the Icacinaceae, the fact that 223 overlapping venation occurs in other families means that we cannot be fully confident in the 224 225 identification of fossil leaves in this family. This probably explains the poor record of fossil 226 leaves attributed to this family (only 13 occurrences). We choose to consider the leaves as dubious if no clear diagnostic characters are given. 227

The genus *Goweria*, attributed to Icacinaceae by Wolfe (1977) and primarily to Menispermaceae (Wolfe, 1968), accommodates oval to lanceolate leaves with palmate primarily venation (in fact pinnatifid, see discussion in Allen et al., 2015), five primaries, brochydodromous secondary venation and percurrent tertiary veins. This diagnosis could

correspond to Icacinaceae family but not exclusively. Indeed, we found a similar type of 232 233 leaves for at least the Metteniusaceae family (Fig. 1) in some genera previously considered as Icacinaceae (Emmotum, Dendrobangia, Platea, Apodytes, Pittosporopsis, personal 234 235 observation). We decided to consider all Goweria species as dubious. Goweria linearis, in particular, seems to be very ambiguous (Wolfe, 1968). Actually, the specimen has a strongly 236 237 asymmetrical apex, which is uncommon in Icacinaceae species. In our opinion, Goweria 238 *bluerimensis* is the most convincing assignment (Allen et al., 2015). These specimens possess 239 expanded petioles with a narrow, resistant vascular strand that could be a diagnostic character. Huziokaea eoutilus partially resembles Icacinaceae, but is closer to Gomphandra 240 241 (Tanai, 1990), which is currently in Stemonuraceae (Kårehed, 2001). The assignment of this extinct genus to Icacinaceae s.s is here considered as wrong. 242

243 *Merrilliodendron ezoanum* identification seems to be based on traits (wide-elliptic 244 shape, intercostal tertiary veins, diverging perpendicular from the midvein), now shared by 245 several families including Icacinaceae and Stemonuraceae (Tanai, 1990); we consider it as 246 dubious.

247 The fossil record for *Phytocrene* leaves is composed of three species. However, Phytocrene acutissima from Eocene of Alaska corresponds to a partial fossil leaf (Wolfe, 248 1977; pl. 12, 1–6). *Phytocrene sordida* from Eocene of USA seems to be more convincing 249 250 (MacGinitie, 1941), but the main traits correspond to the common characters discussed above 251 for Goweria. The author of this species notes the similarity between the fossil species and modern Phytocrene blancoi and Hyperbaena hondurensis (Menispermaceae), showing the 252 253 ambiguity of the determination at the family level. Finally, Phytocrene ozakii from the 254 Eocene of Japan is based on a partial fossil (Tanai, 1990; pl. 6) and no diagnostic characters 255 were clearly used.

Both descriptions of *Pyrenacantha* sp. from Alaska and Japan (Wolfe, 1977; Tanai, 1990) are based on fragmentary material. They have mainly in common a pinnate venation and a weak pair of basal secondary venes that form marginal loops whose apical sides are oriented basally. This shape seems to appear many times in Asterids (e.g., Stemonuraceae, *Lasianthera*) and thus we consider these occurrences as dubious.

261 The genus *Icaciniphyllum* (not shown in Table 1.) was described for simple leaves, 262 with a non-entire margin (undulate, crenulate, and irregularly dentate) and with 263 craspedodromous venation (Kvaček & Bůžek, 1995). These characteristics are rare in Icacinaceae. In fact, only Hosiea and Natsiatum could roughly correspond to this description, 264 265 but the non-entire margin is very different. Other genera (e.g., *Iodes, Pyrenacantha...*) have entire margins and brochidodromous venation. Indeed, the type species Icaciniphyllum 266 artocarpites (Ettingshausen) Kvaček & Bůžek is now considered as a member of Sloanea L., 267 268 (Elaeocarpaceae), which invalidates the previous assignment (Kvaček et al., 2001; Allen et al., 2015). Thus, the affinity is probably wrong for all the species attributed to the genus. 269 270 To sum up, all fossil leaves are considered as dubious or as rejected for our 271 paleogeographical reconstruction. We are convinced that some occurrences (e.g., Goweria *bluerimensis*) are adequately documented; however, we show that the main traits of leaves 272 273 correspond mostly to the Icacinaceae s.l. are now shared by some more or less closely related 274 families.

The fossil record of potential icacinaceous pollen is poor in term of species diversity but rich in occurrences, especially in Europe. To our knowledge, only one pollen type was considered as "*Iodes* type" in the USA (Leopold & MacGinitie, 1972). This specimen was not represented nor described so we rejected this occurrence. The two other occurrences outside Europe are from the Miocene of Cameroon (Salard-Cheboldaeff, 1975). *Echiperiporites minor* is described as a periporate (5 pores) and echinulate pollen, which is apparently close to

Phytocrene pollen. However, a re-examination of the original material (KWI-1 lame B, 281 282 W35/W36) led us to recognize about ten pores (J. Dejax com. Pers.), which allow the rejection of this species, because no modern Icacinaceae pollen shows more than eight pores 283 284 (Lobreau-Callen, 1972). Echiperiporites icacinoides is described as three (4-5?) porate and echinulate pollen which could correspond to Stachyanthus, Deptaulus (Leptaulus?, 285 Aquifoliaceae) and (mostly) *Iodes*. The original material is decayed. We are waiting for its 286 287 potential restoration to eliminate the doubt and we decided here to reject this occurrence. Four species of the fossil pollen genus, Compositoipollenites, have been considered 288 related to the Icacinaceae family. Manchester (2015) considers that "a comprehensive 289 290 comparative investigation remains to be done with attention to other families that also share echinate pollen", in order to confidently conclude about the affinity of these pollen 291 292 specimens. We decided to follow the same caution and we used the dubious status for all 293 Compositoipollenites pollen. The wood attributed to Icacinaceae possesses a wide occurrence and stratigraphic 294 295 range (100.5–5.3 Ma). Unfortunately, they do not correspond to Icacinaceae s.s. and we 296 rejected them all. Indeed, Icacinaceae s.s. have simple perforation plates (Lens et al., 2008), whereas *Icacinoxylon* has scalariform perforation plates. The explanation of this paradoxical 297

nomenclature consists in the initial affinity of this fossil genus was made with *Citronella*,

which is now in Cardiopteridaceae (Fig.1, Appendix), (Stull et al., 2015). The other wood

300 taxon, Apodytoxylon hamamelidoides has Apodytes affinity, which is now in Metteniusaceae

301 family (Fig. 1, Appendix).

302 **3.2 Paleogeographic reconstruction**

We accepted 92 occurrences (mainly from endocarps), considered 45 occurrences as dubious (mainly leaves and pollen species) and rejected 46 occurrences (mainly from the wood species). According to our selection criteria, it has to be considered that all dubious species could be false identifications. We represent them in the following part in order to provide information about the potential range of the family, which could be a subject of discussion, only accepted species will be used in the biogeographic discussions.

309 The occurrences of Icacinaceae all belong to a warm temperate or boreotropical and 310 tropical climate, except for the Paleocene Egyptian occurrence, which apparently belongs to an arid climate (Icacinicarya youssefi). The Northern Hemisphere, in particular European and 311 312 North American areas, has almost all confident fossil occurrences. We have only one occurrence for the Cretaceous period and the most of the occurrences are from the early and 313 middle-late Eocene. The Oligocene record is very poor. In term of (accepted) species richness 314 315 (Table 2.), only one species belongs to the Cretaceous period, 14 are Paleocene, 59 are 316 Eocene (38 from the early Eocene and 24 from the middle-late) and only 3 species are from 317 the Oligocene epoch.

318 Records of Iodes extend from the Upper Cretaceous until the Oligocene. Phytocrene appears in the fossil record during the Paleocene or during the Eocene. Endocarp fossils of 319 320 Pyrenacantha and Natsiatum were found during the Eocene, but only Pyrenacantha is known 321 from the Oligocene. We did not identify other extant genera represented in the fossil record. 322 No extinct genus is present during the Upper Cretaceous: almost all extinct records are found in the Paleogene, mainly during the Eocene epoch. Removing the unnatural genera (based on 323 324 characters that allow us to assign specimens to Icacinaceae family, but not to a genus, e.g., Icacinicarya, Icacinicarytes), we can consider only one extinct genus during the Paleocene, 325 326 six genera from the Eocene and one genus from the Oligocene.

329 **4.1 Age of the family**

The age of Icacinaceae was estimated at 96.7 Ma with the help of molecular phylogeny 330 (Magallón et al., 2009). In another recent study, the estimated age was 104 Ma (Wikström et 331 al., 2015). This analysis used 17 fossil calibration points, but these are from various asterids, 332 no Icacinaceae fossil records were included. In another molecular study, the estimated age is 333 between 103 and 110 Ma, with the maximum likelihood method and between 65.5 and 100.3 334 Ma with a Bayesian analysis (Magallón et al., 2015). This last publication used numerous 335 336 fossils as calibration points and in particular three fossils attributed to Icacinaceae family: 337 *Icacinoxylon alternipuncata, Icacinicarya papillaris, and Iodes germanica.* The icacinaeous affinity of Icacinoxylon is rejected in this review, and Icacinicarya papillaris was considered 338 339 as dubious. Only *Iodes germanica* seems to be an unambiguous Icacinaceae fossil. The authors used 65.5 Ma as the age of calibration base on this record. Two reviews of the fossil 340 record of asteridae were made (Martínez-Millán, 2010; Manchester, 2015), highlighting a 341 342 Maastrichtian minimum age for the Icacinaceae family. In our review of the fossil record, 13 species belong to the Cretaceous epoch but only *Iodes germanica* is accepted. The minimum 343 344 appearance age selected for *Iodes* is 66 Ma (Upper Maastrichtian, following the International 345 chronostratigraphic chart v2017/02).

According to the previous studies, we consider the Upper Maastrichtian as the minimum age given by the fossil record for the Icacinaceae family. However, the calibration point for future molecular studies could be more precise including this minimum age at the internal node *Iodes-Mappianthus* (Stull et al., 2015) rather than at the Icacinaceae divergence.

351 **4.2 Major changes in the family Icacinaceae**

352 In our review, only *Iodes germanica* species already occurs during the Upper 353 Cretaceous of Europe, showing that the Icacinaceae family is present in Europe (Fig. 2A). 354 This species is also found during the Upper Paleocene, at least this species crosses the Cretaceous-Palaeogene boundary and persists during the Paleocene in the same area. This 355 could be a clue to the resilience of some families, and could explained by the previous results 356 357 based on the family level showing that the land plants did not suffer from global extinction 358 (but local turnover) during the Cretaceous-Palaeogene boundary (Cascales-Miñana & Cleal, 359 2014). In addition, we show an increase of occurrences, genera, and species for the Icacinaceae family during the Paleocene and new occurrences outside of Europe, mainly in 360 North America (Fig. 2B, Fig. 3), following the expansion of the boreotropical forest area 361 362 (Mai, 1989; Wolfe, 1975). The genus *Iodes* seems to be limited to Europe during the Upper 363 Cretaceous. With seven new *Iodes* species described for the Paleocene epoch, we show an increase of species richness. All species of *Iodes* from the Paleocene are very similar in term 364 of shape and only differ for some precise traits (size of the endocarp, reticular pattern, shape 365 of the ridges, Del Rio et al. 2019a). 366

We show a clear increase in Icacinaceae diversity during the early Eocene, in term of 367 species, new genera with new shapes, (Fig. 3). Among the 14 species present from the 368 369 Paleocene, only three species were found among the 38 early Eocene species. This shows a 370 turnover and an increase of diversity in the Icacinaceae family. Two genera were more 371 diverse, *Palaeophytocrene* and *Iodes*, which are probably climber species. This increase could be due to the new evergreen thermophilic flora, highly diversified and favourable to climbers, 372 373 flora that occurs during the PETM (Upchurch Jr & Wolfe, 1987; Collinson & Hooker, 2003) and rapid turnover but not catastrophic changes for plants (Cavagnetto, 2000; Wing et al., 374 2005; Collinson et al., 2009; Pigg & DeVore, 2009). In terrestrial diversity, studies show less 375

perturbation but migrations of the fauna probably following the flora change (Godinot & de
Broin, 2003; Gingerich, 2006).

The genus *Palaeophytocrene* only has one species from Bogota (South America) 378 379 during the Paleocene and eight species during the Eocene from North America and Europe. This could indicate a diversification and an expansion of the area of *Paleophytocrene* during 380 381 the PETM, following the expansion of the megathermal vegetation. Indeed, a geological 382 connection between North and South America for the *Paleophytocrene* was proposed during 383 the Paleocene (Stull et al., 2012). This connection became possible by the renewal of the magmatic arc of the Northern Andes during the Paleocene-Eocene (Bayona et al., 2011; 384 385 Cardona et al., 2011). Therefore, this genus shows a strong evidence of potential exchanges between tropical South America and boreotropical flora during this period. Changes during 386 387 the PETM in North America could be partially due to the flora spreading from South 388 America, followed by an expansion of these floras to Europe during the late Ypresian. New endocarp shapes appeared at the end of early Eocene in Europe, as Faboidea, 389 390 Perforatocarpum, Sphaeriodes, Stizocarya, all of these genera are now extinct. Among the 391 eight *Iodes* species from the Paleogene, three were found from the early Eocene showing a partial but important change in Icacinaceae floristic composition (Del Rio & De Franceschi, 392 393 in press). In particular, among the 14 species that occur during the early Eocene, four have no 394 horn-like protrusions at the apex (associated to the vascularisation of the fruit), whereas in the 395 Paleocene, all Iodes species have it (Del Rio et al. 2019a). Thus, we found new shapes and new endocarp types during the early Eocene. These new shapes could be due to local 396 397 diversifications or migrations of new floras from other areas. All species from the Paleocene are in Europe, whereas at least three species of *Iodes* (one with horn-like protrusions, two 398 399 without) are in North America during the early Eocene. The connection between North America and Europe during the Paleocene and early Eocene was well-documented (Wolfe, 400

1975; Manchester, 1994, 1999). This connexion remains possible via the North Atlantic land 401 402 bridge (McKenna, 1975; Tiffney, 1985) and following the extension of the boreotropical forest during the global warming. The migrating direction of exchanges (Asia-North America-403 404 Europe or Asia-Europe-North America) was intensively discussed in mammal palaeontology (Godinot & de Broin, 2003; Smith et al., 2006; Beard, 2008). Here, we have clues for an 405 exchange from Europe during the early Eocene. In fact, all species with or without horn-like 406 407 protrusions are older in Europe than in North America. However, an *Iodes multireticulata* specimen, dated from 55.5 Ma was found in the USA, the occurrence of this species in 408 Europe was dated to about 51 Ma (Reid & Chandler, 1933; Tiffney, 1999). Nevertheless, we 409 410 suspected that the specimen from the Fisher/Sullivan site can be a new species instead of a specimen of I. *multireticulata*. In fact, the size of the specimens from Fisher and London Clay 411 412 site seems distinctive. We show that the size of the endocarps could be used to differentiate 413 modern species when it exists a clear gap between the sizes within species (Del Rio, 2018). 414 This is here the case. With *Iodes* and *Paleophytocrene*, we have in consequence two examples 415 of interchanges between North America and Europe, showing that the dispersion of floras was 416 bilateral.

Due to particularly warm climate, the widest fossil extension of the Icacinaceae (in 417 418 term of species richness and probable area) was reached during the early Eocene. However, 419 this is from the middle-late Eocene that were found the fossil records attributed to modern 420 genera as Natsiatum (Stull et al., 2011), Phytocrene, at first in Europe and then in the USA (Stull et al., 2011; Collinson et al., 2012) and Pyrenacantha (Manchester, 1994). Therefore, it 421 422 is probably during the Eocene that the main modern genera appeared and diversified (Fig. 2D, Table 2). In *Iodes*, we show less species richness during the middle-late Eocene. These 423 424 "species richness regression" could be attributed to the cooling climate (Collinson et al., 1981; Keller, 1983; Zachos et al., 2008), what Tiffney (1985) called a "climatic 425

deterioration". This climate induced a change in floral structure, less favourable to the
climber species (Collinson & Hooker, 2003).

Only three occurrences of Icacinaceae were found during the Oligocene (Fig. 2E, Fig. 428 429 3B). This result shows an important decrease of diversity in the fossil record and is consistent with a regression of the boreotropical forest. At the Oligocene-Eocene boundary, the climate 430 431 was probably the coolest of the Paleogene and remained cold throughout the Oligocene 432 (Zachos et al., 2001, 2008). The boreotropical flora regressed and the deciduous vegetation, with less diversity, expanded (Tiffney, 1985). There are only two extant genera represented, 433 Pyrenacantha and Iodes and one extinct genus, Palaeophytocrene. The species Pyrenacantha 434 435 austroamericana is from the Oligocene of Peru (Stull et al., 2012). The oldest fossil remains of this genus is Pyrenacantha occidentalis of Lutetian age from Oregon, USA (Manchester, 436 437 1994). Thus, these two species show a connection between North and South America during 438 the Eocene. On the other side, the Oligocene *Iodes* species is from Europe. This presence of this species allows for the consideration of *Iodes* as being present in Europe throughout the 439 440 Palaeogene. This genus was probably part of a remnant of the boreotropical forest, also which 441 is almost completely replaced in Europe by cold-tolerant species during the Neogene. (Wolfe, 1975, 1985, Tiffney, 1985). In this way, the *Iodes* fossil from Miocene of Turkey (DDF obs.) 442 could also correspond to a late remain of this thermophilic European flora. Finally, 443 Palaeophytocrene sp. from Oregon, USA also shows a symmetrical pattern in North 444 445 American than *Iodes* in Europe (Meyer & Manchester, 1997).

446 **4.3. Comparison with the history of other angiosperm families**

447 Our results are in accordance with the boreotropical regression hypothesis (Lavin & Luckow,

448 1993). In particular, Wen (1999) establishes a disjunction of 65 genera between North

American and Asian floras, which occurred during the Miocene. This pattern could be
explained by an expansion of a boreotropical flora during the Paleocene and Eocene followed
by a high regression during the Neogene.

452 Some families, including Icacinaceae and Menispermaceae, were considered as closely associated with the boreotropical flora (Wolfe, 1975). These two families have in 453 common a majority of species with a climber habit and a similar present-day distribution. 454 455 Thus, we can hypothesize that both families could have a similar history linked with the 456 boreotropical expansion and regressions. A fossil history of the Menispermaceae was documented (Jacques, 2009). The Menispermaceae appeared during the Paleocene in the 457 458 fossil record (recorded from Cretaceous but dubiously), diversified during the Eocene and then regressed during the Oligocene in Europe and North America. In particular, the 459 maximum of diversity was found during the PETM event. A new species of Menispermaceae, 460 461 Sinomenium macrocarpum Liu et Jacques, was found after this review from the Miocene of North America (Liu & Jacques, 2010). However, this is probably a late refuge of the 462 463 boreotropical flora. New studies show Menispermaceae in South America during the 464 Paleocene (Herrera et al., 2011; Jud et al., 2018). These discoveries reveal a complex relationship between North and South America during the Paleocene. The same pattern is 465 466 observed for the Icacinaceae family during the Paleocene (Stull et al., 2011). In the Vitaceae, also a climber family, the oldest known fossil found is from India 467

before the continental collision (Manchester et al., 2013), followed by a rapid widespread
radiation during the Paleocene and Eocene in Northern Hemisphere. Other routes of
dispersion have been highlighted from South America (Manchester et al., 2012). In addition, a
study on the genus *Parthenocissus* (climber) shows a disjunction between Asia and North
America in the Early Miocene, which is congruent with the boreotropical regression (Nie et
al., 2010, 2012).

474	Some other evidence of expansion during the Paleogene and regression/extinction of
475	tropical floras in some areas (mainly North America and Europe) during the Neogene are
476	found for Paliurus (Rhamnaceae), Sabiaceae family, Canarium (Burseraceae), Deviacer (?
477	Polygalaceae), Ulmaceae, Carya (Juglandaceae), Sapindaceae and Anacardium
478	(Anacardiaceae), (McClain & Manchester, 2001; Manchester et al., 2007; Burge &
479	Manchester, 2008; Wang et al., 2013; Zhang et al., 2013; Chen & Manchester, 2015; Jia et al.,
480	2015; Han et al., 2018). Therefore, the results in the dynamic of flora on the Menispermaceae
481	and other families are very congruent with what we found in the Icacinaceae family.
482	

483 Conclusion

484

485 Among the 183 occurrences previously described as Icacinaceae, only 92 were accepted as 486 belonging to this family in this study and are mainly from endocarp remains. The minimum 487 age of the Icacinaceae family is probably the Upper Maastrichtian given by the fossil record 488 for the Icacinaceae family. With the accepted samplings, we show an increase of the species 489 richness during the Paleocene, probably due to the increase of temperature during the Thanetian. A great increase of diversity in term of genera, species, and morphology is shown 490 491 through the Paleocene-Eocene boundary and during the Ypresian. Exchanges were shown 492 between North America and Europe during the PETM in both ways. This is during the middle 493 and late Eocene that most of the modern genera appear in the fossil record as Natsiatum, Phytocrene, and Pyrenacantha. Finally, the cooling period during the Oligocene, less 494 495 favourable to climbers, could explain the decrease in the diversity of the Icacinaceae in the 496 fossil record. We show the same pattern of diversification and regression in other 497 megathermal families (Menispermaceae, Vitaceae...) showing a global dynamic of the North Hemisphere forest (boreotropical sensu. Wolfe, 1975) during the Paleogene. 498

499

500 Acknowledgments

502	This work was supported by a grant from Agence Nationale de la Recherche under the LabEx
503	ANR-10-LABX-0003-BCDiv, in the program "Investissements d'avenir" n_ANR-11-IDEX-
504	0004-02 and by the CR2P (UMR 7207).
505	
506	Author Contributions
507	DDF designed the project, CDR did the references work, CDR & DDF wrote the manuscript
508	and the discussion part.
509	
510	References
511	
512	Allen, S.E., Stull, G.W., Manchester, S.R., 2015. Icacinaceae from the Eocene of western
513	North America. Am. J. Bot. 102(5), 725-744.
514	Akkiraz, M.S., Kayseri, M.S., Akgün, F., 2008. Palaeoecology of coal-bearing Eocene
515	sediments in Central Anatolia (Turkey) based on quantitative palynological data. Turk.
516	J. Earth Sci. 17(2), 317–360.
517	APG, 2016. An update of the Angiosperm Phylogeny Group classification for the orders and
518	families of flowering plants: APG IV. Bot. J. Linn. Soc. 141(4), 399-436.
519	Bailey, I.W., Sinnott, E.W., 1916. The climatic distribution of certain types of angiosperm
520	leaves. Am. J. Bot. 3(1), 24–39.
521	Bayona, G., Montes, C., Cardona, A., Jaramillo, C., Ojeda, G., Valencia, V., Ayala-Calvo, C.,
522	2011. Intraplate subsidence and basin filling adjacent to an oceanic arc-continent
523	collision: a case from the southern Caribbean-South America plate margin: Intraplate

- subsidence and basin filling adjacent to an oceanic arc-continent collision. Basin Res.
 23(4), 403–422.
- Beard, K.C., 2008. The oldest North American primate and mammalian biogeography during
 the Paleocene–Eocene Thermal Maximum. Proc. Natl. Acad. Sci. 105(10), 3815–3818.
- 528 Blanchard, J., Wang, H., Dilcher, D. L., 2016. Fruits, seeds and flowers from the Bovay and
- 529 Bolden clay pits (early Eocene Tallahatta Formation, Claiborne Group), northern
- 530 Mississippi, USA. Palaeontologia Electronica, 19(3), 1-59.
- Boucot, A.J., Xu, C., Scotese, C.R., Morley, R.J., 2013. Phanerozoic paleoclimate: an atlas of
- 532 lithologic indicators of climate. SEPM (Society for Sedimentary Geology) 11, 1–30.
- 533 Bowerbank, J.-S., 1840. A history of the fossil fruits and seeds of the London Clay. The
- 534 Zoological society, London. 144 pp.
- Brown, R.W., 1962. Paleocene flora of the Rocky Mountains and Great Plains. US
 Government Printing Office, Washington. 375, 179 pp.
- Burge, D.O., Manchester, S.R., 2008. Fruit morphology, fossil history, and biogeography of *Paliurus* (Rhamnaceae). Int. J. Plant Sci. 169(8), 1066–1085.
- 539 Burnham, R.J., 1989. Relationships between standing vegetation and leaf litter in a
- 540 paratropical forest: implications for paleobotany. Rev. Palaeobot. Palynol., 58(1), 5–32.
- 541 Burnham, R.J., Wing, S.L., Parker, G.G., 1992. The reflection of deciduous forest
- 542 communities in leaf litter: implications for autochthonous litter assemblages from the
 543 fossil record. Paleobiology18(1), 30–49.
- 544 Byng, J.W., Bernardini, B., Joseph, J.A., Chase, M.W., Utteridge, T., 2014. Phylogenetic
- relationships of Icacinaceae focusing on the vining genera. Bot. J. Linn. Soc. 176(3),
 277–294.
- 547 Cardona, A., Valencia, V.A., Bayona, G., Duque, J., Ducea, M., Gehrels, G., Jaramillo, C.,
- 548 Montes, C., Ojeda, G., Ruiz, J., 2011. Early-subduction-related orogeny in the northern

- 549Andes: Turonian to Eocene magmatic and provenance record in the Santa Marta Massif
- and Rancheria Basin, northern Colombia: Palaeogene subduction in the northern Andes.

551 Terra Nova 23(1), 26–34.

- 552 Cascales-Miñana, B., Cleal, C.J., 2014. The plant fossil record reflects just two great
- extinction events. Terra Nova 26(3), 195–200.
- 554 Cavagnetto, C.G., 2000. La palynoflore d'un gisement d'ambre de l'Eocène basal du Bassin
- 555 Parisien (Le Quesnoy, France). Palaeontographica Abteilung B 255(4-6), 147–171.
- 556 Chandler, M.E.J., 1925. The Upper Eocene flora of Hordle, Hants, Palaeontographical
- 557 Society monographs, London. 52 pp.
- 558 Chandler, M.E.J., 1954. Some Upper Cretaceous and Eocene fruits from Egypt (with
- appendices by M.Y. Hassan & M. I. Youssef.) Br. Mus. Nat. Hist. Bull. Geol. 2, 149–
 187.
- 561 Chandler, M.E.J. 1961a. Plant remains of the Hengistbury and Barton Beds. British Museum
 562 of Natural History, London. 238 pp.
- 563 Chandler, M.E.J., 1961b. The Lower Tertiary floras of Southern England. I. British Museum
 564 of Natural History, London. 354 pp.
- Chandler, M.E.J., 1962. The Lower Tertiary floras of Southern England. II. Flora of the PipeClay Series of Dorset (Lower Bagshot). British Museum of Natural History, London.
 176 pp.
- 568 Chandler, M.E.J., 1964. The lower tertiary floras of Southern England IV. In: A Summary
- and Survey of Findings in the Light of Recent Botanical Observations. British Museum
 (Natural History), London. 151 pp.
- 571 Chandler, M.E.J., 1978. Supplement to the Lower Tertiary Floras of Southern England, Part
 572 5. Tertiary Research Special Paper 4, 1–47.

- 573 Chen, Y., Manchester, S.R., 2015. Winged Fruits of Deviacer in the Oligocene from the 574 Ningming Basin in Guangxi, South China. PloS one 10(12), e0144009
- 575 Chester, K., I., M., 1955. Some plant remains from the Upper Cretaceous and Tertiary of

576 West Africa. Ann Mag Nat Hist 8(91), 498–504.

- 577 Cleal, C.J., Thomas, B.A., Batten, D.J., Collinson, M., E., 2001. Mesozoic and Tertiary
- 578 paleobotany of Great Britain. Peterborough: Geological Conserv. Rev. Ser. 22, 335 pp.
- 579 Collinson, M.E., 1983. Fossil Plants of the London Clay. The Paleaeontological association,
 580 London. 121 pp.
- Collinson, M.E., Hooker, J.J., 2003. Paleogene vegetation of Eurasia: framework for
 mammalian faunas. Deinsea 10(1), 41–84.
- Collinson, M.E., Fowler, K., Boulter, M., 1981. Floristic changes indicate a cooling climate in
 the Eocene of southern England. Nature 291(5813), 315–317.
- 585 Collinson, M.E., Steart, D.C., Harrington, G.J., Hooker, J.J., Scott, A.C., Allen, L.O.,
- 586 Glasspool, I.J., Gibbons, S.J., 2009. Palynological evidence of vegetation dynamics in
- 587 response to palaeoenvironmental change across the onset of the Paleocene-Eocene

588 Thermal Maximum at Cobham, Southern England. Grana 48(1), 38–66.

- 589 Collinson, M.E., Manchester, S.R., Wilde, V., 2012. Fossil fruit and seeds of the Middle
- 590 Eocene Messel biota, Germany, Abh. Senckenb. Ges. Naturforsch, Stuttgart. 251 pp.
- 591 Crane, P.R., Manchester, S.R., Dilcher, D.L., 1990. A preliminary survey of fossil leaves and
- 592 well-preserved reproductive structures from the Sentinel Butte Formation (Paleocene)
- near Almont, North Dakota. Geology 20, 1–63.
- 594 Del Rio, C., 2018. Les Icacinaceae du Paléogène du Bassin de Paris. PhD diss. Muséum
- 595 national d'Histoire naturelle, Paris.

- 596 Del Rio, C., De Franceschi, D., Icacinaceae fossil fruits from three sites of the Paris Basin
 597 (early Eocene, France): local diversity and global biogeographic implications.
 598 Geodiversitas, In press (accepted 2019-06-25).
- Del Rio, C., Haevermans, T., De Franceschi, D., 2017. First record of an Icacinaceae Miers
 fossil flower from Le Quesnoy (Ypresian, France) amber. Scientific reports, 7(1),
- 601 **11099**.
- Del Rio, C., Thomas, R., De Franceschi, D., 2019a. Fruits of Icacinaceae Miers from the
 Palaeocene of the Paris Basin (Oise, France). Earth and Environmental Science
- Transactions of the Royal Society of Edinburgh, 108(4), 459-469.
- 605 Del Rio, C., Stull, G.W., De Franceschi, D., 2019b. New species of *Iodes* fruits (Icacinaceae)
- from the early Eocene Le Quesnoy locality, Oise, France. Review of Palaeobotany and
 Palynology. 262, 60-71.
- Duperon, J., 1976. A propos de quelques bois fossiles du Sud-Ouest de la France. Bull.
 Société Bot. Fr. 123(9), 533–540.
- 610 Fairon-Demaret, M., Smith, T., 2002. Fruits and seeds from the Tienen Formation at
- Dormaal, Palaeocene–Eocene transition in eastern Belgium. Rev. Palaeobot. Palynol.
 122(1-2), 47–62.
- Gingerich, P.D., 2006. Environment and evolution through the Paleocene–Eocene thermal
 maximum. Trends Ecol. Evol. 21(5), 246–253.
- 615 Godinot, M., de Broin, F. de L., 2003. Arguments for a mammalian and reptilian dispersal
- 616 from Asia to Europe during the Paleocene-Eocene boundary interval. Deinsea 10(1),
 617 255–276.
- 618 Gottwald, H.P., 2000. Gymnosperme und dicotyle Hölzer (67) aus dem" Aachener Sanden"
- 619 der Oberen Kreide von NO-Belgien und NW-Deutschland: mit anatomischer

- Bestimmung der Arten, Kommentaren, Tafeln und 96 Mikrofotos. Verlag Documenta
 Naturae 131, 1–44.
- 622 Grambast Fessard, N., 1969. Contribution à l'étude des flores tertiaires des régions
- provençales et alpines. V. Deux bois de dicotylédones à caractères primitifs du Miocène
 supérieur de Castellane. Nat. Monspel. 20, 105–118.
- 625 Greguss, P., 1969. Tertiary angiosperm woods in Hungary. Bp. Akad. Kiado. 151 pp.
- Gruas-Cavagnetto, C., Tambareau, Y., Villatte, J., 1992. Découverte de pollens, Dinoflagellés
 et Forminifères dans le Danien des Petites Pyrénées: Implications sur la position de La
- 628 Limite Crétacé/Tertiaire. Geobios 25, 19–28.
- Han, M., Manchester, S.R., Wu, Y., Jin, J., Quan, C., 2018. Fossil fruits of *Canarium*
- 630 (Burseraceae) from Eastern Asia and their implications for phytogeographical history. J.
- 631 Syst. Palaeontol. 16(10), 841–852.
- Heer, O., 1870. Die Miocene flora und fauna Spitzbergens. PA Norstedt & söner, Stockholm.
 98 pp.
- Herendeen, P.S., 1991. Charcoalified angiosperm wood from the Cretaceous of eastern North
 America and Europe. Rev. Palaeobot. Palynol. 70(3), 225–239.
- 636 Herrera, F., Manchester, S.R., Hoot, S.B., Wefferling, K.M., Carvalho, M.R., Jaramillo, C.,
- 637 2011. Phytogeographic implications of fossil endocarps of Menispermaceae from the
 638 Paleocene of Colombia. Am. J. Bot. 98(12), 2004–2017.
- 639 Herrera, F., Manchester, S.R., Carvalho, M.R., Jaramillo, C., Wing, S.L., 2014. Paleocene
- 640 wind-dispersed fruits and seeds from Colombia and their implications for early
- 641 Neotropical rainforests. Acta Palaeobot. 54(2), 197–229.
- 642 Hofmann, C.-C., Gregor, H.-J., 2018. Scanning electron microscope and light microscope
- 643 investigations of pollen from an atypical mid-Eocene coal facies in Stolzenbach mine

- 644 (Preu\s senElektra) near Borken (Kassel, Lower Hesse, Germany). Rev. Palaeobot.
 645 Palynol. 252, 41–63.
- 646 Hofmann, C.-C., Mohamed, O., Egger, H., 2011. A new terrestrial palynoflora from the
- 647 Palaeocene/Eocene boundary in the northwestern Tethyan realm (St. Pankraz, Austria).
- 648 Rev. Palaeobot. Palynol. 166(3-4), 295–310.
- 649 Hollick, C.A., 1936. The tertiary floras of Alaska. US Gov't. Print. Off. 182, 185 pp.
- Jacques, F.M.B., 2009. Fossil history of the Menispermaceae (Ranunculales). Ann. Paléontol.
 95(2), 53–69.
- Jaramillo, C., Cárdenas, A., 2013. Global warming and neotropical rainforests: a historical
- 653 perspective. Annual Review of Earth and Planetary Sciences, 41, 741–766.
- Jia, L.-B., Manchester, S.R., Su, T., Xing, Y.-W., Chen, W.-Y., Huang, Y.-J., Zhou, Z.-K.,
- 655 2015. First occurrence of *Cedrelospermum* (Ulmaceae) in Asia and its biogeographic
 656 implications. J. Plant Res. 128(5), 747–761.
- Jud, N.A., Nelson, C.W., Herrera, F., 2016. Fruits and wood of *Parinari* from the early
- Miocene of Panama and the fossil record of Chrysobalanaceae. Am. J. Bot. 103(2),
 277–289.
- Jud, N.A., Iglesias, A., Wilf, P., Gandolfo, M.A., 2018. Fossil moonseeds from the Paleogene
 of West Gondwana (Patagonia, Argentina). Am. J. Bot. 105(5), 927–942
- Kårehed, J., 2001. Multiple origin of the tropical forest tree family Icacinaceae. Am. J. Bot.
 88(12), 2259–2274.
- 664 Kedves, M., 1970. Études palynologiques des couches du Tertiaire inférieur de la région
- 665 Parisienne.V. Pollen triporés, subtriporés et intratriporés. Pollen & Spores 12(1), 83–97.
- 666 Kedves, M., Endredi, L., 1968. Palynologie investigations in the strata of «Buda Marl» with
- 667 plantremains. Acta Biol 14, 1–4.

- Kedves, M., Pradutz, A., 1970. Études palynologiques des couches du tertiaire inférieur de la
 région Parisienne. VI. Ultrastructure de quelques pollens d'Angiospermes de l'Éocène
 inférieur (Sparnacien). Pollen & Spores 12(4), 553–575.
- Kedves, M., Russell, D.E., 1982. Palynology of the Thanetian layers of Menat. The geology
 of the Menat Basin, France. Palaeontogr. Abt. B 182(4-6), 87–150.
- 673 Keller, G., 1983. Paleoclimatic analyses of middle Eocene through Oligocene planktic
- 674 foraminiferal faunas. Palaeogeogr. Palaeoclimatol. Palaeoecol. 43(1-2), 73–94.
- Knobloch, E., Mai, D.H., 1986. Monographie der Fruchte und Samen in der Kreide von
- 676 Mitteleuropa. Vydal Ustredni ustav geologicky, Praha. 219 pp.
- 677 Knobloch, E., Konzalova, M., 1998. Comparison of the Eocene plant assemblages of
- Bohemia (Czech Republic) and Saxony (Germany). Rev. Palaeobot. Palynol. 101(1-4),
 29–41.
- Krutzsch, W., Vanhoorne, R., 1977. Die Pollenflora von Epinois und Loksbergen in Belgien.
 Palaeontogr. Abt. B 163, 1–110.
- Kvaček, Z., Bůžek, Č., 1995. Endocarps and foliage of the flowering plant family Icacinaceae
 from the Tertiary of Europe. Tert. Res. 15(3), 121–138.
- Lavin, M., & Luckow, M, 1993. Origins and relationships of tropical North America in the
 context of the boreotropics hypothesis. Am. Jour. of Botany, 80(1), 1–14.
- Lens, F., Kårehed, J., Baas, P., Jansen, S., Rabaey, D., Huysmans, S., Hamann, T., Smets, E.,
- 687 2008. The wood anatomy of the polyphyletic Icacinaceae s.l., and their relationships
 688 within asterids. Taxon 57(2), 525–552.
- 689 Lenz, O.K., Wilde, V., Riegel, W., 2007. Recolonization of a Middle Eocene volcanic site:
- 690 quantitative palynology of the initial phase of the maar lake of Messel (Germany). Rev.
- 691 Palaeobot. Palynol. 145(3-4), 217–242.

- 692 Leopold, E.B., MacGinitie, H.D., 1972. Development and affinities of Tertiary floras in the
- Rocky Mountains. In A. Graham, ed., Floristics and paleofloristics of Asia and eastern
 North America. Elsevier Pubi. Co., Amsterdam. 147–200.
- Liu, Y.-S. (Christopher), Jacques, F.M.B., 2010. Sinomenium macrocarpum sp. nov.
- 696 (Menispermaceae) from the Miocene–Pliocene transition of Gray, northeast Tennessee,
- 697 USA. Rev. Palaeobot. Palynol. 159(1-2), 112–122.
- MacGinitie, H.D., 1941. Middle Eocene flora from the central Sierra Nevada. Publ. Carnegie
 Inst. Washington 534, 1-178.
- 700 MacGinitie, H.D., 1969. The Eocene Green River flora of northwestern Colorado and
- 701 northeastern Utah. University of California Press. 83, 1-202.
- Magallón, S., Castillo, A., 2009. Angiosperm diversification through time. Am. J. Bot. 96(1),
 349–365.
- Magallón, S., Gómez-Acevedo, S., Sánchez-Reyes, L.L., Hernández-Hernández, T., 2015. A
 metacalibrated time-tree documents the early rise of flowering plant phylogenetic
 diversity. New Phytol. 207(2), 437–453.
- Mai, D.H., 1987. Neue Fruchte und Samen aus palaozanen Ablagerungen Mitteleuropas.
 Feddes Repert. 98, 197–229.
- Mai, D.H., 1989. Development and regional differentiation of the European vegetation during

the Tertiary, in: Woody Plants evolution and Distribution since the Tertiary.
Springer, Vienna. 79–91.

- 712 Manchester, S.R., 1994. Fruits and seeds of the Middle Eocene nut beds flora, Clarno
- 713 Formation, Oregon, Palaeontolographica Americana. Paleontological Research
- 714 Institution, New York. 205 pp.
- 715 Manchester, S.R. 1999. Biogeographical relationships of North American tertiary floras.
- Annals of the Missouri Botanical Garden 86(2), 472–522.

717	Manchester, S.R., Wilde, V., Collinson, M.E., 2007. Fossil cashew nuts from the Eocene of
718	Europe: biogeographic links between Africa and South America. Int. J. Plant Sci.
719	168(8), 1199–1206.

- 720 Manchester, S.R., Chen, I., Lott, T.A., 2012. Seeds of Ampelocissus, Cissus, and Leea
- (Vitales) from the Paleogene of Western Peru and their biogeographic significance. Int.
 J. Plant Sci. 173(8), 933–943.
- Manchester, S.R., Kapgate, D.K., Wen, J., 2013. Oldest fruits of the grape family (Vitaceae)
 from the Late Cretaceous Deccan Cherts of India. Am. J. Bot. 100(9) 1849–1859.
- 725 Manchester, S.R., Grímsson, F., Zetter, R., 2015. Assessing the fossil record of Asterids in the
- 726 context of our current phylogenetic framework 1. Ann. Mo. Bot. Gard. 100(4), 329–
- 727 363.
- Martínez-Millán, M., 2010. Fossil record and age of the Asteridae. Bot. Rev. 76(1), 83–135.
- 729 McClain, A.M., Manchester, S.R., 2001. *Dipteronia* (Sapindaceae) from the tertiary of North
- America and implications for the phytogeographic history of the Aceroideae. Am. J.
- 731 Bot. 88(7), 1316–1325.
- 732 McKenna, M.C., 1975. Fossil mammals and early Eocene North Atlantic land continuity.
- 733 Ann. Mo. Bot. Gard. 62(2), 335–353.
- Meyer, H. W., Manchester, S. R., 1997. Oligocene Bridge Creek flora of the John Day
 Formation, Oregon. University of California Press 354 pp.
- Morley, R.J., Dick, C.W., 2003. Missing fossils, molecular clocks, and the origin of the
 Melastomataceae. Am. J. Bot. 90(11), 1638–1644.
- 757 Micrasionalaceae. Ani. J. Dol. 90(11), 1050–1044.
- 738 Nie, Z.-L., Sun, H., Chen, Z.-D., Meng, Y., Manchester, S.R., Wen, J., 2010. Molecular
- 739 phylogeny and biogeographic diversification of *Parthenocissus* (Vitaceae) disjunct
- between Asia and North America. Am. J. Bot. 97(8), 1342–1353.

- Nie, Z.-L., Sun, H., Manchester, S.R., Meng, Y., Luke, Q., Wen, J., 2012. Evolution of the
- 742 intercontinental disjunctions in six continents in the *Ampelopsis* clade of the grape
 743 family (Vitaceae). BMC Evol. Biol. 12(17), 1–13.
- 744 Pérez-Consuegra, N., Cuervo-Gómez, A., Martínez, C., Montes, C., Herrera, F., Madriñán, S.,
- Jaramillo, C., 2017. Paleogene *Salvinia* (Salviniaceae) from Colombia and their
- 746 paleobiogeographic implications. Rev. Palaeobot. Palynol. 246, 85–108.
- 747 Pérez-Consuegra, N., Góngora, D.E., Herrera, F., Jaramillo, C., Montes, C., Cuervo-Gómez,
- A.M., Hendy, A., Machado, A., Cárdenas, D., Bayona, G., 2018. New records of
- 749 Humiriaceae fossil fruits from the Oligocene and Early Miocene of the western Azuero
- 750 Peninsula, Panamá. Bol. Soc. Geológica Mex. 70(1), 223–239.
- Petrescu, J., Dragastan, O., 1972. Asupra unor lemne de Icacinoxylon Shilkina din oligocenul
 de la Telega-Prahova. Stud. Cerc Geol Geofiz Geog Ser Geol 17, 445–451.
- Petrescu, I., Nutu, A., 1972. Asupra unui lemn de Icacinoxylon Sh. in micenul superior de la
 Pravaleni. Brad. Sargetia Ser Sc 9, 77–79.
- 755 Petrescu, I., Codrea, V., 2004. The signification of the Paleocene microflora from Jibou (Sălaj
- 756 District) for the Paleogene palinological heritage of Romania. Acta Palaeontol.
- 757 Romaniae 4, 351–360.
- Petrescu, I., Ghiurca, V., Nica, V., 1989. Paleobotanical and palynological researches on the
- 759 lower-Oligocene amber and amber-bearing formation at Colti–Buzau. Oligocene
 760 Transylv. Basin 183–198.
- Pigg, K.B., Wher, W.C., 2002. Tertiary flowers, fruits, and seeds of Washington State and
 adjacent area-Part III. Washington Geology 30(3/4), 3–16.
- Pigg, K.B., DeVore, M.L., 2009. Floristic composition and variation in late Paleocene to early
 Eocene floras in North America. Bull. Geosci. 85(1), 135–154.

765	Pigg, K.B., Manchester, S.R., DeVore, M.L., 2008. Fruits of Icacinaceae (Tribe Iodeae) from
766	the Late Paleocene of western North America. Am. J. Bot. 95(7), 824–832.

- Potonié, R., 1960. Synopsis der Gattungen der Sporae dispersae: III. Beib. Geol. Jahrb. 39, 1-189.
- Potter, J.F., 1977. Eocene, lower Bracklesham Beds, iron workings in Surrey. Proc. Geol.
 Assoc. 88(4), 229–241.
- Rankin, B.D., Stockey, R.A., Beard, G., 2008. Fruits of Icacinaceae from the Eocene Appian
 Way Locality of Vancouver Island, British Columbia. Int. J. Plant Sci. 169(2), 305–
 314.
- Reid, E.M., Chandler, M.E.J., 1933. The London Clay Flora. The British Museum (Natural
 History), London. 561 pp.
- Salard-Cheboldaeff, M., 1975. Quelques grains de pollen périporés tertiaires du Cameroun.
 Rev. Micropaleontol. 17(4), 182–190.
- Scott, R.A., 1954. Fossil fruits and seeds from the Eocene Clarno Formation of Oregon.
 Palaeontogr. Abt. B 66–97.
- 780 Scott, R.A., Barghoorn, E.S., 1957. *Phytocrene microcarpa*-a new species of Icacinaceae
- based on Cretaceous fruits from Kreischerville, New York. The Paleobotanist. 25–28.
- Sleumer, H., 1942. Icacinaceae. In Engler, A. (ed) Die Natürlichen Pflanzenfamilien, Leipzig:
 Engelmann 20b, 322–396.
- 784 Sleumer, H., 1971. Icacinaceae, in: Flora Malesiana. CGGJ van Steenis, Noordhoff (ed),
- 785 Leiden. 7(1), 1–87.
- Smith, T., Rose, K.D., Gingerich, P.D., 2006. Rapid Asia–Europe–North America geographic
- 787 dispersal of earliest Eocene primate Teilhardina during the Paleocene–Eocene thermal
- 788 maximum. Proc. Natl. Acad. Sci. 103(30), 11223–11227.

Soudry, D., Gregor, H.-J., 1997. Jodes israelii sp. nov.: a huge phosphate-mineralized 789 790 icacinacean fructification from the Late Cretaceous of the Negev, southern Israel. Cretac. Res. 18(2), 161–178.

791

Stull, G.W., Moore, B.R., Manchester, S.R., 2011. Fruits of Icacinaceae from the Eocene of 792 southeastern North America and their biogeographic implications. Int. J. Plant Sci. 793

794 172(7), 935–947.

- Stull, G.W., Herrera, F., Manchester, S.R., Jaramillo, C., Tiffney, B.H., 2012. Fruits of an 795 796 "Old World" tribe (Phytocreneae; Icacinaceae) from the Paleogene of North and South America. Syst. Bot. 37(3), 784–794. 797
- Stull, G.W., Duno de Stefano, R., Soltis, D.E., Soltis, P.S., 2015. Resolving basal lamiid 798
- phylogeny and the circumscription of Icacinaceae with a plastome-scale data set. Am. J. 799 Bot. 102(11), 1794–1813. 800
- 801 Stull, G.W., Adams, N.F., Manchester, S.R., Sykes, D., Collinson, M.E., 2016. Revision of

Icacinaceae from the Early Eocene London Clay flora based on X-ray micro-CT. 802 803 Botany 94(9), 713–745.

- 804 Takahashi, K., Suzuki, M., 2003. Dicotyledonous fossil wood flora and early evolution of
- wood characters in the Cretaceous of Hokkaido, Japan. Iawa Journal 24(3), 269-309. 805
- Takhtajan, A., 1966. Major phytochoria of the Late Cretaceous and the Early Tertiary on the 806 807 territory of the USSR and adjacent countries. Bot. Zh. 51, 1217–1230.
- Tanai, T., 1990. Euphorbiaceae and Icacinaceae from the Paleogene of Hokkaido, Japan. Bull. 808 Natn. Sci. Mus. Tokyo Ser C. 16(3), 91-118. 809
- 810 Thayn, G.F., Tidwell, W.D., Stokes, W.L., 1985. Flora of the Lower Cretaceous Cedar
- 811 Mountain Formation of Utah and Colorado. Part III: Icacinoxylon pittiense n. sp. Am. J. 812 Bot. 72(2), 175.

- Thiele-Pfeiffer, H., 1988. Die Mikroflora aus dem mitteleozänen Ölschiefer von Messel bei
 Darmstadt. Palaeontogr. Abt. B 211(1-3), 1–86.
- Thomson, P.W., Pflug, H.D., 1953. Pollen und Sporen des mitteleuropäischen Tertiärs.
 Palaeontogr. Abt. B 1–138.
- Tichy, G., 1979. On the occurrence of fruits of Icacinaceae from the Eocene of St. Prankraz
- near Salzburg and the Kressenburg in Upper Bavaria. Verh Geol B-A 3: 415–421.
- 819 Tiffney, B.H., 1999. Fossil fruit and seed flora from the early Eocene fisher/sullivan site, in:
- 820 Early Eocene Vertebrates and Plants from the Fisher/Sullivan Site (Nanjemoy
- Formation) Stafford County, Virginia. Virginial Division of Mineral Ressources. 139–
 159.
- Tiffney, B.H., 1985. The Eocene North Atlantic land bridge: its importance in Tertiary and
- modern phytogeography of the Northern Hemisphere. J. Arnold Arbor. 66(2), 243–273.
- Tuzson, J., 1906. A balatoni fossilis fak monografiaja. [A monograph of the fossil plant
- remains from the Balaton region.]. Bal. Tud Tanulmanyozasanak Eredmenyei 1, 1–56.
- 827 Upchurch Jr, G., Wolfe, J., 1987. Mid-Cretaceous to Early Tertiary vegetation and climate:
- 828 evidence from fossil leaves and woods, in: The Origins of Angiosperms and Their
- Biological Consequences, EM Friis, WG Chaloner, and PR Crane (ed), England. 75–
- **830 105**.
- Utteridge, T., Nagamasu, H., Teo, S., White, L., Gasson, P., Brummitt, R., 2007. Icacinaceae.
 Flower. Plant Fam. World 173.
- Villiers, J.-F., 1973. Icacinaceae, in: Flore Du Cameroun. A. Aubréville et J-F. Leroy (ed),
 Paris, France. 3–100.
- 835 Von Ettingshausen, C., 1869. Die fossile flora des Tertiär-Beckens von Bilin. K. Gerold's
 836 Sohn. 110 pp.

- 837 Wang, Q., Manchester, S.R., Gregor, H.-J., Shen, S., Li, Z.-Y., 2013. Fruits of Koelreuteria
- (Sapindaceae) from the Cenozoic throughout the northern hemisphere: Their ecological,
 evolutionary, and biogeographic implications. Am. J. Bot. 100(2), 422–449.
- Wen, J., 1999. Evolution of eastern Asian and eastern North American disjunct distributions
 in flowering plants. Annu. Rev. Ecol. Syst. 30(1), 421–455.
- 842 Wheeler, E.A., Lee, M., Matten, L.C., 1987. Dicotyledonous woods from the Upper
- 843 Cretaceous of southern Illinois. Bot. J. Linn. Soc. 95(2), 77–100.
- 844 Wikström, N., Kainulainen, K., Razafimandimbison, S.G., Smedmark, J.E.E., Bremer, B.,
- 845 2015. A revised time tree of the asterids: establishing a temporal framework for
- evolutionary studies of the coffee family (Rubiaceae). PLoS ONE 10(5), 1–26,
- e0157206.
- 848 Wing, S.L., Harrington, G.J., Smith, F.A., Bloch, J.I., Boyer, D.M., Freeman, K.H., 2005.
- 849 Transient floral change and rapid global warming at the Paleocene-Eocene boundary.
 850 Science 310(5750), 993–995.
- Wolfe, J.A., 1968. Paleogene biostratigraphy of nonmarine rocks in King County,
- Washington. U.S. Geol. Surv.Prof. Paper 571.
- Wolfe, J.A., 1975. Some aspects of plant geography of the Northern Hemisphere during the
 late Cretaceous and Tertiary. Ann. Mo. Bot. Gard. 62(2), 264–279.
- Wolfe, J.A., 1977. Paleogene floras from the Gulf of Alaska region. U.S.GS. Prof. Paper 997.
- 856 Wolfe, J.A., 1985. Distribution of major vegetational types during the Tertiary. The carbon
- cycle and atmospheric CO2. Nat. Var. Archean Present 32, 357–375.
- Zachos, J.C., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and
 aberrations in global climate 64 Ma to Present. Science 292(5517), 686–693.
- 860 Zachos, J.C., Dickens, G.R., Zeebe, R.E., 2008. An early Cenozoic perspective on greenhouse
- warming and carbon-cycle dynamics. Nature 451(7176), 279–283.

- 862 Zhang, J.-B., Li, R.-Q., Xiang, X.-G., Manchester, S.R., Lin, L., Wang, W., Wen, J., Chen,
- 863 Z.-D., 2013. Integrated fossil and molecular data reveal the biogeographic
- 864 diversification of the Eastern Asian-Eastern North American disjunct Hickory Genus
- 865 (*Carya* Nutt.). PLoS ONE 8, e70449.
- 866

867 **Figure caption**

868

Figure 1. Phylogeny of Asterids following APG IV. In red: Position of the genera of
Icacinaceae s.l. (Sleumer, 1942). The asterix indicates the emplacement of the Icacinaceae s.s.
as circumscribed by Stull et al. (2015).

- Figure 2. Paleomaps showing the distribution of the Icacinaceae family in the fossil record.
- 873 (A) Upper Cretaceous, (B) Paleocene, (C) early Eocene, (D) middle-late Eocene and (E)
- 874 Oligocene. Red dots: accepted occurrence; grey squares: dubious occurrence. Paleomaps
- modified from Boucot et al., 2013.
- Figure 3. Diversity of Icacinaceae (A) Number of genera by ages, (B) Number of species by
- ages, (C) Number of *Iodes* species by ages. C: Cretaceous, P: Paleocene, E: Eocene, O:
 Oligocene.









Croomiocarpon missippiensis Stull, Manchester et Moore		USA	Minimum Age 43	Series Eocene	Stage Lutetian	StatusReferencesDubiousManchester, 1994
Faboidea crassicutis Bowerbank	Endocarp Tallahatta Fm., Claiborne Group (MI) Endocarp Herne bay, Sheppey, London Clay	USA United Kingdom	37 ≈ 51– 47,8	Eocene Eocene	Early-Middle Ypresian	Accepted Stull et al., 2011 Accepted Bowerbank, 1840; Reid & Chandler, 1933
Hosiea pterojugata Mai Icacinicarya mucronata Chandler	Endocarp Gonna, Abschnitt IV, graue Tone Endocarp Bognor, London Clay	Germany United Kingdom	56 47.8	Paleocene Eocene	Upper Ypresian	Accepted Mai 1987 Dubious Chandler, 1961b; Cleal et al., 2001
Icacinicarya amygdaloidea Chandler Icacinicarya bartonensis Chandler Icacinicarya becktonensis Chandler	Endocarp Herne Bay, London Clay Endocarp Headon bed, London Clay Endocarp Hordle Cliffs, London Clay	United Kingdom United Kingdom United Kingdom	≈ 51 41.3 37.8	Eocene Eocene Eocene	Ypresian Lutetian Bartonian	 Rejected Chandler, 1961b; Collinson, 1983; Cleal et al., 2001 Accepted Chandler, 1961a Accepted Chandler, 1961b
Icacinicarya bognorensis Reid and Chandler Icacinicarya budvarensis Knobloch and Mai	Endocarp Bognor, London Clay Endocarp České Bude [*] jovice Budvar	United Kingdom Germany	47.8 83.6	Eocene	Ypresian Santonian	DubiousReid & Chandler 1933; Cleal et al., 2001RejectedKnobloch & Mai, 1986
Icacinicarya collinsonae Pigg, Manchester et Devore Icacinicarya davisii (Chandler) Stull, Adams, Manchester et Collinson	Endocarp Almont (ND) Endocarp Suffolk, London Clay	USA United Kingdom	56.8	Paleocene	Tiffanian Ti3 Ypresian	Accepted Pigg et al., 2008 Accepted Chandler, 1961b; Collinson, 1983
Icacinicarya densipunctata Collinson, Manchester et Wilde Icacinicarya dictyota Pigg, Manchester et Devore	Endocarp Messel biota Endocarp Beicegel (ND)	Germany USA	≈47 56.8	Eocene Paleocene	Lutetian Tiffanian Ti3	Accepted Collinson et al., 2012 Accepted Pigg et al., 2008
Icacinicarya elegans (Bowerbank) Reid and Chandler Icacinicarya emarginata Chandler Icacinicarya forbesii Chandler	Endocarp Sheppey, London Clay Endocarp Sheppey London Clay	United Kingdom United Kingdom	47.8 47.8	Eocene Eocene	Ypresian Ypresian Ypresian	AcceptedReid & Chandler, 1933; Cleal et al., 2001AcceptedChandler, 1961b; Cleal et al., 2001AcceptedChandler, 1961b; Collinger, 1082
Icacinicarya foveolata Reid and Chandler Icacinicarya glabra Chandler	Endocarp Herne Bay, Sheppey, London Clay Endocarp Sheppey, London Clay	United Kingdom	~77.8 ≈ 51–47,8 47.8	Eocene	Ypresian Ypresian	 Cleal et al., 2001 Dubious Reid & Chandler, 1933; Collinson, 1983; Cleal et al., 2001 Accepted Chandler 1961, Cleal et al. 2001
Icacinicarya inornata Chandler Icacinicarya jenkinsi Reid and Chandler	Endocarp Lake, London Clay Endocarp Herne Bay, London Clay	United Kingdom United Kingdom	47.8 ≈51	Eocene Eocene	Ypresian Ypresian	Accepted Chandler, 1962; Cleal et al., 2001 Accepted Reid & Chandler, 1933; Cleal et al., 2001
Icacinicarya minima Reid and Chandler Icacinicarya nodulifera Reid and Chandler	Endocarp Bognor, Sheppey, London Clay Endocarp Herne Bay, Sheppey, London Clay	United Kingdom United Kingdom	47.8 ≈ 51–47,8	Eocene	Ypresian Ypresian	 Accepted Reid & Chandler, 1933; Collinson, 1983; Cleal et al., 2001 Accepted Reid & Chandler, 1933; Collinson, 1983; Cleal et al., 2001
Icacinicarya ovalis Ried and Chandler Icacinicarya ovoidea Reid and Chandler Icacinicarya papillaris Knobloch and Mai	Endocarp Sheppey, London Clay Endocarp Herne Bay, Sheppey, London Clay Endocarp Eisleben	United Kingdom United Kingdom Germanv	47.8 ≈51–47,8 66	Eocene Eocene Cretaceour	Ypresian Ypresian Maastrichian	Accepted Reid & Chandler, 1933Accepted Reid & Chandler, 1933; Cleal et al., 2001Dubious Knobloch & Mai. 1986
Icacinicarya platycarpa Reid and Chandler Icacinicarya pygmaea Chandler	Endocarp Herne Bay, Bognor, Sheppey, London Clay Endocarp Headon beds, London Clay	United Kingdom United Kingdom	00 ≈ 51–47,8 41.3	Eocene	Ypresian Lutetian	 Accepted Reid & Chandler, 1933; Chandler 1978; Collinson, 1983; Accepted Chandler, 1961a; Cleal et al., 2001
Icacinicarya reticulata Chandler Icacinicarya rotundata (Bowerbank) Reid and Chandler	Endocarp Herne Bay, Bognor, London Clay Endocarp Sheppey, London Clay	United Kingdom United Kingdom	≈ 51–47,8 47.8	Eocene Eocene	Ypresian Ypresian	Accepted Chandler, 1961b; Collinson, 1983; Cleal et al., 2001 Dubious Reid & Chandler, 1933; Cleal et al., 2001
Icacinicarya sp. Icacinicarya sp11	Endocarp Igala Division, Kabba Province Endocarp Sheppey, London Clay	Nigeria United Kingdom	66? 47.8	Cretaceous? Eocene	Maastrichian? Ypresian	Rejected Chesters, 1955 Dubious Reid & Chandler, 1933
Icacinicarya sp12 Icacinicarya youssefi Chandler Icacinicarvites corruga (Brown) Pice March (Endocarp Sheppey, London Clay Endocarp Gebel Atshai and Kosseir Area, Red Sea Endocarp Almy Animas EM Cont	United Kingdom Egypt USA	47.8 61.6	Eocene Paleocene Paleoce	Ypresian Danian ?	AcceptedReid & Chandler, 1933AcceptedChandler, 1954DubiousBrown 1962: Pigg et al. 2000
et Devore Icacinicaryites israelii (Soudry & Gregor) Del Rio, Thomas & De Franceschi Icacinicaryites linchensis Pigg, Manchester et Devore	Endocarp Linch, Sand Draw (WY),	Israel USA	about 72,1 56	Cretaceous	Early Maastrichtian Thanetian	Dubious Soudry & Gregor, 1997 Accepted Pigg et al., 2008
Icacinicaryites lottii Stull, S.E. Allen, & Manchester Icacinicarytes sp.	Animas Fm. (CO) Endocarp Kinsinger Lakes (WY) Endocarp Prémontré	USA France	48.5 49	Eocene	Ypresian Cuisian Sup.	Accepted Allen et al., 2015 Accepted Del Rio & De Franceschi in prep.
<i>Iodes acuta</i> Del Rio & De Franceschi <i>Iodes acutiformis</i> Chandler	Endocarp Le Quesnoy, Oise Endocarp Lake, Arne, London Clay	France United Kingdom	55-56 47.8	Eocene Eocene	Basal sparnatian Ypresian	Accepted Del Rio et al., 2019b Accepted Chandler, 1962; Cleal et al., 2001
<i>Iodes bilinica</i> (Ettingshausen) Stull, Adams, Manchester et Collinson <i>Iodes bilinica</i> (Ettingshausen) Stull, Adams, Manchester et Collinson	Endocarp Messel biota Endocarp Bohemia	Gremany Czech Republic	≈ 47 33.9	Eocene Eocene	Lutetian Upper	Accepted Collinson, 2012 Accepted Kvacek & Buzek, 1995
<i>Iodes bilinica</i> (Ettingshausen) Stull, Adams, Manchester et Collinson <i>Iodes brownii</i> (Berry) Stull, Adams, Manchester et Collinson <i>Iodes chandlerae</i> Manchester	 Endocarp Hordle Cliffs, London Clay Endocarp Tipperary flora (WY), Kisinger Lakes (WY), Blue Rim (WY), Bonanza (UT) Endocarp Clarno Nut Bods (OD) 	United Kingdom USA , USA	47.8 49,5 - 43,0	Eocene Eocene	Ypresian End Ypresian - Lutetian Lutetian	 Accepted Von Ettingshausen, 1866; Chandler, 1925; Takhtajan, 1966; Kvacek & Buzek, 1995 Accepted Manchester, 1994; Allen et al., 2015; Stull et al 2016 Accepted Manchester, 1994
Iodes corniculata Reid and Chandler Iodes eocenica Reid and Chandler	Endocarp Herne bay, Sheppey, London Clay Endocarp Sheppey, London Clay	United Kingdom United Kingdom	4.5 ≈ 51–47,8 47.8	Eocene	Ypresian Ypresian	AcceptedReid & Chandler, 1933; Chandler 1978; Collinson, 1983; Cleal et al., 2001AcceptedReid & Chandler, 1933; Chandler 1964;
<i>Iodes germanica</i> Knobloch and Mai <i>Iodes germanica</i> Knobloch and Mai	Endocarp Eisleben Endocarp Gonna	Germany Germany	66 56	Cretaceous Paleocene	Maastrichtian Upper	Cleal et al., 2001 Accepted Knobloch & Mai, 1986; Mai, 1987 Accepted Mai, 1987
Iodes hordwellensis ? Chandler Iodes multireticulata Reid and Chandler	Endocarp Hordle Cliffs, London Clay Endocarp Sheppey, Herne bay, London Clay	United Kingdom United Kingdom	37.8 ≈51–47,8	Eocene Eocene	Bartonian Ypresian	Accepted Chandler, 1961b Accepted Reid & Chandler, 1933; Cleal et al., 2001
Iodes multireticulata Reid and Chandler	Endocarp Clarno Nut Beds (OR) Endocarp Fisher/Sullivan Site (VI)	USA USA	43 55.5	Eocene Eocene	Lutetian Early	Accepted Manchester, 1994 Accepted Tiffney, 1999
Iodes occidentalis S.E. Allen, Stull, & Manchester Iodes parva Del Rio, Thomas & De Franceschi Iodes parva Del Rio, Thomas & De Franceschi	Endocarp Blue Rim (WY) Endocarp Petit Pâtis, Rivecourt, Oise Endocarp Le Quespeu Oise	USA France France	49.5 57-56	Eocene Paleocene	Ypresian Thanetian Basal space i	Accepted Allen et al., 2015 Accepted Del Rio et al., 2019a Accepted Del Rio et al. 2010b
Iodes sp. nov. Iodes reidii Del Rio, Thomas & De Franceschi	Endocarp Passy Endocarp Petit Pâtis, Rivecourt, Oise	France	57-56 s7-56	Eocene Paleocene	معمع sparnatian Middle sparnatian Thanetian	Accepted Del Rio et al., 2019b Accepted Del Rio et al., 2019a
<i>Iodes reidii</i> Del Rio, Thomas & De Franceschi <i>Iodes rigida</i> Del Rio & De Franceschi	Endocarp Passy Endocarp Le Quesnoy, Oise	France France	about 55 55-56	Eocene	middle sparnatian Basal sparnatian	AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio et al., 2019b
<i>Iodes rivecourtensis</i> Del Rio, Thomas & De Franceschi <i>Iodes rivecourtensis</i> Del Rio, Thomas & De Franceschi	Endocarp Petit Pâtis, Rivecourt, Oise Endocarp Petit Pâtis, Rivecourt, Oise	France France	57-56 57-56	Paleocene Paleocene	Thanetian Thanetian	Accepted Del Rio et al., 2019a Accepted Del Rio et al., 2019a
<i>Iodes sinuosa</i> Del Rio, Thomas & De Franceschi <i>Iodes</i> sp.	Endocarp Petit Pâtis, Rivecourt Endocarp Chuckanut Formation (WA)	France USA	57-56 33.9	Paleocene Eocene	Thanetian "Late"	Accepted Del Rio et al., 2019a Accepted Pigg & Wehr, 2002
Iodes sp.	Endocarp Bognor, London Clay Endocarp Grès de Belleu	United Kingdom France	47.8 49	Eocene	Ypresian Cuisian Sup.	Accepted Chandler, 1961b Accepted Del Rio & De Franceschi in prep.
Iodes sp. Iodes tubulifera Del Rio, Thomas & De Franceschi	Endocarp Dormaal Endocarp Petit Pâtis, Rivecourt, Oise Endocarp Paccy	Belgium France France	55 57-56	Eocene Paleocene	Ypresian Thanetian	Accepted Fairon-Demaret & Smith, 2002 Accepted Del Rio et al., 2019a
<i>Iodes tubulifera</i> Del Rio, Thomas & De Franceschi <i>Iodicarpa ampla</i> Manchester <i>Iodicarpa lenticularis</i> Manchester	Endocarp Passy Endocarp Clarno Nut Beds (OR) Endocarp Clarno Nut Beds (OR)	France USA USA	about 55 43 43	Eocene Eocene Eocene	middle sparnatian Lutetian Lutetian	Accepted Del Rio & De Franceschi in prep. Accepted Manchester, 1994 Accepted Manchester, 1994
Natsiatum wilcoxiana (Bery) Stull, Moore & Manchester (cf) Natsiatum wilcoxiana (Bery) Stull, Moore & Manchester	Endocarp Fayette County, Wilbanks II Clay Pits (TE) Endocarp Tallahatta Fm., Claiborne Group (MI)	USA USA	43 37.8 37	Eocene	Middle	Accepted Stull et al., 2011 Dubious Blanchard et al., 2016
Palaeohosiea marchiaca (Mai) Kvacek & Buzek Palaeohosiea marchiaca (Mai) Kvacek & Buzek	Endocarp Bohemia Endocarp Shepey, Bognor, London Clay	Czech Republic United Kingdom	59.2 47.8	Paleocene	Middle Lower	Accepted Mai, 1987; Kvacek & Buzek 1995 Accepted Kvacek & Buzek, 1995
Palaeohosiea suleticensis (Mai) Kvacek & Buzek Palaeophytocrene ambigua Reid and Chandler	Endocarp Suletice, Bohemia Endocarp Bognor, Sheppey, London Clay	Czech Republic United Kingdom	28.1 47.8	Oligocene Eocene	Lower Ypresian	Accepted Kvacek & Buzek, 1995 Accepted Read & Chandler, 1933
Palaeophytocrene foveolata Reid and Chandler Palaeophytocrene foveolata var. minima Reid and Chandler	Endocarp Herne Bay, Bognor, Sheppey, London Clay Endocarp Herne Bay, Bognor, Sheppey, London Clay	United Kingdom United Kingdom	$\approx 51-47,8$ $\approx 51-47,8$	Eocene Eocene	Ypresian Ypresian	Accepted Read & Chandler, 1933 Accepted Reid & Chandler, 1933
Palaeophytocrene hammenii Stull	Endocarp Bogota fm. Checua clay pit Endocarp Clarno Nut Beds (OR)	Colombia USA	58 43	Paleocene Eocene	Middle-Late Lutetian	Accepted Stull et al., 2012 Accepted Scott, 1954; Manchester, 1994
		TICA	27.9	Eocene	Middle	Accepted MacGinitie, 1941; MacGinitie, 1969
Palaeophytocrene kressenbergensis (Schafhäutl)	Endocarp Independence Hill (SN) Endocarp St, Pankraz, Bavaria	Germany	33.9-56	Eocene		Accepted Tichy, 1979
Palaeophytocrene hancocku (Scou) Manchester Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäutl) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester	Endocarp Independence Hill (SN) Endocarp St, Pankraz, Bavaria Endocarp Vancouver island Endocarp Clarno Nut Beds (OR)	Germany Canada USA	33.9-56 33.9 43	Eocene Eocene	? Lutetian	Accepted Tichy, 1979 Accepted Rankin et al., 2008 Accepted Scott, 1954; Manchester, 1994
 Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäutl) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. Palaeophytocrene sp. Palaeophytocrene sp. 	 Endocarp Independence Hill (SN) Endocarp St, Pankraz, Bavaria Endocarp Vancouver island Endocarp Clarno Nut Beds (OR) Endocarp Ronald, Kittitas County (WA) Endocarp Almont (ND) Endocarp John Day Formation (Or) 	Germany Canada USA USA USA USA	37.8 33.9-56 33.9 43 33.9 56 32.2.22.5	Eocene Eocene Eocene Paleocene Oligooor	? Lutetian Late	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMever & Manchester, 1007
 Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäutl) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. 	EndocarpIndependence Hill (SN)EndocarpSt, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpSt, Pankraz, BavariaEndocarpGrès de Belleu	Germany Canada USA USA USA USA Germany France	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49	Eocene Eocene Eocene Paleocene Oligocene Eocene	? Lutetian Late Cuisian Sup.	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.
 Palaeophytocrene hancocku (SCOU) Manchester Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäutl) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. Palaeop	EndocarpIndependence Hill (SN)EndocarpSt, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpSt, Pankraz, BavariaEndocarpGrès de BelleuEndocarpVancouver islandEndocarpYakutat Frm.	Germany Canada USA USA USA USA Germany France Canada Alaska	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9	Eocene Eocene Eocene Paleocene Oligocene Eocene Eocene Eocene	 ? Lutetian Late Cuisian Sup. ? Middle-Late 	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870
 Palae ophytocrene lacunosum (MacGinitie) Wolfe Palae ophytocrene kressenbergensis (Schafhäutl) Palae ophytocrene manchesteri Rankin, Stockey et Beard Palae ophytocrene pseudopersica Scott emend Manchester Palae ophytocrene sp. <l< td=""><td>EndocarpIndependence Hill (SN)EndocarpSt, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpJohn Oay Formation (Or)EndocarpSt, Pankraz, BavariaEndocarpGrès de BelleuEndocarpVancouver islandEndocarpYakuta Frm.EndocarpSteppey, London ClayEndocarpWilbanks II Clay Pits (TE)</td><td>Germany Canada USA USA USA USA Germany France Canada Alaska United Kingdom</td><td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9 47.8 37.8</td><td>Eocene Eocene Eocene Paleocene Oligocene Eocene Eocene Eocene Eocene</td><td> ? Lutetian Late Cuisian Sup. ? Middle-Late Ypresian Bartonian </td><td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Stull et al., 2011</td></l<>	EndocarpIndependence Hill (SN)EndocarpSt, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpJohn Oay Formation (Or)EndocarpSt, Pankraz, BavariaEndocarpGrès de BelleuEndocarpVancouver islandEndocarpYakuta Frm.EndocarpSteppey, London ClayEndocarpWilbanks II Clay Pits (TE)	Germany Canada USA USA USA USA Germany France Canada Alaska United Kingdom	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9 47.8 37.8	Eocene Eocene Eocene Paleocene Oligocene Eocene Eocene Eocene Eocene	 ? Lutetian Late Cuisian Sup. ? Middle-Late Ypresian Bartonian 	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Stull et al., 2011
 Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäutl) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. Phytocrene sp. Phytocrene densipunctata Stull, Moore et Manchester Phytocrene microcarpa Scott & Barghoorn Phytocrene punctilinearis Collinson, Manchester et Wilde 	EndocarpIndependence Hill (SN)EndocarpS, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpSt, Pankraz, BavariaEndocarpGrès de BelleuEndocarpVancouver islandEndocarpYakutat Frm.EndocarpSheppey, London ClayEndocarpMibanks II Clay Pits (TE)EndocarpAndrovette Pit, Kreischerville (NY)EndocarpMessel biota	Germany Canada USA USA USA USA USA Germany France Canada Alaska United Kingdom USA USA USA	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9 47.8 37.8 66 \approx 47	Eocene Eocene Eocene Paleocene Oligocene Eocene Eocene Eocene Eocene Eocene Cretaceous	? Lutetian Late Cuisian Sup. ? Middle-Late Ypresian Bartonian Early-Late Lutetian	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Stull et al., 2016AcceptedStull et al., 2011RejectedScott & Barghoorn, 1957AcceptedCollinson, 2012
 Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäutl) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. Phytocrene elytraeformis (Holl.) Wolfe Perforatocarpum echinatum (Chandler) Stull, Adams, Manchester et Collinson Phytocrene densipunctata Stull, Moore et Manchester Phytocrene microcarpa Scott & Barghoorn Phytocrene sp. Pyrenacantha austroamericana Stull 	EndocarpIndependence Hill (SN)EndocarpS, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpJohn Day Formation (Or)EndocarpS, Pankraz, BavariaEndocarpGrès de BelleuEndocarpVancouver islandEndocarpYakuta Frm.EndocarpSheppey, London ClayEndocarpMibanks II Clay Pits (TE)EndocarpAndrovette Pit, Kreischerville (NY)EndocarpGuajira peninsulaEndocarpBelen flora	Germany Canada USA USA USA USA USA Germany France Canada Alaska United Kingdom USA USA Gremany Colombia Peru	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5	EoceneEoceneEocenePaleoceneOligoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneEoceneOligoceneOligocene	? Lutetian Late Cuisian Sup. ? Middle-Late Ypresian Bartonian Early-Late Lutetian Thanetian Late early	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Stull et al., 2016AcceptedScott & Barghoorn, 1957AcceptedStull et al., 2012AcceptedStull et al., 2012
 Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäut)) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. Palaeophytocrene elytraeformis (Holl.) Wolfe Perforatocarpum echinatum (Chandler) Stull, Adams, Manchester Phytocrene microcarpa Scott & Barghoorn Phytocrene punctilinearis Collinson, Manchestet et Wilde Cf. Phytocrene sp. Pyrenacantha austroamericana Stull Pyrenacantha occidentalis Manchester Pyrenacantha sp. 1 	EndocarpIndependence Hill (SN)EndocarpS, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpJohn Day Formation (Or)EndocarpJohn Day Formation (Or)EndocarpGrès de BelleuEndocarpYancouver islandEndocarpYakutat Frm.EndocarpSheppey, London ClayEndocarpWilbanks II Clay Pits (TE)EndocarpMacovette Pit, Kreischerville (NY)EndocarpGuajira peninsulaEndocarpBelen floraEndocarpClarno Nut Beds (OR)EndocarpKessel biotaEndocarpKessel biotaEndocarpKess	USAGermanyCanadaUSAUSAUSAUSAGermanyFranceCanadaAlaskaUISAUSAGermanyColombiaPeruUSAGermany	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47	Eocene Eocene Eocene Paleocene Oligocene Eocene Eocene Eocene Eocene Eocene Eocene Eocene Eocene Eocene	? Lutetian Late Cuisian Sup. ? Middle-Late Ypresian Bartonian Early-Late Lutetian Thanetian Late early Lutetian	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedStull et al., 2016AcceptedStull et al., 2011RejectedScott & Barghoorn, 1957AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedManchester, 1994AcceptedCollinson et al., 2012
 Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäut) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. Phytocrene densipunctata Stull, Moore et Manchester Phytocrene punctilinearis Collinson, Manchestet et Wilde Cj. Phytocrene sp. Pyrenacantha austroamericana Stull Pyrenacantha occidentalis Manchester Pyrenacantha sp. 1 Pyrenacantha sp. 2 Sphaeriodes ventricosa (Bowerbank) Reid and Chandler Sitzocarya communis Reid and Chandler 	EndocarpIndependence Hill (SN)EndocarpSt, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpSt, Pankraz, BavariaEndocarpGrès de BelleuEndocarpYakuta Frm.EndocarpSheppey, London ClayEndocarpMibanks II Clay Pits (TE)EndocarpMessel biotaEndocarpGuajira peninsulaEndocarpBelen floraEndocarpClarno Nut Beds (OR)EndocarpMessel biotaEndocarpMessel biotaEndocarpSheppey, London ClayEndocarpSheppey Condon Clay	Germany Ganada USA USA USA USA USA USA Germany France Canada Alaska United Kingdom USA USA Gremany Colombia Peru USA Germany Colombia	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 ≈ 47 ≈ 47 ≈ 47 ≈ 47 ≈ 7 ≈ 7	Eocene Eocene Eocene Paleocene Oligocene Eocene Eocene Eocene Eocene Eocene Oligocene Eocene Eocene Eocene Eocene	? Lutetian Late Cuisian Sup. ? Middle-Late Ypresian Bartonian Early-Late Lutetian Lutetian Lutetian Lutetian Lutetian Lutetian	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedStull et al., 2016AcceptedStull et al., 2011RejectedScott & Barghoorn, 1957AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedManchester, 1994AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1932
 Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäutl) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. Phytocrene densipunctata Stull, Moore et Manchester Phytocrene sp. Pyrenacantha austroamericana Stull Pyrenacantha occidentalis Manchester Pyrenacantha sp. 1 Pyrenacantha sp. 2 Sphaeriodes ventricosa (Bowerbank) Reid and Chandler Stizocarya oviformis Reid and Chandler Stizocarya oviformis Reid and Chandler Icacinanthium tainiaphorum Del Rio & De Franceschi 	EndocarpIndependence Hill (SN)EndocarpS, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpAmont (ND)EndocarpJohn Day Formation (Or)EndocarpS, Pankraz, BavariaEndocarpGrès de BelleuEndocarpVancouver islandEndocarpYakuta Frm.EndocarpSheppey, London ClayEndocarpMissel biotaEndocarpGuajira peninsulaEndocarpGarno Nut Beds (OR)EndocarpSesel biotaEndocarpSheppey, London ClayEndocarpMissel biotaEndocarpMissel biotaEndocarpSheppey InsulaEndocarpSheppey InsulaEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpSh	USAGermanyCanadaUSAUSAUSAUSAGermanyFranceCanadaAlaskaUNIED KingdomUSAUSAGermanyColombiaPeruUSAGermanyColombiaPeruUSAUSAUSAUSAUSAUSAUSAUSAUSAUSAFranceUSAFrance	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 8 47.8	EoceneEoceneEoceneEoceneOligoceneEocene	?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly-LateLutetianLutetianLutetianLutetianLutetianYpresianSate earlyLutetianLutetianLutetianYpresian	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Stull et al., 2016AcceptedStull et al., 2011RejectedScott & Barghoorn, 1957AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedManchester, 1994AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler,
Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäut) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene pseudopersica Scott emend Manchester Palaeophytocrene sp. Pytocrene microcarpa Scott & Barghoorn Phytocrene sp. Pyrenacantha austroamericana Stull Pyrenacantha sp. 1 Pyrenacantha sp. 2 Sphaeriodes ventricosa (Bowerbank) Reid and Chandler Stizocarya coiformis Reid and Chandler Goweria alas	EndocarpIndependence Hill (SN)EndocarpS. Pankraz, BavariaEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittias County (WA)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpS. Pankraz, BavariaEndocarpGreis de BelleuEndocarpYakuta Frm.EndocarpShopey, London ClayEndocarpMissel biotaEndocarpGuajira peninsulaEndocarpGuajira peninsulaEndocarpShepey, London ClayEndocarpShesel biotaEndocarpMissel biotaEndocarpShepey, London ClayEndocarpShepey, London ClayEndocar	USAGermanyCanadaUSAUSAUSAUSAGermanyFranceCanadaAlaskaUNited KingdomUSAUSAGermanyIUSAGarmanyUSAUSAUnited KingdomUSAGermanyUSAIUSAGremanyIUSAColombiaPeruUSAGermanyIUSAJapan	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 ≈ 47 47.8 47.8 3.9 ≈ 47 ≈ 47 47.8 47.8 3.9 ≈ 47 ≈ 47 47.8 47.	EoceneEoceneEoceneEoceneOligoceneEocene	?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly -LateLutetianThanetianLutetianLutetianYpresianJpresianSatonianSatonianLutetianLutetianSasal sparnatianMiddle-LateMiddle	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedKull et al., 2008AcceptedStull et al., 2016AcceptedStull et al., 2011RejectedScott & Barghoorn, 1957AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousWolfe, 1977
Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäut) Palaeophytocrene speudopersica Scott emend Manchester Palaeophytocrene sp. Phytocrene densipunctata Stull, Moore et Manchester Phytocrene punctilinearis Collinson, Manchestet et Wilde Cf. Phytocrene sp. Pyrenacantha austroamericana Stull Pyrenacantha sp. 1 Pyrenacantha sp. 2 Splaeriodes ventricosa (Bowerbank) Reid and Chandler Stizocarya oviformis Reid and Chandler Stizocarya oviformis Reid and Chandler	EndocarpIndependence Hill (SN)EndocarpS, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpAmont (ND)EndocarpJohn Day Formation (Or)EndocarpGrès de BelleuEndocarpGrès de BelleuEndocarpGrès de JelleuEndocarpStattat Frm.EndocarpSheppey, London ClayEndocarpGuajira peninsulaEndocarpGuajira peninsulaEndocarpGlaron Nut Beds (OR)EndocarpGlaron Nut Beds (OR)EndocarpGlaron Nut Beds (OR)EndocarpGuajira peninsulaEndocarpGlaron Nut Beds (OR)EndocarpSheppey, London ClayEndocarpSheppey, London Clay <td>USAGermanyCanadaUSAUSAUSAUSAGermanyFranceCanadaAlaskaUNIAUSAGermanyUSAUSAGremanyUSAGremanyUSAUSAUSAJapanJapanUSAJapanUSAJapanJapanJAlaskaJapanJAlaskaJapanJAlaskaJapan</td> <td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 47.8 33.9 ≈ 46 49.5 33.9</td> <td>EoceneEoceneEoceneEoceneOligoceneEocene</td> <td>?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly-LateLutetianItate earlyLutetianYpresianYpresianSasal sparnatianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-Late</td> <td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedKouffe, 1977; Hollick, 1936; Heer, 1870AcceptedStull et al., 2016AcceptedStull et al., 2011RejectedScott & Barghoorn, 1957AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousTanai, 1990DubiousAllen et al., 2015DubiousWolfe, 1977</td>	USAGermanyCanadaUSAUSAUSAUSAGermanyFranceCanadaAlaskaUNIAUSAGermanyUSAUSAGremanyUSAGremanyUSAUSAUSAJapanJapanUSAJapanUSAJapanJapanJAlaskaJapanJAlaskaJapanJAlaskaJapan	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 47.8 33.9 ≈ 46 49.5 33.9	EoceneEoceneEoceneEoceneOligoceneEocene	?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly-LateLutetianItate earlyLutetianYpresianYpresianSasal sparnatianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-Late	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedKouffe, 1977; Hollick, 1936; Heer, 1870AcceptedStull et al., 2016AcceptedStull et al., 2011RejectedScott & Barghoorn, 1957AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousTanai, 1990DubiousAllen et al., 2015DubiousWolfe, 1977
Palaeophytocrene lacunosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäut) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene speudopersica Scott emend Manchester Palaeophytocrene sp. Phytocrene densipunctata Sull, Moore et Manchester Phytocrene microcarpa Scott & Barghoorn Phytocrene sp. Pyrenacantha austroamericana Stull Pyrenacantha sp. 1 Pyrenacantha sp. 2 Sphaeriodes ventricosa (Bowerbank) Reid and Chandler Stizocarya ovijormis Reid and Chandler Goweria alaskana Wolfe Goweria bibaiensis Tanai Goweria bibaiensis Tanai Goweria dilleri (Knowlon) Wolfe Goweria lineari	EndocarpNdependence Hill (SN)EndocarpSt, Pankraz, BavariaEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittias County (WA)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpGrès de BelleuEndocarpGrès de BelleuEndocarpVancouver islandEndocarpSheppey, London ClayEndocarpNibanks II Clay Pits (TE)EndocarpNibanks II Clay Pits (TE)EndocarpScale biotaEndocarpGuajira peninsulaEndocarpGuajira peninsulaEndocarpSheppey, London ClayEndocarpSheppey, London Clay <trr>Endocarp</trr>	CSAGermanyCanadaUSAUSAUSAUSAGarmanyFranceCanadaAlaskaUSAUSAGermanyUSAUSAUSAQanadaQanadaInited KingdomQarmanyUSAGermanyUSAGermanyQarmanyQarmanyQarmanyUSAQarmanyQarmanyQarmanyUSAQarmany <td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 47.8 47.8 47.8 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8</td> <td>EoceneEoceneEoceneColeoceneOligoceneEocene<!--</td--><td>?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly -LateLutetianThanetianLutetianYpresianYpresianSasal sparnatianMiddle-LateMiddle-LateMiddleSasal sparnatianMiddle-LateMiddleSasal sparnatianMiddleSasal sparnatianMiddleSasal sparnatianSasal sparnatian<</td><td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedKodfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Stull et al., 2016AcceptedStull et al., 2016AcceptedStull et al., 2011AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousVolfe, 1977DubiousAllen et al., 2015DubiousWolfe, 1968; Wolfe 1977DubiousWolfe, 1968RejectedStuller et al., 2015</td></td>	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 47.8 47.8 47.8 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8	EoceneEoceneEoceneColeoceneOligoceneEocene </td <td>?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly -LateLutetianThanetianLutetianYpresianYpresianSasal sparnatianMiddle-LateMiddle-LateMiddleSasal sparnatianMiddle-LateMiddleSasal sparnatianMiddleSasal sparnatianMiddleSasal sparnatianSasal sparnatian<</td> <td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedKodfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Stull et al., 2016AcceptedStull et al., 2016AcceptedStull et al., 2011AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousVolfe, 1977DubiousAllen et al., 2015DubiousWolfe, 1968; Wolfe 1977DubiousWolfe, 1968RejectedStuller et al., 2015</td>	?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly -LateLutetianThanetianLutetianYpresianYpresianSasal sparnatianMiddle-LateMiddle-LateMiddleSasal sparnatianMiddle-LateMiddleSasal sparnatianMiddleSasal sparnatianMiddleSasal sparnatianSasal sparnatian<	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedKodfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Stull et al., 2016AcceptedStull et al., 2016AcceptedStull et al., 2011AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousVolfe, 1977DubiousAllen et al., 2015DubiousWolfe, 1968; Wolfe 1977DubiousWolfe, 1968RejectedStuller et al., 2015
 Palaeophytocrene lacunosum (MacGinitic) Wolfe Palaeophytocrene kressenbergensis (Schafhäut) Palaeophytocrene manchesteri Rankin, Stockey et Beard Palaeophytocrene sp. Palaeophytocrene vancouverensis Rankin, Stockey et Beard Paleophytocrene elytraeformis (Holl.) Wolfe Perforatocarpum echinatum (Chandler) Stull, Adams, Manchester et Collinson Phytocrene densipunctata Stull, Moore et Manchester Phytocrene sp. Pyrenacantha austroamericana Stull Pyrenacantha sp. 1 Pyrenacantha sp. 1 Pyrenacantha sp. 2 Sphaeriodes ventricosa (Bowerbank) Reid and Chandler Stizocarya oviformis Reid and Chandler Stizocarya oviformis Reid and Chandler Goweria alaskana Wolfe Goweria bilaeinsis Tanai Goweria dilleri (Knowhon) Wolfe Goweria dilleri (Knowhon) Wolfe Huziokaeae eoutilus (Endo) Tanai Huziokaeae eoutilus (Endo) Tanai Huziokaeae eoutilus (Endo) Tanai 	EndocarpIndependence Hill (SN)EndocarpS, Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpS, Pankraz, BavariaEndocarpGracouver islandEndocarpVancouver islandEndocarpYakutat Frm.EndocarpVancouver islandEndocarpValtatas I'L Clay Pits (TE)EndocarpMibanks II Clay Pits (TE)EndocarpMessel biotaEndocarpGalira peninsulaEndocarpBelen floraEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpMessel biotaEndocarpSheppey, London ClayEndocarpSheppey, London Clay	OSA Germany Canada USA Germany France Canada VIA VISA USA USA USA OBERNANC VIA VISA OCIOMBIA Peru VIA	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 4	EoceneEoceneEoceneOligoceneCoceneEocene	?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly -LateLutetianThanetianLutetianYpresianJpresianSasal sparnatianMiddle-LateMiddle-LateMiddle-LateMiddleMiddleMiddleSasal sparnatianMiddleMiddl	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedPolfe, 1977; Hollick, 1936; Heer, 1870AcceptedChandler, 1961b; Cleal et al., 2001; Sull et al., 2016AcceptedStull et al., 2012AcceptedStull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousWolfe, 1977DubiousMoler, 1968; Wolfe 1977DubiousWolfe, 1968; Wolfe 1977DubiousWolfe, 1968; Wolfe 1977DubiousKolfe, 1968; Wolfe 1977DubiousKolfe, 1968NubiousKolfe, 1976NubiousKolfe, 1976NubiousKolfe, 1976NubiousKolfe, 1968; Wolfe 1977NubiousKolfe, 1976NubiousKolfe, 1976NubiousKolfe, 1977NubiousKolfe, 1976NubiousKolfe, 1977NubiousKolf
Palaeophytocrene kacanosum (MacGinitie) Wolfe Palaeophytocrene kressenbergensis (Schafhäut)) Palaeophytocrene pseudopersica Scott ement Manchester Palaeophytocrene sp. Palaeophytocrene sp. Phytocrene microcarpa Scott & Barghoorn Phytocrene microcarpa Scott & Barghoorn Phytocrene punctilinearis Collinson, Manchestert et Wilde Cf. Phytocrene sp. Pyrenacantha austroamericana Stull Pyrenacantha austroamericana Stull Pyrenacantha sp. 1 Pyrenacantha sp. 1 Staocarya communis Reid and Chandler Staocarya oviformis Reid and Chandler Staocarya oviformis Reid and Chandler Goweria diaskana Wolfe Goweria diaskana Wolfe Inziokaeae eoutikus (Endo) Tanai Merrilliodendron ezoanum Tanai Phytocrene acutissima Wolfe Phytocrene acutissima Wolfe	EndocarpIndependence Hill (SN)EndocarpS. Pankraz, BavariaEndocarpVancouver islandEndocarpClarno Nut Beds (OR)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpS. Pankraz, BavariaEndocarpS. Pankraz, BavariaEndocarpS. Pankraz, BavariaEndocarpS. Pankraz, BavariaEndocarpVancouver islandEndocarpYakuta Frm.EndocarpVancouver islandEndocarpNeppey, London ClayEndocarpGuajira peninsulaEndocarpBelen floraEndocarpBelen floraEndocarpSheppey, London ClayEndocarpSheppey, London Clay	OSA Germany USA Oarnany ISA USA USA USA Oarnany USA Oarnany Oarnany <td< td=""><td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 33.9 ≈ 40 33.9 ≈ 40 33.9 ≈ 40 33.9 47.8 40 33.9 40 4</td><td>EoceneEoceneEoceneOligoceneCoceneEoceneEoceneEoceneEoceneEoceneCretaceousFoceneEocene<trr>Eocene<trr>Eocen</trr></trr></td><td>?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly -LateLutetianThanetianLate earlyLutetianYpresianYpresianYpresianMiddle-Late<t< td=""><td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedScott & Barghoorn, 1957AcceptedScott & Barghoorn, 1957AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousWolfe, 1968; Wolfe 1977DubiousWolfe, 1968; Wolfe 1977DubiousKolfe, 1968RejectedTanai, 1990DubiousWolfe, 1977DubiousWolfe, 1977<t< td=""></t<></td></t<></td></td<>	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 33.9 ≈ 40 33.9 ≈ 40 33.9 ≈ 40 33.9 47.8 40 33.9 40 4	EoceneEoceneEoceneOligoceneCoceneEoceneEoceneEoceneEoceneEoceneCretaceousFoceneEocene <trr>Eocene<trr>Eocen</trr></trr>	?LutetianLateCuisian Sup.?Middle-LateYpresianBartonianEarly -LateLutetianThanetianLate earlyLutetianYpresianYpresianYpresianMiddle-Late <t< td=""><td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedScott & Barghoorn, 1957AcceptedScott & Barghoorn, 1957AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousWolfe, 1968; Wolfe 1977DubiousWolfe, 1968; Wolfe 1977DubiousKolfe, 1968RejectedTanai, 1990DubiousWolfe, 1977DubiousWolfe, 1977<t< td=""></t<></td></t<>	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedScott & Barghoorn, 1957AcceptedScott & Barghoorn, 1957AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousWolfe, 1968; Wolfe 1977DubiousWolfe, 1968; Wolfe 1977DubiousKolfe, 1968RejectedTanai, 1990DubiousWolfe, 1977DubiousWolfe, 1977 <t< td=""></t<>
Palacophytocrene lacanosum (MacGinitie) Wolfe Palacophytocrene kressenbergensis (Schafhäut) Palacophytocrene manchesteri Rankin, Stockey et Beard Palacophytocrene sp. Palacophytocrene sp. Phytocrene densipuerata Stull, Moore et Manchester Phytocrene punctilinearis Collinson, Manchestet et Wilde G. Phytocrene sp. Pyrenacantha occidentalis Manchester Pyrenacantha sp. 1 Pyrenacantha sp. 1 Stocarya communis Reid and Chandler Stocarya communis Reid and Chandler Stocarya ovformis Reid and Chandler Goweria bibaiensis Tanai Goweria bibaiensis Tanai Goweria bibaiensis Tanai Goweria linearis Wolfe Huziokaeae eoutilus (Endo) Tanai Phytocrene acutissima Wolfe Phytocrene acutistima Wolfe Phytocrene	EndocarpIndependence Hill (SN)EndocarpSt, Pankraz, BavariaEndocarpClarno Nut Beds (OR)EndocarpRonald, Kittitas County (WA)EndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpSt, Pankraz, BavariaEndocarpGrès de BelleuEndocarpVancouver islandEndocarpVancouver islandEndocarpValatat Frm.EndocarpSheppey, London ClayEndocarpMessel biotaEndocarpGuajira peninsulaEndocarpGlarno Nut Beds (OR)EndocarpBelen floraEndocarpMessel biotaEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpMessel biotaEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarpBlue Rim (WY)LeafQiauno-sawa, Bibai, HokkaidoLeafGreen river (WY)LeafKushtaka Frm.LeafKushtaka Frm.LeafKushtaka Frm.LeafKushtaka Frm.LeafKushtaka Frm.LeafKushtaka Frm.LeafKushtaka Frm.LeafKushtaka Frm.LeafKushtaka Frm.LeafKushtaka Frm.Leaf<	Germany Ganada USA Qarmany France Canada Alaska United Kingdom Qermany ISA QisA QisAsa QisAsa <td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 33.9 ≈ 40 33.9 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.9 37.8 37.8 37.8 37.9 37.8 37.8</td> <td>EoceneEoceneEoceneOligoceneEocene</td> <td>?LutetianLateCuisian Sup.?Middle-LateYpresianEarly-LateIutetianLutetianLutetianYpresianYpresianYpresianYpresianYpresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleYpresianMiddle-Late<trtr>Middle-LateMiddl</trtr></td> <td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRunkin et al., 2008AcceptedSuill et al., 2016AcceptedSuill et al., 2011AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousAllen et al., 2015DubiousVolfe, 1977DubiousVolfe, 1968Reij et al.Suill et al., 2015DubiousYolfe, 1968Reij et al.Suill et al., 2015DubiousTanai, 1990DubiousYolfe, 1977DubiousVolfe, 1977DubiousYolfe, 1977DubiousYolfe, 1977DubiousYolfe, 1977DubiousYolfe, 1977DubiousYolfe, 1977DubiousYol</td>	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 33.9 ≈ 40 33.9 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.8 37.9 37.8 37.8 37.8 37.9 37.8 37.8	EoceneEoceneEoceneOligoceneEocene	?LutetianLateCuisian Sup.?Middle-LateYpresianEarly-LateIutetianLutetianLutetianYpresianYpresianYpresianYpresianYpresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleYpresianMiddle-Late <trtr>Middle-LateMiddl</trtr>	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRunkin et al., 2008AcceptedSuill et al., 2016AcceptedSuill et al., 2011AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousAllen et al., 2015DubiousVolfe, 1977DubiousVolfe, 1968Reij et al.Suill et al., 2015DubiousYolfe, 1968Reij et al.Suill et al., 2015DubiousTanai, 1990DubiousYolfe, 1977DubiousVolfe, 1977DubiousYolfe, 1977DubiousYolfe, 1977DubiousYolfe, 1977DubiousYolfe, 1977DubiousYolfe, 1977DubiousYol
Palacophytocrene lacunosum (MacGinitic) Wolfe Palacophytocrene lacunosum (MacGinitic) Wolfe Palacophytocrene wanchesteri Rankin, Stockey et Beard Palacophytocrene sp. Palacophytocrene sp. Phytocrene sp. Phytocr	EndocarpK.Pankraz, BavariaEndocarpK.Pankraz, BavariaEndocarpClarno Nut Beds (OR)EndocarpK.Pankraz, BavariaEndocarpAlmont (ND)EndocarpJohn Day Formation (Or)EndocarpK.Pankraz, BavariaEndocarpK.Pankraz, BavariaEndocarpK.Pankraz, BavariaEndocarpK.Pankraz, BavariaEndocarpK.Pankraz, BavariaEndocarpVancouver islandEndocarpYalutat Frm.EndocarpNeppey, London ClayEndocarpMibanks II Clay Pits (TE)EndocarpMarovette Pit, Kreischerville (NY)EndocarpGuajira peninsulaEndocarpGuajira peninsulaEndocarpKessel biotaEndocarpMessel biotaEndocarpSheppey, London ClayEndocarpSheppey, London ClayEndoKushtaka Frm. </td <td>OSA Germany Canada USA USA USA USA USA USA USA USA USA Germany Fance Canada Alaska UNited Kingdom Qermany USA Germany QiSA QiSA QiSA QiSA Qisamany Qisamany Qisamany Qisamany Qisamany Qisamany Qisaka Qisapan<td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 3.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 5.5-56 33.9 ≈ 40 33.9 ≈ 40 5.6</td><td>EoceneEoceneEoceneOligoceneCoceneEocene</td><td>?LutetianLateCuisian Sup.?Middle-LateQuresianBartonianEarly-LateLutetianLutetianLutetianQuresianQuresianYpresianMiddle-LateMiddle.Late<tr< td=""><td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedChandler, 1961b; Cleal et al., 2001; Suff et al., 2012AcceptedStull et al., 2016AcceptedStull et al., 2012AcceptedSuff et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousJen et al., 2015DubiousJen et al., 2015DubiousMolfe, 1968; Wolfe 1977DubiousJen et al., 2015DubiousTanai, 1990DubiousTanai, 1990DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977Dubious<t< td=""></t<></td></tr<></td></td>	OSA Germany Canada USA USA USA USA USA USA USA USA USA Germany Fance Canada Alaska UNited Kingdom Qermany USA Germany QiSA QiSA QiSA QiSA Qisamany Qisamany Qisamany Qisamany Qisamany Qisamany Qisaka Qisapan <td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 3.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 5.5-56 33.9 ≈ 40 33.9 ≈ 40 5.6</td> <td>EoceneEoceneEoceneOligoceneCoceneEocene</td> <td>?LutetianLateCuisian Sup.?Middle-LateQuresianBartonianEarly-LateLutetianLutetianLutetianQuresianQuresianYpresianMiddle-LateMiddle.Late<tr< td=""><td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedChandler, 1961b; Cleal et al., 2001; Suff et al., 2012AcceptedStull et al., 2016AcceptedStull et al., 2012AcceptedSuff et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousJen et al., 2015DubiousJen et al., 2015DubiousMolfe, 1968; Wolfe 1977DubiousJen et al., 2015DubiousTanai, 1990DubiousTanai, 1990DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977Dubious<t< td=""></t<></td></tr<></td>	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 3.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 5.5-56 33.9 ≈ 40 33.9 ≈ 40 5.6	EoceneEoceneEoceneOligoceneCoceneEocene	?LutetianLateCuisian Sup.?Middle-LateQuresianBartonianEarly-LateLutetianLutetianLutetianQuresianQuresianYpresianMiddle-LateMiddle.Late <tr< td=""><td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedChandler, 1961b; Cleal et al., 2001; Suff et al., 2012AcceptedStull et al., 2016AcceptedStull et al., 2012AcceptedSuff et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousJen et al., 2015DubiousJen et al., 2015DubiousMolfe, 1968; Wolfe 1977DubiousJen et al., 2015DubiousTanai, 1990DubiousTanai, 1990DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977Dubious<t< td=""></t<></td></tr<>	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedChandler, 1961b; Cleal et al., 2001; Suff et al., 2012AcceptedStull et al., 2016AcceptedStull et al., 2012AcceptedSuff et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDel Rio et al., 2017DubiousJen et al., 2015DubiousJen et al., 2015DubiousMolfe, 1968; Wolfe 1977DubiousJen et al., 2015DubiousTanai, 1990DubiousTanai, 1990DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977Dubious <t< td=""></t<>
Palaeophytocrene lacunosam (MacGinitie) Wolfe Palaeophytocrene lacunosam (MacGinitie) Wolfe Palaeophytocrene iressenbergensis (Schafhäut) Palaeophytocrene sp. Palaeophytocrene sp. Phytocrene punctilmeatra Sull, Moore et Manchester Phytocrene punctilmeatra Sull, Moore et Manchester Phytocrene sp. Pyrenacantha astroamericana Sull Pyrenacantha occidentalis Manchester Pyrenacantha sp. 1 Pyrenacantha sp. 1 Pyrenacantha sp. 2 Sphaeriodes ventricosa (Bowethanh) Reid and Chandler Sitzocarya ovijornis Reid and Chandler Goweria alaskana Wolfe Goweria dibaiensis Tanai Goweria dibaiensis Tanai Phytocrene acutistima Wolfe Huziokaeae eoutilus (Endo) Tanai Hervillodendron ezoanum Tanai Phytocrene acutistima Wolfe Phytocrene sordida (Lesuereux.) MacGinitie Phytocrene sordida (Lesuereux.) MacGinitie Phytocrene acutistima cellas Krutzch & Vanhoorne 1977 Compositoipolenites medius Krutzch & Vanhoorne 1977	EndocarpK.Pankraz, BavariaEndocarpK.Pankraz, BavariaEndocarpClarno Nut Beds (OR)EndocarpK.Pankraz, BavariaEndocarpJohn Day Formation (Or)EndocarpJohn Day Formation (Or)EndocarpS.Pankraz, BavariaEndocarpGras de BelleuEndocarpYalutat Frm.EndocarpSheppey, London ClayEndocarpMorovette Pit, Kreischerville (NY)EndocarpMoseel biotaEndocarpGuajira peninsulaEndocarpGlarno Nut Beds (OR)EndocarpGlarno Nut Beds (OR)EndocarpGlarno Nut Beds (OR)EndocarpGlarno Nut Beds (OR)EndocarpGlarno Nut Beds (OR)EndocarpSheppey, London ClayEndocarpSheppey, London ClayEndocarp	Germany Ganada USA Germany France Canada Alaska USA USA USA Germany USA Gremany USA Germany USA Germany USA Germany USA Germany USA Germany USA Germany USA Japan USA Japan Alaska USA Japan Japan Japan Japan Japan Japan Japan Japan Japan <td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9 47.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 ≈ 47 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 33.9 ≈ 40 33.9 ≈ 40 56</td> <td>EoceneEoceneEoceneOligoceneEocene</td> <td>?LutetianLateCuisian Sup.?Middle-LateYpresianEarly-LateLutetianLutetianLutetianYpresianSasal sparnatianMiddle-LateMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidel<td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedSuill et al., 2011AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousJene et al., 2015DubiousWolfe, 1968RejectedTanai, 1990DubiousWolfe, 1977DubiousWolfe, 1977DubiousWolfe, 1977DubiousMacginitie, 1941DubiousAccepted, 1990DubiousCavagnetto, 2000DubiousKause, K Vanhoorne, 1977DubiousCavagnetto, 2000DubiousKrutzsch & Vanhoorne, 1977DubiousCavagnetto, 2000</td></td>	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 33.9 47.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 ≈ 47 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 33.9 ≈ 40 33.9 ≈ 40 56	EoceneEoceneEoceneOligoceneEocene	?LutetianLateCuisian Sup.?Middle-LateYpresianEarly-LateLutetianLutetianLutetianYpresianSasal sparnatianMiddle-LateMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidelMidel <td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedSuill et al., 2011AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousJene et al., 2015DubiousWolfe, 1968RejectedTanai, 1990DubiousWolfe, 1977DubiousWolfe, 1977DubiousWolfe, 1977DubiousMacginitie, 1941DubiousAccepted, 1990DubiousCavagnetto, 2000DubiousKause, K Vanhoorne, 1977DubiousCavagnetto, 2000DubiousKrutzsch & Vanhoorne, 1977DubiousCavagnetto, 2000</td>	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedSuill et al., 2011AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousJene et al., 2015DubiousWolfe, 1968RejectedTanai, 1990DubiousWolfe, 1977DubiousWolfe, 1977DubiousWolfe, 1977DubiousMacginitie, 1941DubiousAccepted, 1990DubiousCavagnetto, 2000DubiousKause, K Vanhoorne, 1977DubiousCavagnetto, 2000DubiousKrutzsch & Vanhoorne, 1977DubiousCavagnetto, 2000
Palacophytocrene lacunoum (MacGinitic) Wolfe Palacophytocrene lacunoum (MacGinitic) Wolfe Palacophytocrene pseulopersica Scott emend Manchester Palacophytocrene sp. Phytocrene panetilinearis Collinson, Manchester et Wilde Cy. Phytocrene sp. Pyrenacantha accidentalis Manchester Stizocarya communis Reid and Chandler Stizocarya ovijornis Reid and Chandler Gaveria diskana Wolfe Goweria bherimensis S.F. Allen, Sull, & Mancheberser Phytocrene acutistima Wolfe Phytocrene acutistima Wolfe <t< td=""><td>Endocary Rodener Hill (SN) Endocary Sepankraz, Bavaria Endocary Carno-Nut Beds (OR) Endocary Sobept - London Clay Endocary Carno-Nut Beds (OR) Endocary Sobept - London Clay Endocary Sobept - London Clay <</td><td>OSAGermanyCanadaUSAUSAUSAUSAGermanyFanceCanadaAlaskaUNICH KingdomUSAOremanyUSAGermanyUSAGermanyUSAGermanyUSAGermanyUSAGermanyUSAGermanyUSAJapanJapanJapanJapanAlaskaJapanAlaskaJapanAlaskaJapanFranceAlaskaJapanJapanFranceAlaskaJapanFranceAlaskaJapanFranceFranceBelgiumFranceBelgiumBelgiumFranceBelgiumFrance<</td><td>37.8 33.9-56 33.9 43 3.9 56 32,2-33,6 49 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 33.9 ≈ 40 33.9 ≈ 40 56 5</td><td>EoceneEoceneEoceneOligoceneEocene</td><td>?LutetianLateCuisian Sup.?Middle-LateQuresianBartonianEarly-LateLutetianThanetianLutetianQuresianQuresianYpresianYpresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddleYpresianMiddleYpresianYiresianYiresianMiddleYiresianYires</td><td>AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRandner, 1961b: Cleal et al., 2001AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedSuill et al., 2016AcceptedSuill et al., 2011AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2013AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousAlen et al., 2017DubiousAlen et al., 2015DubiousAlen et al., 2015DubiousAlen et al., 2015DubiousMolfe, 1968; Wolfe 1977DubiousMolfe, 1977<t< td=""></t<></td></t<>	Endocary Rodener Hill (SN) Endocary Sepankraz, Bavaria Endocary Carno-Nut Beds (OR) Endocary Sobept - London Clay Endocary Carno-Nut Beds (OR) Endocary Sobept - London Clay Endocary Sobept - London Clay <	OSAGermanyCanadaUSAUSAUSAUSAGermanyFanceCanadaAlaskaUNICH KingdomUSAOremanyUSAGermanyUSAGermanyUSAGermanyUSAGermanyUSAGermanyUSAGermanyUSAJapanJapanJapanJapanAlaskaJapanAlaskaJapanAlaskaJapanFranceAlaskaJapanJapanFranceAlaskaJapanFranceAlaskaJapanFranceFranceBelgiumFranceBelgiumBelgiumFranceBelgiumFrance<	37.8 33.9-56 33.9 43 3.9 56 32,2-33,6 49 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 33.9 ≈ 40 33.9 ≈ 40 56 5	EoceneEoceneEoceneOligoceneEocene	?LutetianLateCuisian Sup.?Middle-LateQuresianBartonianEarly-LateLutetianThanetianLutetianQuresianQuresianYpresianYpresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddleYpresianMiddleYpresianYiresianYiresianMiddleYiresianYires	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedRandner, 1961b: Cleal et al., 2001AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedSuill et al., 2016AcceptedSuill et al., 2011AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedSuill et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2013AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousAlen et al., 2017DubiousAlen et al., 2015DubiousAlen et al., 2015DubiousAlen et al., 2015DubiousMolfe, 1968; Wolfe 1977DubiousMolfe, 1977 <t< td=""></t<>
Palacophytocrene lacunosum (MacCimitic) Wulfe Palacophytocrene lacunosum (MacCimitic) Wulfe Palacophytocrene pseudopersica Scottemend Manchester Palacophytocrene sp. Palacophytocrene sp. Palacorafia austroamericana Sull, Moare et Manchester Phytocrene sp. Pyrenacamba sp. 1 Pyrenacamba Sp. 2 Sitocarya outformis Reid and Chandler Goweria bibatensis Tamai Goweria bibatensis Tamai Phytocrene soutific (Eado) Tamai Accimathiama iniaphorum Del Rio & De Franceschi Goweria bibatensis SE. Allen, Sull, & Man- Chester Goweria dilleri (Knowlon) Walfe Phytocrene soutific (Lesuercux,) MacGimitic Phytocrene soutific (Lesuercux,) MacGimitic	EndocaryNaceaurent Hill (SN)EndocarySa Pankraz BavariaEndocaryCancouver islandEndocaryRonald, Kittilas County (WA)EndocaryAlmont (ND)EndocaryOlm Day Formation (Or)EndocarySa Pankraz BavariaEndocarySa Pankraz BavariaEndocaryOlanouver islandEndocaryVancouver islandEndocarySabepey London ClayEndocaryNator (Day Formation (NT))EndocaryNacouver Bala (Sate)EndocaryNacouver Bala (Sate)EndocaryOlaria penissulaEndocarySabepey London ClayEndocaryOlaria penissulaEndocarySate) ForaEndocarySabepey London ClayEndocarySabepey London ClayLagíCanLagíSabe franLagíSabe franLagíSabe franLagíSabe franLagí	OSAGermanyCanadaUSAUSAUSAUSACananyFranceCanadaCanadaOaraadaUNAUNAUSAOaraanaOaraanaUSAOaraanay<	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 ≈ 47 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 47.8 33.9 ≈ 40 33.9 ≈ 40 56	EoceneEoceneEoceneOligoceneEoceneForeneEoceneEoceneForeneEoceneForeneEocene	?LutetianLateCuisian Sup.?Middle-LateQuresianEarly - LateLutetianLutetianLutetianQuresianQuresianYpresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleYpresianYpresianMiddle-LateMiddleMiddle-LateMiddle-LateMiddleYpresianYpresianYpresianMiddleMiddleMiddleMiddleDanianYpresianQuresianDanianNiddleMiddle	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedMey er & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedSull et al., 2016AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCollinson, 2012AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2013AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNolfe, 1977DubiousAllen et al., 2015DubiousAllen et al., 2015DubiousTanai, 1990DubiousTanai, 1990DubiousTanai, 1990DubiousColler, 1977DubiousCavagnetto, 2000DubiousTanai, 1990DubiousTanai, 1990DubiousTanai, 1990DubiousCavagnetto, 2000DubiousTanai, 1990DubiousTanai, 1990DubiousTanai, 1990DubiousTanai, 1990DubiousTanai, 1990D
Palacophytocrene lacunosum (MacCimito) Walfe Palacophytocrene lacunosum (MacCimito) Walfe Palacophytocrene manchester Rankin, Stuckey et Beard Palacophytocrene sp. Phytocrene ankerocarpa Solt & Barghorn Phytocrene sp. Phytocrene sp. Phytocrene sp. Pyrenacontha acstroumericana Sull Pyrenacontha acstroumericana Sull Pyrenacontha acstroumericana Sull Goweria akakara Wolfe Goweria bihatentis Tami Goweria bihatentis Tami Goweria bihatentis Tami Phytocrene coratitis (Knowhon) Wolfe <	EndocaryIndependence Hill (SN)EndocarySA Pankaz, BavariaEndocaryCamo Nut Beds (OR)EndocaryAmout (NU)EndocaryAmout (NU)EndocaryAmout (NU)EndocarySA Pankaz, BavariaEndocaryCarso de BelleuEndocaryCarso de BelleuEndocarySoneouver islandEndocarySoneouver islandEndocarySoneouver islandEndocaryAndrouteu Prix (Norteu Prix (No	OSAGermanyCanadaUSAUSAUSAUSAGermanyFanceCanadaAlaskaUSAUSAOremanyISAOremanyISAOremanyISAOremanyISAOremanyISAOremanyISAOremanyISAOremanyISAOremanyISAI	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 49 33.9 47.8 37.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 57-56 33.9 ≈ 40 33.9 $\approx 55-56$ 56	EoceneEoceneEoceneOligoceneEocene	?LutetianLateCuisian Sup.?Middle-Late?Middle-LateCutetianLutetianLutetianLutetianLutetianYpresianYpresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddle-LateMiddle-	AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedCran et al., 1990AcceptedTichy, 1979AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedWolfe, 1977; Hollick, 1936; Heer, 1870AcceptedSull et al., 2008AcceptedSull et al., 2011AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedSull et al., 2017DubiousTanai, 1990DubiousAllen et al., 2015DubiousYolfe, 1968; Wolfe 1977DubiousYolfe, 1977
 Palacophytocrene lacunosum (MacCimits) Wolfe Palacophytocrene lacunosum (MacCimits) Wolfe Palacophytocrene preudapersica Scott emend Manchester Palacophytocrene sp. Phytocrene densipunctam Stall, Moore et Manchester Phytocrene sp. Phytocrene sp. Phytocrene sp. Phytocrene sp. Pyrenacantha austroanericana Stall Pyrenacantha sp. 1 Pyrenacantha sp. 2 Sphaeriades ventricoa (Bowerbank) Reid and Chandler Sizocarya commanis Reid and Chandler Goweria alackana Wolfe Goweria alackana Wolfe Goweria alackana Wolfe Phytocrene acutistina Tanai Goweria alackana Wolfe Phytocrene acutistina Wolf	Radocary Ridependence Hill (SN) Radocary Ranara, Bavaria Radocary Carno Nut Beds (OR) Radocary Almont (ND) Radocary Almont (ND) Radocary Salancary (Nara, Bavaria) Radocary Carno Nut Beds (OR) Radocary </td <td>OSAGermanyCanadaUSAUSAUSAUSAGermanyFanceCanadaAlaskaUSAUSAOarnanyUSAOarnany</td> <td>37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 49 33.9 47.8 49 33.9 47.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 55-56 33.9 ≈ 40 33.9 ≈ 33.9 ≈ 40 33.9 ≈ 33.9 ≈ 33.9 $\propto 33.9$ ≈ 33.9 $\propto 33.9$ ≈ 33.9 $\propto 33.9$ $\propto 33.9$ $\propto 33.9$ $\propto 33.9$ $\propto 33.9$ ≈ 33.9 $\propto 33.9$ ≈ 33.9 $\propto 33.$</td> <td>EoceneEoceneEocenePaleoceneEocene</td> <td>? Luetian Lae Cuisian Sup. ? Middle-Late Paresian Luetian Tarly-Late Luetian Luetian Luetian Luetian Luetian Luetian Ypresian Kiddle-Late Widdle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle Middle Middle Middle Middle Middle Ypresian Ypresian</td> <td>AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedNotife, 1977; Hollick, 1936; Heer, 1870AcceptedSout & Barghoorn, 1957AcceptedSout & Barghoorn, 1957AcceptedSout & Barghoorn, 1957AcceptedSout & Barghoorn, 1957AcceptedSout et al., 2012AcceptedSout et al., 2012AcceptedSout et al., 2012AcceptedSout et al., 2012AcceptedSout et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDiel Ko et al., 2017AcceptedDiel Ko et al., 2017DubiousAllen et al., 2015DubiousAllen, 1963DubiousAllen, 1973DubiousNofe, 1977DubiousMolfe, 1977DubiousCanaj, 1990DubiousYolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousCanagentio, 2000DubiousCanagentio, 2000DubiousCanagentio, 2000DubiousCanagentio, 2000DubiousCanagentio, 2000<tr< td=""></tr<></td>	OSAGermanyCanadaUSAUSAUSAUSAGermanyFanceCanadaAlaskaUSAUSAOarnanyUSAOarnany	37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 49 33.9 47.8 49 33.9 47.8 66 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 58 28.5 43 ≈ 47 55-56 33.9 ≈ 40 33.9 ≈ 33.9 ≈ 40 33.9 ≈ 33.9 ≈ 33.9 $\propto 33.9$ ≈ 33.9 $\propto 33.9$ ≈ 33.9 $\propto 33.9$ $\propto 33.9$ $\propto 33.9$ $\propto 33.9$ $\propto 33.9$ ≈ 33.9 $\propto 33.9$ ≈ 33.9 $\propto 33.$	EoceneEoceneEocenePaleoceneEocene	? Luetian Lae Cuisian Sup. ? Middle-Late Paresian Luetian Tarly-Late Luetian Luetian Luetian Luetian Luetian Luetian Ypresian Kiddle-Late Widdle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle-Late Middle Middle Middle Middle Middle Middle Ypresian	AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedScott, 1954; Manchester, 1994AcceptedCrane et al., 1990AcceptedMeyer & Manchester, 1997AcceptedTichy, 1979AcceptedTichy, 1979AcceptedRankin et al., 2008AcceptedRankin et al., 2008AcceptedNotife, 1977; Hollick, 1936; Heer, 1870AcceptedSout & Barghoorn, 1957AcceptedSout & Barghoorn, 1957AcceptedSout & Barghoorn, 1957AcceptedSout & Barghoorn, 1957AcceptedSout et al., 2012AcceptedSout et al., 2012AcceptedSout et al., 2012AcceptedSout et al., 2012AcceptedSout et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedDiel Ko et al., 2017AcceptedDiel Ko et al., 2017DubiousAllen et al., 2015DubiousAllen, 1963DubiousAllen, 1973DubiousNofe, 1977DubiousMolfe, 1977DubiousCanaj, 1990DubiousYolfe, 1977DubiousMolfe, 1977DubiousMolfe, 1977DubiousCanagentio, 2000DubiousCanagentio, 2000DubiousCanagentio, 2000DubiousCanagentio, 2000DubiousCanagentio, 2000 <tr< td=""></tr<>
 Palacophytocrone lacanosam (MacCimitie) Wolfe Palacophytocrone lacanosam (MacCimitie) Wolfe Palacophytocrone spesialogenica Scott emend Manchester Palacophytocrone sp. Patrophytacrone sp. Phytocrone and changer (Chandler) Stall, Adams, Manchester et Collision Phytocrone and construction Stall. Phytocrone sp. Pyrenacantha austroomericana Stall Pyrenacantha austroomericana Stall Pyrenacantha austroomericana Stall Sicocarya comunais Reid and Chandler Sicocarya comunais Reid and Chandler Sicocarya comunais Reid and Chandler Goweria blachona Wolfe Goweria blachona Wolfe Goweria blachona Wolfe Goweria blachona Wolfe Goweria blachone econoum Tanai Phytocrene sordiar (Leasereux.) MacGimite Phytocrene sordiar sintanus Krutch & Vanhoorne 1977 Compositopolenites mediar Krutch & Vanhoorne 1977 Compositopolenites minimus Krutch & Vanhoorne 1977 Compositopolenites minimus Krutch & Vanhoorne 1977	Radecae Indecae Radecae Radecae Radecae Cancouver island Radocae Cancouver island R	CSAGermanyCanadaUSAUSAUSAUSAUSACanadaFanceCanadaCanadaDavaAlaskaUSACombiaColombiaPeruUSAGermanyGarmanyQiada AisagaQiada Aisaga <td> 37.8 33.9-56 33.9 43 33.9 56 32.2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈47 58 28.5 43 ≈47 ≈47 47.8 47.8</td> <td>EoceneEoceneEoceneOligoceneEocene</td> <td>? ? Lutetian Late Cuisian Sup. ? Middle-Late Paresian Iate early Lutetian Lutetian Lutetian Uateian Jare early Viresian Juetian Iate early Viresian Middle-Late Yiresian Middle-Late Middle Middle Middle Middle Middle Mide Mide Mide Mide Mide Mide Mide Mide <</td> <td>AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedPigg & Wehr, 2002AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedChandier, 1961b C (Scal et al., 2001)AcceptedCollisson, 2012AcceptedSuti et al., 2018AcceptedSuti et al., 2012AcceptedSuti et al., 2012AcceptedCollisson et al., 2012AcceptedCollisson et al., 2013AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNofic, 1967AcceptedNofic, 1968DubiousAloret al., 2017DubiousAloret al., 2017DubiousNofic, 1968AcceptedSuti et al., 2017DubiousNofic, 1967DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic,</td>	 37.8 33.9-56 33.9 43 33.9 56 32.2-33,6 33.9-56 49 33.9 33.9 47.8 37.8 66 ≈47 58 28.5 43 ≈47 ≈47 47.8 47.8	EoceneEoceneEoceneOligoceneEocene	? ? Lutetian Late Cuisian Sup. ? Middle-Late Paresian Iate early Lutetian Lutetian Lutetian Uateian Jare early Viresian Juetian Iate early Viresian Middle-Late Yiresian Middle-Late Middle Middle Middle Middle Middle Mide Mide Mide Mide Mide Mide Mide Mide <	AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedPigg & Wehr, 2002AcceptedMeyer & Manchester, 1997AcceptedDel Rio & De Franceschi in prep.AcceptedDel Rio & De Franceschi in prep.AcceptedChandier, 1961b C (Scal et al., 2001)AcceptedCollisson, 2012AcceptedSuti et al., 2018AcceptedSuti et al., 2012AcceptedSuti et al., 2012AcceptedCollisson et al., 2012AcceptedCollisson et al., 2013AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNofic, 1967AcceptedNofic, 1968DubiousAloret al., 2017DubiousAloret al., 2017DubiousNofic, 1968AcceptedSuti et al., 2017DubiousNofic, 1967DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic, 1977DubiousNofic,
 Palacophytocrane lacunomo (MacGinitie) Wolfe Palacophytocrane lacunomo (MacGinitie) Wolfe Palacophytocrane specialoperrica Scott emend Marchester Palacophytocrane sp. Phytocrane densignacobia Sull, Moore et Manchester Phytocrane densignacobia Sull, Moore et Manchester Phytocrane and coldentalis Manchester Phytocrane and coldentalis Manchester Sizocarya oriformia Reid and Chandler Sizocarya oriformia Reid and Chandler Sizocarya oriformia Reid and Chandler Generia ibustanis Volfe Generia ibustanis Volfe Phytocrane conflux (Endo) Tanai Compositophytolenites medias Krutch & Vanhoorne 1977 Compositophytolenites me	Radecan Indecand Radecan Cancouver island Rad	OSAGermanyUSAUSAUSAUSAUSAUSAGermanyFanceCanadaAlaskaUSAUSAUSAOremanyISAUSAUSAUSAOremanyISAOremanyISAOremanyISAOremanyINICH KingdomINICH KingdomI	 37.8 33.9-56 33.9 43 33.9 56 32,2-33,6 32,2-33,6 33.9 33.9 33.9 33.9 47.8 37.8 66 ≈47 58 28.5 43 ≈47 57-56 33.9 ≈46 49.5 33.9 ≈40 56 5	EoceneEoceneEoceneOligoceneEocene	? Lutetian Late Cuisian Sup. ? Middle-Late Paresian Lutetian Iarly-Late Lutetian Lutetian Lutetian Lutetian Lutetian Lutetian Lutetian Ypresian Kalde-Late Widdle-Late Middle-Late Middle	AcceptedFichy: 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedPigg & Wehr, 2002AcceptedDere et al., 1990AcceptedDere & Manchester, 1997AcceptedTichy: 1979AcceptedDer & Baka et al., 2008AcceptedPiele & Sall et al., 2001AcceptedOut & Bake et al., 2003AcceptedOut & Barghoorn, 1957AcceptedSall et al., 2012AcceptedSall et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedSall et al., 2013AcceptedReid & Chandler, 1933AcceptedSall et al., 2017AcceptedNarie, 1997AcceptedNarie, 1997AcceptedNarie, 1990DubiousAlore, 1967AcceptedSall, 1990DubiousAnai, 1990DubiousAnai, 1990DubiousAnai, 1990DubiousMarie, 1947AubiousNarie, 1947DubiousAnai, 1990DubiousAnai, 1990DubiousAnai, 1990DubiousArai, 1990DubiousArai, 1991DubiousCavagnetio
 Palaeophysicenee lacunome (AueGinike) Wolfe Palaeophysicenee lacunome (AueGinike) Wolfe Palaeophysicenee sp. Physicenee designation (Chandler) Stall, Adams. Marchester of Collision Physicenee designation (Chandler) Stall, Adams. Physicenee designation (Chandler) Stall, Adams. Physicenee sp. Pyrenaeontha sp. 1 Statecarya eophysicenas Stall and Chandler Statecarya eophysicenas Stall and Chandler Generici abadiant Wolfe Generici abadiant Wolfe Generici abadiant Wolfe Physicenee contilis (Endo) Tanai Physicenee contilis (Endo) T	Radeca Rideca Radeca Radeca Rade Radeca Rade Radeca Rade Radeca	OSAGermanyUSAUSAUSAUSAUSAUSAGarmanyFanceAlaskaUSAUSAOarmanyUSAOarmanyINAUSAOarmanyINAOarmanyINAOarmanyINAOarmanyINAOarmanyINAOarmanyINAOarmanyINA <td>37.833.9-5633.94333.95632.2-33,633.9-564933.947.837.866≈475828.543≈475828.543≈475828.543≈4757.5633.9≈4033.9≈4033.9≈4033.9≈4033.9≈40565850505152565656565656585056575658565856565656565656</td> <td>EoceneEoceneEocenePaleoceneEocene</td> <td>?LutetianLateQuisian Sup.?Middle-LateQuisian Sup.?Middle-LateQuietianLutetianLutetianQuietian</td> <td>AcceptedFichy: 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedDel Ko & De Francesch in prep.AcceptedDel Ko & De Francesch in prep.AcceptedNanken et al., 2008AcceptedOut & Nor, 1977AcceptedSull et al., 2010AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2013AcceptedCollinson et al., 2013AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNoffe, 1977AcceptedNoffe, 1977DubiousAlce, 1994AcceptedSull et al., 2015DubiousAlce, 1994AubiousNoffe, 1977DubiousCavagnetio, 2000DubiousSull et al., 2014AubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000D</td>	37.833.9-5633.94333.95632.2-33,633.9-564933.947.837.866≈475828.543≈475828.543≈475828.543≈4757.5633.9≈4033.9≈4033.9≈4033.9≈4033.9≈40565850505152565656565656585056575658565856565656565656	EoceneEoceneEocenePaleoceneEocene	?LutetianLateQuisian Sup.?Middle-LateQuisian Sup.?Middle-LateQuietianLutetianLutetianQuietian	AcceptedFichy: 1979AcceptedRankin et al., 2008AcceptedPigg & Wehr, 2002AcceptedPigg & Wehr, 2002AcceptedCrane et al., 1990AcceptedDel Ko & De Francesch in prep.AcceptedDel Ko & De Francesch in prep.AcceptedNanken et al., 2008AcceptedOut & Nor, 1977AcceptedSull et al., 2010AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2012AcceptedCollinson et al., 2013AcceptedCollinson et al., 2013AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedReid & Chandler, 1933AcceptedNoffe, 1977AcceptedNoffe, 1977DubiousAlce, 1994AcceptedSull et al., 2015DubiousAlce, 1994AubiousNoffe, 1977DubiousCavagnetio, 2000DubiousSull et al., 2014AubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000DubiousCavagnetio, 2000D
 Palaraphysic rane lacansam (McGinitky) Wolfs Palaraphysic rane lacansam (McGinitky) Wolfs Palaraphysic rane lacansam (McGinitky) Wolfs Palaraphysic rane sp. Physic rane and a materian science Manke bester Physic rane and a materian science Mail Pyrenacanha sp. 1 Pyrenacanha sp. 2 Sphaeriades variticions (Bowetbank) Reid and Chandler Sticocary a communit Reid and Chandler Sticocary a communit Reid and Chandler Converta Bhearist Mole Hasianashian ikhisaphoran Dal Ro & De Franceschi Goweria Bhearist Wolfe Hasianawa Wolfe Pyrenacanha sp. <li< td=""><td>Radecan Rindener Radecan Rindener Radecan Rindener Rinderan Rindener Rinder Rindener Rinder Rindener Rinder Rindener Rinder Rindener Rinder Rindener <tr< td=""><td>OSAGermanyCanadaUSAUSAUSAUSAUSACanadaCanadaCanadaCanadaUSAUSACanadaOramanyUSAOramanyO</td><td>37.833.9-5633.94333.95632,2-33,633,933,933,933,947.866≈475828.543≈475828.543≈47₹4033.9≈4649.533.9≈4033.9≈4033.9≈4033.9≈4033.9≈405656585359.2565833.9≈4751.65359.2565833.9≈4751.65356</td><td>EoceneEoceneFaleoceneCaleoceneEocene<!--</td--><td>?LutetianLateCuisian Sup.?Middle-LateParesianEarly-LateLutetianLutetianLutetianLutetianJare arlyJare sianJiresianYiresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMidle<td>AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedPigg, & Wehr, 2002AcceptedOran et al., 1990AcceptedRole, 1990AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedOran et al., 2008AcceptedOran et al., 2001AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1974AcceptedNofe, 1974AubioneNofe, 1974Aubion</td></td></td></tr<></td></li<>	Radecan Rindener Radecan Rindener Radecan Rindener Rinderan Rindener Rinder Rindener Rinder Rindener Rinder Rindener Rinder Rindener Rinder Rindener <tr< td=""><td>OSAGermanyCanadaUSAUSAUSAUSAUSACanadaCanadaCanadaCanadaUSAUSACanadaOramanyUSAOramanyO</td><td>37.833.9-5633.94333.95632,2-33,633,933,933,933,947.866≈475828.543≈475828.543≈47₹4033.9≈4649.533.9≈4033.9≈4033.9≈4033.9≈4033.9≈405656585359.2565833.9≈4751.65359.2565833.9≈4751.65356</td><td>EoceneEoceneFaleoceneCaleoceneEocene<!--</td--><td>?LutetianLateCuisian Sup.?Middle-LateParesianEarly-LateLutetianLutetianLutetianLutetianJare arlyJare sianJiresianYiresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMidle<td>AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedPigg, & Wehr, 2002AcceptedOran et al., 1990AcceptedRole, 1990AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedOran et al., 2008AcceptedOran et al., 2001AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1974AcceptedNofe, 1974AubioneNofe, 1974Aubion</td></td></td></tr<>	OSAGermanyCanadaUSAUSAUSAUSAUSACanadaCanadaCanadaCanadaUSAUSACanadaOramanyUSAOramanyO	37.833.9-5633.94333.95632,2-33,633,933,933,933,947.866≈475828.543≈475828.543≈47₹4033.9≈4649.533.9≈4033.9≈4033.9≈4033.9≈4033.9≈405656585359.2565833.9≈4751.65359.2565833.9≈4751.65356	EoceneEoceneFaleoceneCaleoceneEocene </td <td>?LutetianLateCuisian Sup.?Middle-LateParesianEarly-LateLutetianLutetianLutetianLutetianJare arlyJare sianJiresianYiresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMidle<td>AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedPigg, & Wehr, 2002AcceptedOran et al., 1990AcceptedRole, 1990AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedOran et al., 2008AcceptedOran et al., 2001AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1974AcceptedNofe, 1974AubioneNofe, 1974Aubion</td></td>	?LutetianLateCuisian Sup.?Middle-LateParesianEarly-LateLutetianLutetianLutetianLutetianJare arlyJare sianJiresianYiresianMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddle-LateMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMiddleMidle <td>AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedPigg, & Wehr, 2002AcceptedOran et al., 1990AcceptedRole, 1990AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedOran et al., 2008AcceptedOran et al., 2001AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1974AcceptedNofe, 1974AubioneNofe, 1974Aubion</td>	AcceptedFichy, 1979AcceptedRankin et al., 2008AcceptedPigg, & Wehr, 2002AcceptedOran et al., 1990AcceptedRole, 1990AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedDol Rin & Do Francesch in prep.AcceptedOran et al., 2008AcceptedOran et al., 2001AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1971AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedSoutier, 1973AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedReid, & Chandler, 1933AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1973AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1977AcceptedNofe, 1973AcceptedNofe, 1974AcceptedNofe, 1974AubioneNofe, 1974Aubion
 Representational control (SAUR) Mallice/SHI Palaeophytocerne laceaseum (McCGinike) Wolfe Palaeophytocerne pseudoperate Scott entend Manchester Palaeophytocerne pseudoperate Scott entend Manchester Palaeophytocerne pseudoperate Scott entend Manchester Palaeophytocerne ps. Palaeophytocerne ps. Palaeophytocerne ps. Palaeophytocerne ps. Palaeophytocerne ps. Palaeophytocerne optimus (Holl) Wolfe Palaeophytocerne optimus (Holl Adams, Manchester et Collinson Pathorene optimus (Holl Adams, Manchester Parocerne optimus (Holl Adams, Manchester Parocerne optimus (Holl Adams, Manchester Parocerne optimus (Holl Adams) Gaveria distanta Wolfe Gaveria biolaesis Tamai Palaeopholocore contain Tamai Palaeopholocore optimus (Holl Adams) Palaeopholocore optimus (Holl Adams) Palaeopholocore optimus (Kratch & Vanhoore 1977 Gamposhopholoches molinas Kratch & Vanhoore 1977 Gamposhopholoches moli	Radeace Ridence	CSAGermanyCanadaUSAUSAUSAUSACamanyFanceCanadaAlaskaUSAUSACarmanyISAOramanyISAOramanyISAOramanyISAOramanyISAOramanyISAOramanyISAOramanyISAOramanyISAIS	37.833.9-5633.94333.95632.2-33,633.9-564933.947.847.837.828.543≈475828.543≈475828.543≈4747.847.847.847.847.847.847.833.9≈4033.9≈4033.9≈4033.9≈40565658505658505658565656585658565658565656565656575658565856575658565856585658565856585658565856585658565756585956575658<	EoceneEoceneEoceneOligoceneEocene	?LutetianLutetianLate?Middle-LateQuerianEarly-LateLutetianLutetianLutetianLutetianJuresianQuerianiYiresianMiddle-LateYiresianY	AcceptedFalvy, 1979AcceptedRunkin et al., 2008AcceptedGuot, 1954; Manchester, 1994AcceptedUge & Wehr, 2002AcceptedChan et al., 1990AcceptedDalko & De Francesch in prep.AcceptedBankin et al., 2008AcceptedSull et al., 2018AcceptedSull et al., 2018AcceptedCander, 1961b; Cleal et al., 2011AcceptedCallison, 2012AcceptedSull et al., 2012AcceptedSull et al., 2012AcceptedCallison et al., 2013AcceptedCallison et al., 2013AcceptedCallison et al., 2014AcceptedCallison et al., 2017AcceptedCallison et al., 2015AcceptedCallison et al., 2014AcceptedCallison et al.,
 Langepolence in an encode (MAR) Matter BHT Palaesphysiocrene in anchorent italian, Stuckey et Beard Palaesphysiocrene sp. Physiocrene puncification (Chandke) Stull, Adams. Physiocrene sp. Phy	<table-container><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-container>	CSAGermanyCanadaUSAUSAUSAUSACananaGarmanyCanadaMasaUSAUSACanadaMaraneDasaOamanaUSAUSAUSAOamana <t< td=""><td> 33.9-56 33.9 43 33.9 56 32.2-33,6 33.9-56 49 33.9 47.8 37.8 66 ≈47 58 28.5 43 ≈47 ₹47 47.8 33.9 ≈ 40 56 53 533 533 533 </td><td>ForeeneeFo</td><td>? Lutetian Late Cuisian Sup. ? Middle-Late Paresian Lutetian Late carly Lutetian Lutetian Lutetian Vipresian Lutetian Vipresian Vipresian Widdle-Late Middle-Late Widdle-Late Vipresian Vipresian Widdle-Late Middle-Late Middle-Late Middle-Late Middle Vipresian Vipresian Vipresian Middle Vipresian Vip</td><td>AcceptedFieldy, 1979AcceptedScott, 1954, Manchester, 1994AcceptedQuer et al., 1990AcceptedCancet et al., 1990AcceptedCancet et al., 1990AcceptedDel Kio & De Francesch in prep.AcceptedScott, 1977, Hollick, 1936, Heer, 1870AcceptedScott, 1977, Hollick, 1936, Heer, 1870AcceptedScott, 2012AcceptedScott, 2013AcceptedScott, 2013AcceptedScott, 2014AcceptedScott, 2014<</td></t<>	 33.9-56 33.9 43 33.9 56 32.2-33,6 33.9-56 49 33.9 47.8 37.8 66 ≈47 58 28.5 43 ≈47 ₹47 47.8 33.9 ≈ 40 56 53 533 533 533 	ForeeneeFo	? Lutetian Late Cuisian Sup. ? Middle-Late Paresian Lutetian Late carly Lutetian Lutetian Lutetian Vipresian Lutetian Vipresian Vipresian Widdle-Late Middle-Late Widdle-Late Vipresian Vipresian Widdle-Late Middle-Late Middle-Late Middle-Late Middle Vipresian Vipresian Vipresian Middle Vipresian Vip	AcceptedFieldy, 1979AcceptedScott, 1954, Manchester, 1994AcceptedQuer et al., 1990AcceptedCancet et al., 1990AcceptedCancet et al., 1990AcceptedDel Kio & De Francesch in prep.AcceptedScott, 1977, Hollick, 1936, Heer, 1870AcceptedScott, 1977, Hollick, 1936, Heer, 1870AcceptedScott, 2012AcceptedScott, 2013AcceptedScott, 2013AcceptedScott, 2014AcceptedScott, 2014<
 Comparison of decompoint (Control NUMERINER) Palarophysion rule decompoint (Control Number 1997) Palarophysion rule operatory and control operanoty and control operatory and control operatory and coperan	Index <td>OSAGermanyICAUSAUSAUSAUSAISAISAICA<!--</td--><td> 33.9-56 33.9 43 33.9 56 32.2-33,6 49 33.9 49 33.9 49 33.9 47.8 37.8 66 ≈47 58 28.5 43 ≈47 47.8 33.9 ≈40 56 53 533 46.2 53 533 533 46.2 53 533 533 533 533 533 534 535 54 55 56 58 53 53 53 53 53 53 53 53 53 54 55 56 57 56 57 56 57 56 58 56 56 56 56 5</td><td>EacenesEacenesFaleocenesCacenesEacenesEacenesEacenesEacenesCacenesCacenesEa</td><td>? Lutcian Late Quisian Sup. ? Middle-Late Presian Lutcian Late carly Lutcian Lutcian Vipresian Vipresian Vipresian Vipresian Vipresian Vipresian Vipresian Widdle-Late Widdle Widdle</td><td>AcceptedFalsy, 1979AcceptedSotu, 1954, Manchester, 1994AcceptedQig, & Wahr, 2002AcceptedGrane et al., 1990AcceptedDirab, 2008AcceptedDirab, 2008AcceptedDirab, 2008AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 8 ary, boot, 1957AcceptedSotu, 1931AcceptedSotu, 1932AcceptedSotu, 1</td></td>	OSAGermanyICAUSAUSAUSAUSAISAISAICA </td <td> 33.9-56 33.9 43 33.9 56 32.2-33,6 49 33.9 49 33.9 49 33.9 47.8 37.8 66 ≈47 58 28.5 43 ≈47 47.8 33.9 ≈40 56 53 533 46.2 53 533 533 46.2 53 533 533 533 533 533 534 535 54 55 56 58 53 53 53 53 53 53 53 53 53 54 55 56 57 56 57 56 57 56 58 56 56 56 56 5</td> <td>EacenesEacenesFaleocenesCacenesEacenesEacenesEacenesEacenesCacenesCacenesEa</td> <td>? Lutcian Late Quisian Sup. ? Middle-Late Presian Lutcian Late carly Lutcian Lutcian Vipresian Vipresian Vipresian Vipresian Vipresian Vipresian Vipresian Widdle-Late Widdle Widdle</td> <td>AcceptedFalsy, 1979AcceptedSotu, 1954, Manchester, 1994AcceptedQig, & Wahr, 2002AcceptedGrane et al., 1990AcceptedDirab, 2008AcceptedDirab, 2008AcceptedDirab, 2008AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 8 ary, boot, 1957AcceptedSotu, 1931AcceptedSotu, 1932AcceptedSotu, 1</td>	 33.9-56 33.9 43 33.9 56 32.2-33,6 49 33.9 49 33.9 49 33.9 47.8 37.8 66 ≈47 58 28.5 43 ≈47 47.8 33.9 ≈40 56 53 533 46.2 53 533 533 46.2 53 533 533 533 533 533 534 535 54 55 56 58 53 53 53 53 53 53 53 53 53 54 55 56 57 56 57 56 57 56 58 56 56 56 56 5	EacenesEacenesFaleocenesCacenesEacenesEacenesEacenesEacenesCacenesCacenesEa	? Lutcian Late Quisian Sup. ? Middle-Late Presian Lutcian Late carly Lutcian Lutcian Vipresian Vipresian Vipresian Vipresian Vipresian Vipresian Vipresian Widdle-Late Widdle	AcceptedFalsy, 1979AcceptedSotu, 1954, Manchester, 1994AcceptedQig, & Wahr, 2002AcceptedGrane et al., 1990AcceptedDirab, 2008AcceptedDirab, 2008AcceptedDirab, 2008AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 1977, Holick, 1036, Heer, 1870AcceptedSotu, 8 ary, boot, 1957AcceptedSotu, 1931AcceptedSotu, 1932AcceptedSotu, 1
 Pakeophysicsense lacensem (MacGalle) Wolfe Pakeophysicsense lacensem (MacGalle) Wolfe Pakeophysicsense pesadopertica Scott enered Vanbeneer Pakeophysicsense pesadopertica Scott Eduals, Stockey et Baard Pakeophysicsense advancementeria Rathan, Stockey et Baard Pakeophysicsense advancementeria Statilitation (Chardise) Sull. Advans, Marchester et Collinam Paysocrear advancementeria Statilitation (Chardise) Sull. Advans, Marchester et Collinam Paysocrear advancementeria Statilitation (Chardise) Sull. Advans, Marchester et Mike Paysocrear advances (Bowchards Rati and Chardler Paysoc	Index </td <td>IONA<!--</td--><td> 33.9-56 33.9 43 33.9 56 32.2-33,6 32.2-33,6 32.2-33,6 33.9 49 33.9 33.9 47.8 37.8 28.5 43 ≈47 ₹47 ₹47 ₹47 ₹47 ₹47 ₹40 33.9 ≈40 56 56 56 56 56 58 56 33.9 41.2 33.9 ≈47 5.33 5.33 4.2 5.33 5.33 4.2 5.33 5.33 4.2 5.33 5.33 4.2 5.3 56 41.2 3.9 56 58 56 </td><td>EaceaneEaceaneFaleoceneCaceaneEaceaneEaceaneEaceaneEaceaneCaceaneCaceaneCaceaneCaceaneCaceaneEac</td><td>? Luterian Late Cuisian Sup. ? Middle-Late Bartonian Early-Late Luterian Luterian Luterian Paresian Vipresian Middle-Late Wiresian Vipresian Middle-Late Middle Middle Middle Noresian Vipresian Vipresian</td><td>AcceptedFaky, 1979AcceptedScoti, 1954; Manchester, 1994AcceptedPige & Wehr, 2002AcceptedGrane et al., 1990AcceptedDel Rin & Del Rin & Enserschi in prep.AcceptedDel Rin & Del Ri</td></td>	IONA </td <td> 33.9-56 33.9 43 33.9 56 32.2-33,6 32.2-33,6 32.2-33,6 33.9 49 33.9 33.9 47.8 37.8 28.5 43 ≈47 ₹47 ₹47 ₹47 ₹47 ₹47 ₹40 33.9 ≈40 56 56 56 56 56 58 56 33.9 41.2 33.9 ≈47 5.33 5.33 4.2 5.33 5.33 4.2 5.33 5.33 4.2 5.33 5.33 4.2 5.3 56 41.2 3.9 56 58 56 </td> <td>EaceaneEaceaneFaleoceneCaceaneEaceaneEaceaneEaceaneEaceaneCaceaneCaceaneCaceaneCaceaneCaceaneEac</td> <td>? Luterian Late Cuisian Sup. ? Middle-Late Bartonian Early-Late Luterian Luterian Luterian Paresian Vipresian Middle-Late Wiresian Vipresian Middle-Late Middle Middle Middle Noresian Vipresian Vipresian</td> <td>AcceptedFaky, 1979AcceptedScoti, 1954; Manchester, 1994AcceptedPige & Wehr, 2002AcceptedGrane et al., 1990AcceptedDel Rin & Del Rin & Enserschi in prep.AcceptedDel Rin & Del Ri</td>	 33.9-56 33.9 43 33.9 56 32.2-33,6 32.2-33,6 32.2-33,6 33.9 49 33.9 33.9 47.8 37.8 28.5 43 ≈47 ₹47 ₹47 ₹47 ₹47 ₹47 ₹40 33.9 ≈40 56 56 56 56 56 58 56 33.9 41.2 33.9 ≈47 5.33 5.33 4.2 5.33 5.33 4.2 5.33 5.33 4.2 5.33 5.33 4.2 5.3 56 41.2 3.9 56 58 56 	EaceaneEaceaneFaleoceneCaceaneEaceaneEaceaneEaceaneEaceaneCaceaneCaceaneCaceaneCaceaneCaceaneEac	? Luterian Late Cuisian Sup. ? Middle-Late Bartonian Early-Late Luterian Luterian Luterian Paresian Vipresian Middle-Late Wiresian Vipresian Middle-Late Middle Middle Middle Noresian Vipresian	AcceptedFaky, 1979AcceptedScoti, 1954; Manchester, 1994AcceptedPige & Wehr, 2002AcceptedGrane et al., 1990AcceptedDel Rin & Del Rin & Enserschi in prep.AcceptedDel Rin & Del Ri
 Lance, quences also and particular y Matcheeler Palacophysicscene anochos en Radia. Stocky of Board Palacophysicscene anochos en Radia. Stocky of Board Palacophysicscene appendiques also anothom technology Palacophysicscene appendiques also also anothom technology Palacophysicscene appendiques also also anothom technology Palacophysicscene acylorodynamic (Damber) Multi, Maters also allow also also anothom technology Palacophysicscene acylorodynamic (Damber) Multi, Alarma, Mulatobastic (Damber) Multi, Mulatobastic (Damber) Mulat	IndexIndexIndexIndexNonureinant	DescriptionGermanyUSAUSAUSAUSAUSAGermanyIanacaMaskaUSAUSAUSAUSAUSAOrandarianIARARIDANA	33.9-5633.9335632.2-33,633.9-564933.933.947.837.828.543≈47×47×47×47×47×40>3.9≈4033.9≈4133.9≈4756585633.9≈415.336033.934.25.336633.934.25.336633.934.25.35.365.36 <td>EaceaneEaceaneFaleoceneColoreneEaceaneEaceaneEaceaneEaceaneCoreaceCoreaceCoreaceEaceaneEa</td> <td>? Luterian Late Late Casisian Sup. ? Middle-Late Bartonian Early-Late Luterian Late carly Late carly Luterian Varesian Varesian Varesian Varesian Middle-Late Middle Naracian sup. Middle Naracian Naracian Middle Naracian Naracian Naracian Naracian Naracian Naracian</td> <td>AcceptedFolty, 1979AcceptedScot, 1954; Manchester, 1994AcceptedGara et al., 1990AcceptedGara et al., 1990AcceptedDel No & Del No.AcceptedDel No & Del No.AcceptedDel No.AcceptedCollector & Del No.AcceptedSull et al., 2011AcceptedSull et al., 2014AcceptedSull et al., 2014Accepte</td>	EaceaneEaceaneFaleoceneColoreneEaceaneEaceaneEaceaneEaceaneCoreaceCoreaceCoreaceEaceaneEa	? Luterian Late Late Casisian Sup. ? Middle-Late Bartonian Early-Late Luterian Late carly Late carly Luterian Varesian Varesian Varesian Varesian Middle-Late Middle Naracian sup. Middle Naracian Naracian Middle Naracian Naracian Naracian Naracian Naracian Naracian	AcceptedFolty, 1979AcceptedScot, 1954; Manchester, 1994AcceptedGara et al., 1990AcceptedGara et al., 1990AcceptedDel No & Del No.AcceptedDel No & Del No.AcceptedDel No.AcceptedCollector & Del No.AcceptedSull et al., 2011AcceptedSull et al., 2014AcceptedSull et al., 2014Accepte
 J. J. C. C. S. C. S. S.	<table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-row><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container><table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-container></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row></table-row>	ConversionGermanyUSAUSAUSAUSAUSACanadaGarmanyDiada fangaUSA<	3.3.9.33.9.33.933.933.9.33.9.33.9.33.9.33.9.33.9.33.9.33.9.33.9.47.833.9.33.9.33.9.33.9.33.9.33.9.34.033.9.34.133.9.34.033.9.34.133.9.34.033.9.34.033.9.34.033.9.34.133.9. <trr>34.2.35.3.3.35.3.3.</trr>	EaceaneEaceaneFaleoceneCaleoceneEaceaneEaceaneEaceaneEaceaneEaceaneCaleoceneCaleoceneEaceane<	?LutetianLutetianLateCuisian Sup.?Middle-LateEarly-LateLutetianLutetianLutetianJare sainJyresianUyresianMiddle-LateMiddleMiddle-LateMiddleMiddleJopersianMiddleMidleMidleMidleMidleMidleMidleMidleMidleMidleMidleMidleMidle </td <td>NecepterTedpy, 1979AccepterSout, 1934, Manchester, 1934AccepterGana et al., 1990AccepterDity A. Damebaster, 1937AccepterDity A. Damebaster, 1937AccepterDit M. Damebaster, 1937AccepterDit M. Damebaster, 1937AccepterDit M. Damebaster, 1937AccepterDitti M. Damebaster, 1937AccepterCambebaster, 1937AccepterCambebaster, 1937AccepterDitti A. Damebaster, 1937AccepterDitti M. Damebaster, 1937AccepterCalimon et al., 2012AccepterDitti A. Damebaster, 1937AccepterDitti A. Damebaster, 1933AccepterDitti A. Damebaster, 1937AccepterDitti A. Damebaster,</td>	NecepterTedpy, 1979AccepterSout, 1934, Manchester, 1934AccepterGana et al., 1990AccepterDity A. Damebaster, 1937AccepterDity A. Damebaster, 1937AccepterDit M. Damebaster, 1937AccepterDit M. Damebaster, 1937AccepterDit M. Damebaster, 1937AccepterDitti M. Damebaster, 1937AccepterCambebaster, 1937AccepterCambebaster, 1937AccepterDitti A. Damebaster, 1937AccepterDitti M. Damebaster, 1937AccepterCalimon et al., 2012AccepterDitti A. Damebaster, 1937AccepterDitti A. Damebaster, 1933AccepterDitti A. Damebaster, 1937AccepterDitti A. Damebaster,

Icacinoxylon nishidae Takahashi & Suzuki Wood Rumoi County Japan 89,8? Cretaceous Turonian? Rejected Takahashi & Suzuki, 2003

Icacinoxylon pittiense Thayn, Tidell & Stokes	Wood	Utah	USA	100.5	Cretaceous	Aptian-Albian	Rejected	Thay n et al., 1985
Icacinoxylon platanoides Greguss	Wood	Zirc	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon shilikinae Greguss	Wood	Pomaz	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp.	Wood	Colti-Buzau	Romania	28.1	Oligocene	Lower	Rejected	Petrescu et al., 1989
Icacinoxylon sp.	Wood	Pravaleni	Romania	5.33	Miocene	Sup.	Rejected	Petrescu & Nutu, 1972
Icacinoxylon sp. 1	Wood	Dorog	Hungary	23.03	Oligocene		Rejected	Greguss, 1969
Icacinoxylon sp. 2	Wood	Bajna	Hungary	23.03	Oligocene		Rejected	Greguss, 1969
Icacinoxylon sp. 3	Wood	Cserhathalap	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 4	Wood	Varpalota	Hungary	15.9	Miocene	Lower Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 5	Wood	Cinkota	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 6	Wood	Pecsszabolcs	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 7	Wood	Mor	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 8	Wood	Megyaszo	Hungary	11.6	Miocene	Sarmatian	Rejected	Greguss, 1969
Icacinoxylon sp. 9	Wood	Budapest	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 10	Wood	Harsagy	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 11	Wood	Bodajk	Hungary	11.6	Miocene	Lower Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 12	Wood	Iskaszent-gy orgy	Hungary	15.9	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 13	Wood	Homokbodoge	Hungary	15.9	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 14	Wood	Solymar	Hungary	23.03	Oligocene		Rejected	Greguss, 1969
Icacinoxylon sp. 15	Wood	Penzeskut	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp. 16	Wood	Ratka	Hungary	11.6	Miocene	Samatian	Rejected	Greguss, 1969
Icacinoxylon sp. 17	Wood	Nagybatony	Hungary	15.9	Miocene	Burdigalian	Rejected	Greguss, 1969
Icacinoxylon sp. 18	Wood	Nograd	Hungary	11.6	Miocene	Helvetian	Rejected	Greguss, 1969
Icacinoxylon sp.1	Wood	Potomac group	USA	93.9	Cretaceous	Albian-Cenomanian	Rejected	Herendeen, 1997
Icacinoxylon sp.2	Wood	Asen	Sweden	72.1	Cretaceous	Santonian-	Rejected	Herendeen, 1997
Icacinoxylon sylvaticum (Tuzon) Greguss	Wood	Ipoly tarnoc	Hungary	15.97	Miocene	Burdigalian	Rejected	Tuzson, 1906; Greguss, 1969
Icacinoxylon aff. sylvaticum (Tuzon) Greguss	Wood	Grateloup	France	28.1	Oligocene	Stampien	Rejected	Duperon, 1976

Icacninoxylon cantleyoides Gottwald Wood ? Belgium 83.6 Cretaceous Santonian Rejected Gottwald, 2000

Epoch	Number of species	Extinct genus	Modern genus
Oligocene	3	Palaeophytocrene	Iodes (Palaeohosiea) Pyrenacantha
Eocene	59	Faboidea Icacinanthium Icacinicarya Iodicarpa (Iodes?) Palaeophytocrene Perforatocarpum Sphaeriodes Stizocarya	Iodes (Iodes, Croomiocarpum, Palaeohosiea) Natsiatum Phytocrene Pyrenacantha
Paleocene	16	Icacinicarya Icacinicarytes Palaeophytocrene	Iodes (Iodes, Palaeohosiea, Hosiea) Phytocrene ?
Upper Cretaceous	1	Ø	Iodes

TABLE 2. Number of species of extinct and modern Icacinaceae accepted genera in different epochs. *Palaeohosiea*, *Hosiea* and *Croomiocarpum* species are considered as part of *Iodes*.