



Discovery of the only living population of *Pupoidopsis hawaiiensis* (Gastropoda: Pupillidae) in the last 50 years

Olivier Gargominy, Jean-François Butaud, Benoît Fontaine, Vincent Prié,
Dario Zuccon, Sandrine Tercerie, Joana Hauata, Maxime Hauata

► To cite this version:

Olivier Gargominy, Jean-François Butaud, Benoît Fontaine, Vincent Prié, Dario Zuccon, et al.. Discovery of the only living population of *Pupoidopsis hawaiiensis* (Gastropoda: Pupillidae) in the last 50 years. American malacological Bulletin, 2020, 38 (1), pp.50. 10.4003/006.038.0103 . mnhn-03084456

HAL Id: mnhn-03084456

<https://mnhn.hal.science/mnhn-03084456>

Submitted on 21 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH NOTE

Discovery of the only living population of *Pupoidopsis hawaiiensis* (Gastropoda: Pupillidae) in the last 50 years

Olivier Gargominy¹, Jean-François Butaud², Benoît Fontaines³, Vincent Prié⁴, Dario Zuccon⁴, Sandrine Terceries⁵, Joana Hauata⁶, and Maxime Hauata⁶

¹UMS 2006 PatriNat (OFB CNRS MN HN), Muséum national d'Histoire naturelle, CP41, 36 rue Geoffroy Saint-Hilaire 75005 Paris, gargo@mnhn.fr

²Consultant in forestry and Polynesian botany, BP 52832, 98716 Pirae, Tahiti, French Polynesia

³UMS 2006 PatriNat (OFB CNRS MNHN), Centre d'Écologie et des Sciences de la Conservation (UMR 7204), Muséum national d'Histoire naturelle, CP 135, 43 rue Buffon, 75005 Paris

⁴Institute of Systematics, Evolution, Biodiversity (ISYEB) (MNHN, CNRS, SU, EPHE, UA), Muséum national d'Histoire naturelle, CP 51, 57 rue Cuvier, 75005 Paris, France

⁵UMS 2006 PatriNat (OFB CNRS MNHN), Muséum national d'Histoire naturelle, CP41, 36 rue Geoffroy Saint-Hilaire 75005 Paris

⁶Pu Tahī Haga no Ganaa NGO, BP 60910, 98702 Faaa centre, Tahiti, French Polynesia

Abstract: *Pupoidopsis hawaiiensis* Pilsbry & C.M. Cooke, 1921, and its monospecific genus, is the only representative of Pupillidae in tropical Pacific Islands. Although most of native species from that region are single island endemics, *P. hawaiiensis* has been recorded from Hawaii to the Tuamotus. However, it has not been recorded alive since 1965. Here we report the only modern living population of that species from a single site in Anaa (Tuamotu archipelago). A single COI haplotype and a single 28S allele support the hypothesis of self-fertilization. The species should be listed as Critically Endangered, and included in the list of protected species of French Polynesia. An Action Plan for its conservation should be undertaken, with protection status given to its last known stronghold in Anaa.

Key words: upraised coral reefs, Tuamotu, Anaa, conservation

Pupoidopsis hawaiiensis Pilsbry & C.M. Cooke, 1921

(Pupillidae, Pupoidinae) is noteworthy because this species, the sole representative of the genus, is endemic to the Pacific Islands and has historically been reported with a wide distribution in this area, while most other native species are single island endemics. It is also the unique indigenous representative of the Pupillidae in tropical Pacific islands. First described as an extinct genus and species from Oahu, Molokai, and Maui in the Hawaiian islands (Pilsbry 1920-1921), it is now known from Hawaii, Line islands (Kiribati) and Tuamotu archipelago (French Polynesia). It has only been found alive four times: in Kiritimati – Christmas (Line islands, Kiribati) in 1924 (Anonymous 1926; Cooke and Neal 1928) and 1965 (Christensen 2013) and in Tabuaeran – Fanning (Line islands, Kiribati) and Hao (Tuamotu) in 1934 (Christensen 2013).

All collectors report the species as belonging to the lowland fauna, either on high islands or atolls. The heavy human impact on lowland habitats is thought to have been the cause of extirpation of *P. hawaiiensis* from its former distribution (Gargominy and Meyer 2012).

Here we report the presence of *P. hawaiiensis* from Anaa (Tuamotu) for the first time, and document the only currently known living population of this species.

MATERIALS AND METHODS

A malacological survey of Anaa (Tuamotu Islands, French Polynesia) was conducted January 4 - 18, 2018, on behalf of the NGO Pu Tahī Haga no Ganaa and funded by BEST 2.0 (Gargominy *et al.* 2018). Data were collected from 40 localities, representing a total of 676 lots, 22,000 specimens and 38 taxa. Specimens are deposited in the collection of the Muséum national d'Histoire naturelle (Paris, France), under numbers MNHN-IM-2018-46, -71, -80, -129, -141, -302, -333, -491, -527, -562, -584, -624 and MNHN-IM-2013-76001 to 76005.

Five specimens of *P. hawaiiensis* were prepared for molecular analyses (MNHN-IM-2013-76001 to 76005). Total genomic DNA was extracted using the Qiagen DNA Mini Kit and following the manufacturer's protocol. Two standard genes fragments were selected for PCR amplification: the mitochondrial cytochrome oxidase I (COI, amplified with primers LCO1490 and HCO2198, Folmer *et al.* 1994) and the nuclear large subunit rRNA gene (28S, amplified with primers C1 and D2, Jovelín and Justine 2001). Sequences were compared to other Pupillidae sequences available in Genbank. We assembled two datasets: the COI (655bp) and the 28S (784bp) datasets, including 10 and nine other Genbank Pupillidae sequences, respectively. We used Valloniidae sequences as outgroup. Both datasets were analysed under the maximum likelihood criterion, using RAXML v. 7.0.3 (Stamatakis 2006), selecting a GTR+Γ+I model and random starting tree, with empirical base frequencies

and estimated α -shape parameters and GTR-rates. Nodal support was estimated using 100 bootstrap replicates.

RESULTS

Pupoidopsis hawaiiensis shells were collected in one fourth of the collecting sites (11 stations out of 40). These stations are distributed all over the island (Fig. 1), but the species was recorded in greater abundance on the south-western part of the island (Otepipi, Okuku, Hapuku-Putuahara and Tekahora islets). *Pupoidopsis hawaiiensis* was found alive at a single station on the motu Otepipi (Figs. 1, 2, 3). The area was no more than 100 square meters. It was quite abundant, with up to ten individuals fixed under a single limestone pebble a dozen centimeters in diameter. It was only found on the ground under rocks. At this site, the vegetation was dominated by the native trees and shrubs *Allophylus rhomboidalis* (Nadeaud) Rdlk., *Cyclophyllum barbatum* (G.Forst.) N.Hallé & J.Florence, *Glochidion wilderi* J.Florence, *Guettarda speciosa* L., *Pisonia grandis* R.Br., *Planchonella tahitensis* (Nadeaud) Pierre ex Dubard, *Premna serratifolia* L., *Sophora tomentosa* L., *Timonius uniflorus* (Banks ex C.F.Gaertn.) Govaerts, *Xylosma suaveolens* (J.R.Forst. & G.Forst.) G.Forst. The understory was very scarce on this rocky station, with native ferns *Asplenium polyodon* G.Forst. and *Microsorium grossum* (Langsd. & Fisch.) S.B.Andrews.

Among the five sequenced specimens, we recovered a single COI haplotype and a single 28S allele (GenBank accession numbers MT457074-MT457078 and MT457405- MT457409, respectively). The phylogenetic reconstruction places *Pupoidopsis hawaiiensis* in the Pupillidae (Fig. 4). In the COI tree *Pupoidopsis* is sister to *Pupoides albilabris*, while in the 28S tree it is basal to a clade containing *Pupoides albilabris*, *Pupilla* and *Gastrocopta*, albeit with no bootstrap support.

DISCUSSION

Although the phylogenetic reconstruction confirms that *Pupoidopsis hawaiiensis* belong to the Pupilloidea, the limited number of published sequences available in Genbank prevent us to clarify its place within the family. A better resolution requires a larger, multigene dataset and the inclusion of the type species of *Pupoides* (*Bulimus nitidulus* L. Pfeiffer, 1839, type locality in Cuba).

The fact that among the five sequenced specimens, we recovered a single COI haplotype and a single 28S allele is congruent with Cooke and Neal (1928) who observed that most specimens are female by lacking penis, and we suggest that the species could reproduce by self-fecundation. This probable self-fertilization, allowing colonization of a new island with a single individual, is congruent with the exceptionally wide distribution of the species within the Pacific.

Anaa is an uplifted atoll, i.e. typically composed of lowland habitats either on sand or on rocky calcareous substrate. Preferred habitat for *P. hawaiiensis* in this island is *feo*, which corresponds to areas of upraised limestone (coral reef) up to few meters high, or areas neighbouring *feo* with great density of limestone pebbles on the ground. *Feo* represent areas where vegetation is in the best state of conservation on this atoll, as these areas were not converted to coconut plantations due to the rocky substrate. This relatively good state of conservation of *feo* probably accounts for the fact that *P. hawaiiensis* still survives there. However, we are unable to explain why the species has been extirpated everywhere in Anaa except in a single place on Otepipi. No rat predation on this species is recorded, but it is worth noting that just as elsewhere in Anaa, Otepipi harbours two rat species (*Rattus exulans* and *R. rattus*).

As a comparison, the land snail *Mautodontha daedalea* (A.A. Gould, 1846), endemic to Makatea and Anaa atolls in the Tuamotu, was found alive on five stations in Anaa (with 12 additional stations where shells only were found). Despite the fact it belongs to the Endodontidae, a family which was much impacted by habitat perturbations (Solem 1976 ; Abdou and Bouchet 2000 ; Zimmermann *et al.* 2009 ; Sartori *et al.* 2013 ; 2014), it is still more represented than *P. hawaiiensis*. *Pupoidopsis hawaiiensis* may still survive in Anaa in other stations, but the contrast between the station on Otepipi, where live individuals were abundant, and other stations where only empty shells were found suggests that if it is still extant somewhere else, its abundance is much lower than it used to be. We hypothesize that it has disappeared from 10 out of 11 stations, i.e. a reduction of more than 90% of its former range on this single atoll.

As stated by Christensen (2013), some islands in Kiribati and Tuamotu may still support living populations of *P. hawaiiensis*, as it is now proved for Anaa. However, the species must be highly threatened, since it is extinct in most of its known former range (Kiribati, Hawaiian archipelago – Christensen (2013)) and close to extinction in Anaa. As for most invertebrates which are under-studied, we cannot rule out the fact that it is still extant somewhere else. However, it has never been found in French Polynesia, where extensive collections have been done (for instance, Society islands, Austral islands, Gambier islands, Makatea – e.g. Gargominy 2008, Christensen and Kahn 2017, Sartori *et al.*, 2013, 2014, Abdou and Bouchet 2000). Additional surveys should be

undertaken in neighboring islands (especially Niau, another uplifted atoll) to find the species. *Pupoidopsis hawaiiensis* is currently classified as data deficient DD in the IUCN Red List (Cowie and Pokryszko 1996). We propose that it should be listed as Critically Endangered B1,B2ab (i,ii,iii,iv,v) (only one living population on a single atoll and general decline of extent of occurrence, area of occupancy, quality of habitat, number of locations, number of mature individuals). We recommend that it should be included in the list of protected species of French Polynesia (Anonymous 2018), and an Action Plan for its conservation should be undertaken. The *feo* of Otepipi could also be given a protection status, for instance through the general management plan (PGA) of the Anaa municipality. *Feo* also provide strong protection against natural disasters as barriers to the ocean.

ACKNOWLEDGMENTS

We thank the European Commission which founded part of this work through BEST 2.0 project 1264 “Sauvegarde de la flore menacée de l’atoll de Anaa et définition de zones prioritaires de conservation basées sur la flore et la malacofaune sur les atolls soulevés de Anaa et Niau aux Tuamotu”. We thank also Maima Matai and Mare Paeahi who helped for malacological collections in the field. Frédéric Alain Jacq prepared the map of Figure 1. This work was supported by ‘Service de Systématique Moléculaire’ (UMS2700 Acquisition et Analyse de Données pour l’Histoire naturelle, MNHN). We also thank two anonymous reviewers for helpful comments on the manuscript.

LITERATURE CITED

- Abdou, A. and P. Bouchet. 2000. Nouveaux gastéropodes Endodontidae et Punctidae (Mollusca, Pulmonata) récemment éteints de l’archipel des Gambier (Polynésie). *Zoosystema* **22**: 689–707.
- Anonymous. 1926. *Pupoidopsis hawaiiensis* on Christmas Island. *The Nautilus* **39**: 144.
- Anonymous. 2018. Arrêté n°466 CM du 22 mars 2018. Available at <http://lexpol.cloud.pf/LexpolAfficheTexte.php?texte=515575>. 29 September 2019.
- Christensen, C. 2013. *Pupoidopsis hawaiiensis* Pilsbry & Cooke, 1921 (Gastropoda: Pupillidae): extirpated in Hawai’i but a possible survivor in Kiribati and French Polynesia. *Bishop Museum Occasional Papers* **114**: 43–48.
- Christensen, C. C. and J. G. Kahn. 2017. First records of the invasive predatory land snail *Gulella* (*Huttonella*) *bicolor* (Hutton, 1834) (Gastropoda: Streptaxidae) from the Society Islands, French Polynesia. *Bishop Museum Occasional Papers* **121**: 1–11.
- Cooke, C. M. and M. C. Neal. 1928. Distribution and anatomy of *Pupoidopsis hawaiiensis*. *Bernice P. Bishop Museum Bulletin* **47**: 28–33.
- Cowie, R. H. and B. Pokryszko. 1996. *Pupoidopsis hawaiiensis*. The IUCN Red List of Threatened Species 1996: e.T18920A8700770. Available at: <http://dx.doi.org/10.2305/IUCN.UK.1996.RLTS.T18920A8700770.en>. 12 January 2019.
- Folmer, O., M. Black, W. Hoeh, R. Lutz, and R. Vrijenhoek. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* **3**: 294–299.
- Gargominy, O. 2008. Beyond the alien invasion: A recently discovered radiation of Nesopupinae (Gastropoda: Pulmonata: Vertiginidae) from the summits of Tahiti (Society Islands, French Polynesia). *Journal of Conchology* **39**: 517–536.
- Gargominy, O., B. Fontaine, S. Tercerie, and V. Prié. 2018. *Inventaire des Mollusques continentaux d’Anaa. Rapport de mission sur Anaa du 4 au 18 janvier 2018*. Pu Tahī Haga no Ganaa, Anaa & Muséum national d’Histoire naturelle, Paris, France.
- Gargominy, O. and J. Y. Meyer. 2012. Disjunct distribution in the Pacific Ocean: *Pupoidopsis hawaiiensis* in French Polynesia, Kiribati and Hawaii, South Pacific Islands. *Journal of Conchology* **41**: 129–130.
- Jovelín, R., and J. L. Justine. 2001. Phylogenetic relationships within the polyopisthocotylean monogeneans (Platyhelminthes) inferred from partial 28S rDNA sequences. *International Journal of Parasitology* **31**: 393–401.
- Pilsbry, H. A. 1920-1921. *Manual of conchology. Second series: Pulmonata. Vol. XXVI. Pupillidae (Vertigininae, Pupillinae)*. Academy of Natural Sciences, Philadelphia, USA.
- Sartori, A. F., O. Gargominy, and B. Fontaine. 2013. Anthropogenic extinction of Pacific land snails: A case study of Rurutu, French Polynesia, with description of eight new species of endodontids (Pulmonata). *Zootaxa* **3640**: 343–372.

Sartori, A. F., O. Gargominy, and B. Fontaine. 2014. Radiation and decline of endodontid land snails in Makatea, French Polynesia. *Zootaxa* **3772**: 1–68.

Solem, A. 1976. *Endodontoid Land Snails from Pacific Islands (Mollusca: Pulmonata: Sigmurethra). Part 1. Family Endodontidae*. Field Museum Press, Chicago, USA.

Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**: 2688–2690.

Zimmermann, G., O. Gargominy, and B. Fontaine. 2009. Quatre espèces nouvelles d'Endodontidae (Mollusca, Pulmonata) éteints de Rurutu (I les Australes, Polynésie française). *Zoosystema* **31**: 791–805.




Figure 1. Map showing the locations where *Pupoidopsis hawaiiensis* was found. Black circles: living population, grey circles: shells only, white circles: absent.



Figure 2. Live *Pupoidopsis hawaiiensis* in place under a stone on the ground.



Figure 3. Live *Pupoidopsis hawaiiensis*.




Figure 4. Maximum-likelihood trees based on COI (top) and 28S sequences (bottom). The bootstrap support values (>50%) are indicated at the node.