



Three new combinations and one lectotypification of fern and lycophyte taxa from the French overseas territories

Germinal Rouhan, Vincent Boullet, Ashley Field, Eric Schuettpelz

► To cite this version:

Germinal Rouhan, Vincent Boullet, Ashley Field, Eric Schuettpelz. Three new combinations and one lectotypification of fern and lycophyte taxa from the French overseas territories. *Phytotaxa*, 2021, 497 (1), pp.54-56. 10.11646/phytotaxa.497.1.6 . mnhn-03201065

HAL Id: mnhn-03201065

<https://mnhn.hal.science/mnhn-03201065>

Submitted on 19 Apr 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Three new combinations and one lectotypification of fern and lycophyte taxa from the French overseas territories

GERMINAL ROUHAN^{1,*}, VINCENT BOULLET^{2,3}, ASHLEY FIELD^{4,5} & ERIC SCHUETPELZ⁶

¹ Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, Sorbonne Université, École Pratique des Hautes Études, CNRS, Université des Antilles; CP 39, 57 rue Cuvier, 75005 Paris, France; germinal.rouhan@mnhn.fr; <https://orcid.org/0000-0002-0751-5352>

² Conservatoire Botanique National de Mascarin, 2 rue du Père Georges, Les Colimaçons, 97436 Saint-Leu, La Réunion; France; v.boulet43@orange.fr; <https://orcid.org/0000-0002-3482-1662>

³ EA 7462 Géoarchitecture, Université de Bretagne Occidentale, 29200 Brest.

⁴ Queensland Herbarium, Department of Environment and Science, Brisbane Botanic Gardens, Mount Coot-tha, Mt Coot-tha Road Toowong, Qld 4066, Australia; ashley.field@jcu.edu.au; <https://orcid.org/0000-0002-0139-7914>

⁵ Australian Tropical Herbarium, Sir Robert Norman Building, James Cook University Campus, Smithfield Queensland 4878, P.O .Box 6811 Cairns, Qld 4870, Australia.

⁶ Department of Botany, National Museum of Natural History, Smithsonian Institution, Washington, DC, U.S.A.; SchuettpelzE@si.edu; <https://orcid.org/0000-0003-3891-9904>

*Corresponding author

With the exception of French Guiana and Adélie Land, the French Overseas Territories (FOTs) are islands and their biodiversity is remarkable in many ways. Notably, they harbour numerous unique taxa leading to exceptionally high rates of endemism. Among French endemic plants, 95% occur in the FOTs, accounting for ca. 4000 species of tracheophytes and ‘bryophytes’ (Gargominy *et al.* 2020).

We recently initiated a project entitled "Flore Endémique des Territoires d'Outre-Mer" (i.e., Endemic Flora of the French Overseas Territories; FEnTOM) to summarize the present state of knowledge of plant endemism in the FOTs. Using the most current information and data, we intend to build an up-to-date checklist of all known strict and regional endemic spermatophytes, ‘pteridophytes’ and ‘bryophytes’ known in the FOTs (Veron *et al.* in prep.). In compiling this list, we discovered that three nomenclatural combinations remain outstanding for ferns and lycophytes, and we here present them to avoid the use of invalid combinations in the FEnTOM project and to also make appropriate names available in the French national taxonomic database TAXREF (Gargominy *et al.* 2020), as well as international taxonomic databases. These new combinations are consistent with the classification presented in PPG I (2016).

Antrophyopsis gigantea (Bory) Rouhan, Bouillet & Schuettp., *comb. nov.*

Basionym: *Antrophyum giganteum* Bory (1833: 36)

Type:— Mauritius. Le Pouce, *C. Bélanger s.n.* (lectotype P [P00483382!], hic designatus; isolectotypes P [P00483381!], P [P01482666!]).

Notes:— This species was not included in the study of vittarioid ferns by Schuettpelz *et al.* (2016) and its placement in the phylogeny has yet to be assessed. Nonetheless, it is morphologically most similar to *Antrophyopsis boryana* (Willdenow 1810: 128) Schuettpelz (2016: 717) that was included in the molecular analysis and confirmed as belonging to

Antrophyopsis (Benedict 1907: 447) Schuettpelz (2016: 717). It is distinguished from *A. boryana* by sessile, subelliptic to subob lanceolate fronds, but both share the generic characters, notably spherical apical cells of the soral paraphyses and ecostate laminae (i.e. lacking midribs). Although *A. gigantea* was described from Mauritius, it was never observed there again (Badré 2008) and is today known only from La Reunion. As there are three sheets housed at P, we designated as lectotype the sheet with complete leaves given that the base of fronds is critical for distinguishing the species; also, the same sheet bears the more complete labels.

Lepisorus mucronatus (Fée 1852: 82) Li Wang (2010: 35) var. ***durus*** (Copel.) Rouhan, *comb. nov.*

Basionym: *Hymenolepis dura* Copeland (1938: 69, t.20).

Synonym: *Belvisia mucronata* (Fée 1852: 82) Copeland (1947: 192) var. *dura* (Copel.) Hovenkamp & Franken (1993: 523).

Type:—French Polynesia: “Australes”, Rapa, versant sud du Mont Lekie, 330 m, 20 July 1934, *H. St. John & J. Maireau 15621* (holotype BISH [BISH1000333, online image]; isotypes BISH [BISH1000335, online image], BO, K [K000880383!], MICH [1191320, online image], P [P00717795!], UC [UC 542661, online image], US [00134710, online image], W).

Notes:— *Belvisia* Mirbel (1803: 65) is entirely subsumed in *Lepisorus* (Smith 1846: 13) Ching (1933: 47, 56) following studies by Wang *et al.* (2010). The variety *durus* is known from Rapa Island in French Polynesia.

Phlegmariurus pseudovarius (Brownlie) Rouhan & A.R.Field, *comb. nov.*

Basionym: *Lycopodium pseudovarium* Brownlie (1969: 23, t.1, f.13-15).

Synonym: *Huperzia pseudovaria* (Brownlie) Holub (1991: 93).

Type:— New Caledonia. Prov. Sud: Mont Humboldt, 1200 m, February 1872, *B. Balansa* 3558 (holotype P [P00523013!]).

Notes:— Following the classification proposed by Field *et al.* (2016) and later retained by PPG I (2016), this species formerly placed in *Huperzia* Bernhardi (1801: 126) belongs in *Phlegmariurus* Holub (1964: 21). It is endemic to New Caledonia, but may possibly also occur in French Polynesia according to recent observations (J. Nitta, pers. com.).

The type at P was determined in 2015 by A. Field as *Phlegmariurus varius* (Brown 1810: 165) A.R.Field & Bostock (2013: 49) due to the lack of the combination *P. pseudovarius*, and because he thought it may be a synonym of *Phlegmariurus varius*. More recent work provides evidence supporting *P. pseudovarius* as distinct, along with at least two other species in this complex (Field, unpublished data).

Acknowledgements

We thank the Office Français de la Biodiversité (French Agency of Biodiversity) and the Ministère des Outre-mer (French Ministry of Overseas Territories) for funding and supporting this project.

We are also grateful to RECOLNAT national Research Infrastructure (ANR-11-INBS-0004).

References

- Badré, F. (2008) Ptéridophytes 1. Psilotacées à 26. Marsiléacées. In: Bosser, J., Badré, F. & Guého, J. (eds.) *Flore des Mascareignes - La Réunion, Maurice, Rodrigues*. IRD, MSIRI, RBG-Kew, Paris. 454 pp.
- Benedict, R.C. (1907) The genus *Antrophyum*—I. Synopsis of subgenera, and the American species. *Bulletin of the Torrey Botanical Club* 34(9): 445–458.
<http://dx.doi.org/10.2307/2479065>
- Bernhardi, J.J. (1801) Tentamen alterum filices in genera redigenda. *Journal für die Botanik (Schrader)* 1800: 121–136.
- Bory de Saint-Vincent, J.B.G.M. (1833) In: Bélanger, C., Bory de Saint-Vincent, J.B.G.M. & Montagne, C. (eds.) *Voyages aux Indes-Orientales par le nord de l'Europe : les provinces du Caucase, la Géorgie, l'Arménie et la Perse, suivis de détails topographiques, statistiques et autres sur les Pégot, les îles de Java, de Maurice et de Bourbon, sur le Cap-De-Bonne-Espérance et Sainte-Hélène ; pendant les années 1825–1829, Botanique II*. Paris. 192 pp.
- Brown, R. (1810) *Prodromus Florae Novae Hollandiae et Insulae van-Diemen*, exhibens characteres plantarum quas annis 1802–1805 per oras utriusque insulae collegit et descripts Robertus Brown; insertis passim aliis speciebus auctori hucusque cognitis, seu evulgatis, seu ineditis praesertim Banksianis, in primo itinere navarchi Cook detectis, 1. J. Johnson & Co., London. 590 pp.
- Brownlie, G. (1969) N°3 Ptéridophytes. In: Aubréville, A. (ed.), *Flore de la Nouvelle-Calédonie et Dépendances*. Muséum national d'Histoire naturelle, Paris. 308 pp.
- Ching, R.-C. (1933) The studies of Chinese ferns IX. *Bulletin of the Fan Memorial Institute of Biology* 4(3): 47–110.
- Copeland, E.B. (1938) Ferns of southeastern Polynesia. *Occasional Papers of Bernice Pauahi Bishop Museum of Polynesian Ethnology and Natural History* 14(5): 45–101.
- Copeland, E.B. (1947) *Genera Filicum: the genera of ferns*. Chronica Botanica Company of Waltham, Mass. 247 pp.
- Fée, A.L.A. (1850–1852) *Mémoires sur les Familles des Fougères; Cinquième Mémoire, Genera Filicum*. Paris, Strasbourg, 387 pp.
- Field, A. & Bostock, P. (2013) New and existing combinations in Palaeotropical *Phlegmariurus* (Lycopodiaceae) and lectotypification of the type species *Phlegmariurus phlegmaria* (L.) T.Sen & U.Sen. *PhytoKeys* 20: 33–51.
<https://doi.org/10.3897/phytokeys.20.4007>
- Field, A.R., Testo, W., Bostock, P.D.B., Holtum, J.A.M. & Waycott, M. (2016) Molecular phylogenetics and the morphology of the Lycopodiaceae subfamily *Huperzioideae* supports three genera: *Huperzia*, *Phlegmariurus* and *Phylloglossum*. *Molecular Phylogenetics and Evolution* 94: 635–657. <https://doi.org/10.1016/j.ympev.2015.09.024>
- Gargominy, O., Tercerie, S., Régnier, C., Dupont, P., Daszkiewicz, P., Léotard, G., Antonetti, P., Ramage, T., Vandel, E., Petitteville, M., Leblond, S., Idczak, L., Boullet, V., Denys, G., De Massary, J.C., Lévêque, A., Jourdan, H., Rome, Q., Dusoulier, F., Touroult, J., Savouré-Soubelet, A., Barbut, J., Canard, A., Simian, G., Le Divelec, R., Haffner, P., Meyer, C., Van Es, J., Poncet, R., Demerges, D., Mehran, B., Horellou, A., Moulin, N., Ah-Peng, C., Bernard, J.-F., Caesar, M., Comolet-Tirman, J., Courtecuisse, R., Delfosse, E., Dewynter, M., Hugonnot, V., Kondratyeva, A., Lavocat Bernard, E., Lebouvier, M., Lebreton, E., Malécot, V., Moreau, P.A., Muller, S., Noblecourt, T., Pellens, R., Robbert Gradstein, S., Rodrigues, C., Rouhan, G. & Véron, S. (2020) TAXREF v14.0, référentiel taxonomique pour la France. UMS PatriNat, Muséum national d'Histoire naturelle, Paris. Archive de téléchargement contenant 8 fichiers.

Available from:

<https://inpn.mnhn.fr/telechargement/referentielEspece/taxref/14.0/menu/> (accessed 28 Feb. 2021)

Holub, J. (1964) *Lycopodiella*, novy rod radu Lycopodiales. *Preslia* 36: 16–22.

Holub, J. (1991) Some taxonomic changes within Lycopodiales. *Folia Geobotanica & Phytotaxonomica* 26(1): 81–94. <https://doi.org/10.1007/BF02912943>

Hovenkamp, P.H. & Franken, N.A.P. 1993. An account of the fern genus *Belvisia* Mirbel (Polypodiaceae). *Blumea: Biodiversity, Evolution and Biogeography of Plants* 37(2): 511–527.

Mirbel, C.F.B. de (1803) *Histoire naturelle générale et particulière des plantes* 4. Paris, 431 pp.

PPG I (2016) A community-derived classification for extant lycophytes and ferns: PPG I. *Journal of Systematics and Evolution* 54: 563–603. <https://doi.org/10.1111/jse.12229>

Schuettpelz, E., Chen, S.-W., Kessler, M., Pinson, J.B., Johnson, G., Davila, A., Cochran, A.T., Huiet, L. & Pryer, K.M. (2016) A revised generic classification of vittarioid ferns (Pteridaceae) based on molecular, micromorphological, and geographic data. *Taxon* 65(4): 708–722. <http://dx.doi.org/10.12705/654.2>

Smith, J. (1846) Additions to the “Hortus Kewensis”. *Botanical Magazine* 72 (Comp.): 1–42.

Veron, S., Rodrigues-Vaz, C., Lebreton, E., Ah-Peng, C., Boulet, V., Chevillotte, H., Gradstein, S.R., Jérémie, J., Lavocat Bernard, E., Lebouvier, M., Meyer, J.-Y., Munzinger, J., Poncy, O., Thouvenot, L., Viscardi, G., Léotard, G., Gargominy, O., Leblond, S., Pignal, M., Rouhan, G., Tercerie, S., Invernón, V. & Muller, S. (accepted). *Biodiversity and Conservation*.

Wang, L., Wu, Z.Q., Xiang, Q.P., Heinrichs, J., Schneider, H. & Zhang, X.C. (2010) A molecular phylogeny and a revised classification of tribe *Lepisoreae* (Polypodiaceae) based on an analysis of four plastid DNA regions. *Botanical Journal of the Linnean Society* 162: 28–38.

Willdenow, C.L. (1810) *Species Plantarum*. Editio Quarta, 5. Berolini [Berlin], 542 pp.