



HAL
open science

The middle Permian tracksite of Gonfaron: A complex ecosystem in the midst of the end-Guadalupian mass-extinction

Antoine Logghe, Eudald Mujal, Lorenzo Marchetti, Pierre Barrier, Christian Montenat, André Nel, Jean-Marc Pouillon, Romain Garrouste, Jean-Sébastien Steyer

► To cite this version:

Antoine Logghe, Eudald Mujal, Lorenzo Marchetti, Pierre Barrier, Christian Montenat, et al.. The middle Permian tracksite of Gonfaron: A complex ecosystem in the midst of the end-Guadalupian mass-extinction. European Association of Vertebrate Paleontologists, Jun 2022, Benevento (IT), France. mnhn-03978538

HAL Id: mnhn-03978538

<https://mnhn.hal.science/mnhn-03978538v1>

Submitted on 21 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ABSTRACT BOOK



19th

EAVP

CONFERENCE 2022

Benevento / Pietraroja Italy

27.06 – 02.07.2022

Co-organised by and with the patronage of:



In collaboration with:



and supported by:



How to cite an abstract from this volume:

Yurac Diaz M., Salazar C., Meyer C.A., Suárez M.E, Belvedere M. 2022. The dinosaur ichnological record of northern Chile: a review and its potential development. In: Belvedere M., Mecozzi B., Amore O., Sardella R (eds.). Abstract book of the XIX Annual Conference of the European Association of Vertebrate Palaeontologists, Benevento/Pietraroja, Italy, 27th June-2nd July 2022. *PalaeoVertebrata*, Special Volume 1-2022, 224. Doi: 10.18563/pv.eavp2022

THE MIDDLE PERMIAN TRACKSITE OF GONFARON: A COMPLEX ECOSYSTEM IN THE MIDST OF THE END-GUADALUPIAN MASS EXTINCTION

A. Logghe^{1*}, L. Marchetti¹, E. Mujal^{2,3}, P. Barrier⁴, C. Montenat^{4†}, A. Nel⁵, J.-M. Pouillon⁶,
R. Garrouste⁵, J.-S. Steyer⁷

¹ Museum für Naturkunde Berlin, Leibniz-Institut für Evolutions- und Biodiversitätsforschung, Invalidenstrasse 43, 10115 Berlin, Germany.

² Staatliches Museum für Naturkunde Stuttgart, Rosenstein 1, D-70191 Stuttgart, Germany.

³ Institut Català de Paleontologia M. Crusafont, ICTA-ICP building, c/ de les columnes, s/n, E-08193 Cerdanyola del Vallès, Catalonia, Spain.

⁴ Département géosciences (IGAL), institut polytechnique LaSalle-Beauvais, 19, rue Pierre-Waguet, BP 30313, 60026 Beauvais cedex, France.

⁵ Institut de Systématique, Evolution, Biodiversité, UMR 7205 – CNRS, MNHN, SU EPHE, Muséum National d'Histoire Naturelle, Sorbonne Universités, 57 rue Cuvier, CP 50, Entomologie, F-75005, Paris, France.

⁶ Rhinopolis Association, Nivolas Vermelle, France.

⁷ Centre de Recherches en Paléontologie de Paris, UMR 7202 – CNRS, MNHN, SU, EPHE, Muséum National d'Histoire Naturelle, Sorbonne Universités, 8 rue Buffon, CP38, F-75005 Paris, France.

*presenting author, antoine.logghe@orange.fr

Keywords: *Capitanian, ichnology, locomotion, captorhinomorph, full-body impression*

The Permian outcrops of Provence yield one of the most extensive records of continental ichnofossils, reaching from the Kungurian through the Capitanian (early-middle Permian). The Pélitique Formation of the Gonfaron site (Provence, France) has recently been revised and is marked by a diverse vertebrate ichnofauna: *Batrachichnus salamandroides* (temnospondyls/lepospondyls), *Capitosauroides talus* comb. nov. (therocephalian therapsids), *Dicynodontipus* isp. (cynodont therapsid), *Varanopus* isp. (bolosaurian parareptiles), *Hyloidichnus bifurcatus* (captorhinomorph eureptiles) and *Rhynchosauroides* isp. (neodiapsid eureptiles). This ichnoassociation has been dated to the late Capitanian and assigned to the newly introduced *Dicynodontipus* sub-biochron. It is also the earliest evidence of the post-dinocephalian extinction recovery at low latitudes of Pangaea. Moreover, the Gonfaron site is an excellent site for the study

of Permian ecosystems: it yielded two rare full-body impressions of Hexapoda that testify certain ecological strategies, such as mimetics and jumping adaptation, possibly indicating high trophic pressure in harsh palaeoenvironmental conditions. The trophic pressure was investigated through the analysis of co-occurrences of fossil traces (arthropod and tetrapod tracks and trackways), which show that the association of tetrapod and hexapod imprints was relatively common. Also, the good preservation of the tetrapod tracks of the Pélitique Formation allowed the first synapomorphy-based correlation between *Hyloidichnus* tracks and captorhinomorph trackmakers and a precise reconstruction of the locomotion of captorhinomorph eureptiles. Further studies and new excavations on the site are planned to provide more data/material and strengthen our knowledge of this crucial period in the evolution of tetrapods and hexapods.