



HAL
open science

Temporal analysis of plant community changes documenting climate and urban practice impacts in cities. The case of Paris (France) over a 136-year period

Machon Nathalie, Gallant-Canguilhem Matthieu, Ropars Lise, Jeanne Vallet

► To cite this version:

Machon Nathalie, Gallant-Canguilhem Matthieu, Ropars Lise, Jeanne Vallet. Temporal analysis of plant community changes documenting climate and urban practice impacts in cities. The case of Paris (France) over a 136-year period: Comparison Parisian plant communities 1884-2021. 2023. mnhn-04096935

HAL Id: mnhn-04096935

<https://mnhn.hal.science/mnhn-04096935>

Preprint submitted on 14 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 ORIGINAL ARTICLE

2 **Temporal analysis of plant community changes documenting climate and urban practice**
3 **impacts in cities. The case of Paris (France) over a 136-year period.**

4 **Running title:** Comparison Parisian plant communities 1884-2021

5

6 **Machon Nathalie^{1*}, Gallant-Canguilhem Matthieu¹, Ropars Lise¹, Vallet Jeanne²**

7 ¹ *Centre d'Ecologie et des Sciences de la Conservation, Muséum national d'Histoire*
8 *naturelle, Sorbonne Université, Centre National de la Recherche Scientifique, 57 rue Cuvier,*
9 *75005 Paris, France*

10 ² *Conservatoire Botanique National du Bassin Parisien, Muséum national d'Histoire*
11 *naturelle, 57 rue Cuvier, 75005 Paris, France*

12 *For correspondence: E-mail nathalie.machon@mnhn.fr

13

- 1 • **Background and Aims:** Besides its intrinsic value, urban biodiversity participates to city
2 dwellers' better health and quality of life. One method to define which factors are driving
3 biodiversity is to determine the changes in communities between two time-periods and
4 confront them with the environmental and/or practice changes that have occurred at the
5 same time.
- 6 • **Methods:** We compared the changes that occurred in the Paris (France) plant communities
7 at a 136-years interval, thanks to two inventories performed in exactly the same streets,
8 wharves and squares, according to exactly the same protocol. We examined which species
9 disappeared, endured and appeared from the 19th century onwards, determined the traits of
10 these different groups of species and deduced which factors could have driven the shifts in
11 the composition of the Parisian weed communities.
- 12 • **Key Results:** The total number of species (about 190 species) remained relatively similar
13 over a 136-year timespan. Nevertheless, a turnover of about 50% of the species happened.
14 The differences between the 1884 and 2021 floristic data mostly followed climatic and
15 human activity changes i.e. the current flora seems to be better adapted to a warmer
16 climate and shifts in soil qualities. The differences in weeding practices also modeled the
17 new plant communities.
- 18 • **Conclusions:** This study is documenting changes in plant communities over such a long
19 period. Generally, comparable lists of urban data are scarce because former naturalists
20 rarely studied city areas and seldom detailed reproducible protocols. In our case, the
21 protocol was strictly reproducible and accurate enough to enable a comparison. The
22 results inform on the impact of global and local changes on biodiversity. It also provides
23 important clues about how to better integrate biodiversity in urban projects for the
24 ecosystem to function better and to improve future city dwellers' well-being.

1 **Key words:** Functional traits, streets, docks, squares, Joseph Vallot, Ellenberg indicators,
2 Jaccard's distances.

3

4 **Introduction**

5 Urbanization is one of the principal reason for land use changes worldwide to bring about
6 biodiversity alterations (Maxwell *et al.*, 2016). However, biodiversity is present in cities.
7 Besides its intrinsic value, it also participates to city dwellers' better health and quality of life
8 (Aerts, Honnay and Van Nieuwenhuysse, 2018). Territorial collectivities have to preserve it in
9 order to participate to the general conservation of biodiversity and to give citizens the best
10 possible life-conditions. For these reasons, it is necessary to determine what the main drivers
11 of the composition of natural urban communities are, and particularly the composition of
12 plant communities, which are key elements for biodiversity. Their state shapes the quality of
13 all networks and food chains interactions. In cities, they are composed of both cultivated and
14 spontaneous species, and among them, native and exotic ones. Cultivated species are species
15 planted by humans for food or ornamental purposes. Spontaneous ones are those that grow
16 naturally, originated locally or that landed by themselves from outside the city, or even spilled
17 over from gardens.

18 As conceptualized by (Aronson *et al.*, 2016), spontaneous urban vegetation is the result of
19 processes by which species filter which bare human disturbances and city environment. Most
20 plant species in cities dwell in diverse habitats (Lundholm and Marlin 2006) including parks,
21 public gardens (Shwartz *et al.*, 2013), flowerbeds, lawns, river banks, railways and green
22 spaces at business sites (Serret *et al.*, 2014). Among these habitats, public spaces such as
23 streets, squares and riverbanks offer a multitude of micro-green spaces that can harbor a
24 number of plant species, especially when management and trampling are limited (Pellegrini
25 and Baudry 2014);(Schmidt, Poppendieck, and Jensen 2014). In streets of cities like Paris –

1 except for alignment trees, which are planted mainly to decorate and provide shade in summer
2 – plants are rather wild and grow spontaneously in every crack in pavements, walls and roads.
3 Up till now, few studies have described the species that grow in such public spaces and have
4 determined the drivers of their communities' compositions (Omar et al. 2018) (Lundholm and
5 Marlin 2006).

6 One method to define which factors are driving biodiversity is to determine the changes in
7 communities between two time periods and confront them with the environmental and/or
8 practice changes having occurred at the same time, and examine how adjustments may have
9 taken place (Grimm et al. 2008). Thus, comparing urban plant communities, observed at more
10 than a one-century interval, make up extremely informative case studies of the effects of local
11 and global changes on biodiversity (Salinitro et al. 2019a).

12 Trait-based approaches can even more accurately provide general rules for community
13 ecology (Cadotte, Carscadden, and Mirotnick 2011), as well as a more mechanistic
14 understanding of community assembly and disassembly – particularly in the context of global
15 climate change (Mcgill *et al.*, 2006), (Williams *et al.*, 2019). Thus, in our study, reproductive
16 and ecological traits have been investigated to explain interspecific variations at temporal
17 scale, in response to environmental and anthropogenic changes as well.

18 Among ecological traits, Ellenberg's indicator values (EIV) (Ellenberg 1991) describe in
19 which environmental conditions plant species grow in Central Europe (Diekmann 2003).
20 They can be used as a numerical system to classify species' habitat niches (Silvertown et al.
21 2006) i.e., what their peak of occurrence are in terms of light availability (L), temperature (T),
22 atmospheric (Ma) and soil moisture (Ms), soil reaction (R), soil fertility (N) and salinity (S).
23 Furthermore, dependence of plant species on pollinators is also a trait that can characterize
24 their ecological needs. For each of these traits, their means per community can be calculated.
25 Comparison in time or space can characterize changes among communities.

1 This paper presents the changes that occurred in the Paris plant communities at a 136-year
2 interval, thanks to the existence of an inventory published in 1884 in a book entitled “Essay
3 on the flora of the Paris pavement limited to outer boulevards, or Catalog of plants that grow
4 spontaneously in streets and on wharves” (Vallot 1884), by a well-known French botanist
5 Joseph Vallot. In 2021, we followed in his footsteps, using the same inventory protocol at
6 exactly the same places. We used the trait-based approach to understand how environmental
7 changes may have affected biodiversity from that time onwards.

8 (Salinitro et al. 2019b) have listed different studies published on the same subject in other
9 European cities. None of them held such precise data, performed exactly the same protocol in
10 exactly the same places and used species’ characteristics to inform the recorded changes and
11 thus characterize global changes’ impacts on communities. Our study is more precise than
12 previous ones in that it was based on data related to the abundance of species by their
13 presence in a large sample of streets, wharves and squares in Paris, and not just in general, as
14 was the case, for example, in Italy in Bologna (Salinitro et al. 2019b), Brussels in Belgium
15 (Godefroid 2001), Plzen in the Czeck Republic (Chocholoušková and Pyšek 2003) or Zurich
16 in Switzerland (Landolt E 2000).

17 [Figure 1]

18 Comparing both available datasets, the present paper reports the changes that happened over
19 136 years in the plant communities found in 52 public spaces, all found in Paris, France:
20 streets, squares and Seine river’s wharves. The study only focused on the spontaneous flora
21 and excluded cultivated/ornamental vegetation. The habitats sampled all belonged to the
22 micro-structures that support vegetation, i.e., cracks in asphalt, the base of alignment trees, the
23 feet of walls, the spaces between paving stones... We examined which species were growing
24 in the Parisian public spaces in the late 19th century, which ones disappeared, endured, and
25 emerged from that time on. We determined the traits of these different groups of species and

1 deduced which factors could have driven the shifts in Parisian weed communities'
2 compositions.

3 **MATERIALS AND METHODS**

4 **Study site**

5 The study is located in the capital city of France (the World Geodetic System 1984 reference
6 is 48° 50' 26.91" N, 2° 23' 17.46" E), within the limits Paris used to have just before the
7 annexation of the bordering cities, which took place in 1860 (map fig 1, list Annex 1). The
8 area covered an area of approximately 3.5 km², in the city's densest districts.

9 During the 19th century, on emperor Napoleon III's command, the Paris center was deeply
10 modified by Baron Haussmann (Haussmann 1890). He opened wide avenues, created many
11 grounds and green spaces, built pavements for pedestrians, planted trees along the main
12 streets and erected the famous Paris Haussmann buildings. He resurfaced the streets, wharves
13 and squares by re-paving and asphaltting the main roads. By 1884, the work was completed.

14 In 1884, the human density was approximately 80,000 habitants/km². It was lower in 2021,
15 when there were around 20,500 inhabitants/km² (according to INSEE i.e. the French National
16 Institute for Statistics and Economic Studies, 2017, <http://www.insee.fr>).

17 In 2021, the heritage of Baron Haussmann is still significant. Urbanization has not changed
18 much in 136 years. Districts have more or less the same shape today as they used to at the end
19 of the 19th century. All the streets that existed in late 19th century still exist, except one (rue
20 Bunant).

21 **Floristic data**

22 **1884's inventories**

1 In 1884, Joseph Vallot published the list of the plants growing in 52 public spaces (PS): 35
2 streets, 15 riverbanks and 2 squares in Paris (France), comprised in the study site. He
3 collected the data in all the habitats in streets, squares and river wharves: 1° in the cracks
4 between cobblestones in streets and quays, on banks; 2° stones on quays; 3° plants on small
5 walls; 4° the rare bare soils; 5° the soil around tree bases along boulevards and avenues.

6 Among the plants inventoried, we removed all trees and shrubs because we considered they
7 had probably been planted. We also removed the only mention of a non-angiosperm plant
8 (i.e., *Equisetum arvense*). We have updated each species' names according to the French list
9 TAXREF 15.0 (<https://inpn.mnhn.fr/telechargement/referentielEspece/taxref/15.0/menu>).
10 Only one species namely *Carex muricata* L., observed by Vallot, remains questionable as it is
11 a historical name covering several taxa (*Carex otrubae*, *divulsa*, *leersii*, *pairae*, *spicata*...).

12 Joseph Vallot did not mention when he carried out his inventories. As he did not record any
13 spring or autumn species, we assumed that he collected his data over one growing season
14 between May and August, over a single year, around 1884.

15 **2021's inventories**

16 We researched the names and locations of each public space (PS) (street, wharf and square)
17 visited by Vallot. Some of them had changed name since then, but only one street had to be
18 discarded (Rue Bunant), as it no longer existed in 2021. We thus carried out our inventories in
19 all streets but one of them within the same limits as Vallot's.

20 We listed all the herbaceous angiosperm species between May and August 2021, except
21 ornamental species. When necessary, we used the app Pl@ntNet® to ensure the validity of
22 our identifications.

1 A matrix of the presence/absence of each species, in each of the sampled locations, at each
2 period: in Annex 1, it is presented with species in the rows; and in the columns, the
3 streets/wharves/squares as found in 1884 or 2021.

4 Regarding the native status, we used the database from the National Botanical Conservatory
5 of Parisian Basin CBNBP (<http://cbnbp.mnhn.fr/cbnbp/>), which indicates whether each
6 species is considered as native or exotic in France.

7 **Ellenberg indicator values and species dependency on pollinators**

8 For each species, seven Ellenberg indicator values were collected using the R package TR8
9 (Bocci 2015): temperature, light, nutrient, soil reaction and humidity, atmospheric humidity,
10 and salinity indicator values from the French database Baseflor (Julve Philippe 1998), and
11 also their dependence on pollinators for their reproduction (SPD). Ellenberg's temperature
12 indicator values are species-specific indices describing preferences along climatic gradients
13 (Ellenberg 1991). They characterize the optimum temperature class for a species' growth and
14 survival, among 9 classes ranging from "alpine to nival, altimediterranean floor" (T-number =
15 1) to "thermomediterranean to subarid floor" (T-number = 9). The soil reaction gradient
16 ranges from 'high acidity, never moderately acidic or alkaline' (R- number= 1) to 'alkaline
17 and calcareous conditions, only calcareous soils' (R- number = 9). The soil moisture covers
18 species growing on dry soils (Ms- number = 1) to species growing on wet, hypoxic soils
19 (Ms- number = 9). EIVs for nitrogen/nutrients span species growing on poor soils
20 (N- number 1) to species growing on nutrient- rich soils (N- number = 9). Light species
21 growing in deep shadow have an EIV of (L- number = 1), while species growing in full light
22 have an EIV of (L- number = 9). For all these indicators, we calculated the community-mean
23 preferences per street in 1884 and in 2021 (hereafter CMP) i.e., the mean of EIV of all species
24 inventoried in each street (Garnier *et al.*, 2004).

25 We also classified species on the basis of their dependence level to pollinators for their

1 reproduction. Following the methods of (Duchenne, Martin, and Porcher 2021), plants were
2 scored as more or less pollinator dependent. Pollen vectors (i.e., insects, wind or self-
3 fertilization) were retrieved from the Baseflor (Julve Philippe 1998) Ecoflora
4 (<http://ecoflora.org.uk/> Fitter and Peat 1994) and BiolFlor (www.biolflor.de Klotz, Kühn, and
5 Durka 2002; Kühn, Durka, and Klotz 2004) trait-bases. To integrate the sometimes-
6 contradictory data indicated by the different databases, the authors calculated a single variable
7 as the percentage of times “insects” proved to be a pollen vector for a given species, across all
8 databases. Their “species pollinator dependence” (SPD) was 100% when they were reportedly
9 completely pollinator dependent, i.e., allogamous and self-incompatible. It was 0% when they
10 were reported as only reproduced by selfing and/or pollinated by the wind. For each street,
11 wharf and square, we also calculated the mean dependence to pollinators and compared them
12 between the two periods.

13 All the data were treated using R packages *rstatix* (<https://rpkgs.datanovia.com/rstatix/>) and
14 (<http://haozhu233.github.io/kableExtra/>). The tests performed on mean Ellenberg indicator
15 values and mean dependence to pollinators by public space were Wilcoxon’s paired series
16 median comparison tests, since both normality and homoscedasticity hypotheses were not
17 validated for the 1884 data set.

18 **Floristic distances among streets and among wharves between the two periods**

19 To properly characterize inventories in relation to each other, we calculated pairwise
20 Jaccard’s distances between the 1884 and 2021 street floras, using the R *vegan* package
21 (<https://github.com/vegandevs/vegan>). With those distances, we drew the two dendrograms
22 using the R *dendextend* package (<https://cran.r-project.org/package=dendextend>) with Ward’s
23 method, which minimizes the intraclass inertia and maximizes the interclass inertia in order to
24 obtain the most homogeneous closures and subgroups possible.

25

1 **RESULTS**

2 **General floristic trends**

3 [Figure 2]

4 A total of 282 species had been listed in Paris over both inventories (Fig 2). One third (95 sp.)
5 were only found by Vallot in 1884 and not in the 2021 inventory. Among them, according to
6 the French databases (MNHN & OFB (Ed) 2003), eighteen of them disappeared from Paris
7 during the 20th century. One third (94 sp.) were only found in 2021 and had not been seen by
8 Vallot. Among them, 46 appeared in Paris during the 20th (39 sp.) or 21st centuries (7 sp.). And
9 one third (93 sp.) were found in both inventories. The Dice coefficient (Dice 1945) i.e. their
10 similarity was thus 0.496.

11 In the 1884 pool of species, 10.70 % were exotic (i.e., arrived in France after 1492). In 2021,
12 this rate was 23.53%.

13 The mean number of species growing in these streets and the wharves increased a lot, from
14 5.23 sp and 30.52 sp respectively in 1884 to 32.05; and 40,47 sp in 2021. In both squares,
15 species richness remained relatively identical, with a mean of 31 and 31.5 species.

16 **Mean trait analyses**

17 [Table 1]

18 [Figure 3]

19 Analyses of comparisons of mean Ellenberg indicator values of plants in the Parisian public
20 spaces detected significant differences between street, wharf and square communities between
21 the two periods (Fig 3). In particular, they revealed a significant increase in temperature
22 preference between the one in the 19st century to the other in 21st (the mean EIV passing
23 rising from 5.21 to 5.69, p-value < 0.001). In the same way, analyses revealed that 21st
24 century plants preferred drier (EIV passing from 4.75 to 4.73, p-value=0.013) and more
25 alkaline soils (mean EIV passing from 5.82 to 6.24, p < 0.001). Dependence on pollinators

1 also seemed to have increased between these periods (the mean dependence increasing from
2 36.39% to 45.17%). Preferences for light, salinity, atmospheric humidity and nitrate did not
3 seem to have changed significantly between these two periods (tab 1 and figure 3).

4

5 Floristic distances

6 [Figure 4]

7 [Figure 5]

8 Except for one street inventoried in 2021, we found a clear differentiation of floristic
9 composition between these two periods in streets and Seine wharves (Figures 4 and 5). Each
10 street and wharf were floristically more similar to the other streets and wharves of its period
11 than to itself at a 136-year interval. The distribution of species is completely changed between
12 the two periods. The streets that were floristically similar are not the same currently. The
13 average floristic distance is greater in streets in 1884 than in those in 2021, probably due to
14 the low number of species per street found in the 19th century.

15

16 **Discussion**

17 The inventory performed by Vallot in 1884 followed an intense urbanization process that
18 occurred over the 19th century and completely changed the shape of the city. Our inventories
19 occurred in a very similar territory, from an urbanistic point of view. Thus, differences
20 between the 1884 and 2021 floristic data mostly followed climatic and human activity
21 changes. Contrary to other studies on other cities conducted worldwide, the present results
22 pointed out that the total number of species (about 190 species) remained relatively similar
23 over a 136-year timespan. Nevertheless, there was a turnover of about 50% of species, only
24 half of them identical, and the other half exclusive to one of the two period inventories.
25 However, even if some of those exclusive species already existed or still exist in Paris, they

1 were only found in one of the two inventories. To explain such a turnover, it should be noted
2 that beyond their mere presence, species abundance, and thus detectability, may have changed
3 with time. Usually, variations of species abundances are due to stochastic processes or are
4 linked to environmental changes or practices. In any case, a study of the databases containing
5 all the data from inventories carried out in Paris over time from the 17th century onwards,
6 suggested this: 18 species became extinct in Paris; 39 (species) emerged during the 20th
7 century and 7 in the 21st century. In particular, the number of alien species doubled
8 (representing from 10.70% to 23.53% of total species).

9 This 50% turnover is lower than turnovers found in Bologna and in Plzeň in Czech Republic
10 (Salinitro et al. 2019b) and (Chocholoušková and Pyšek 2003) where similarity indices were
11 only 35% and 43% in 120 years and in 60 years exceeded the ones in Brussels (a 79.48%
12 similarity) (Godefroid 2001).

13 In contrast to these previous studies in other cities, we directly compared the plant community
14 indices in public spaces between the two time periods. This provides more accurate
15 information than simply comparing species' average preferences in the overall lists. Actually,
16 we incorporated variations in species abundance at the city level and highlighted more subtle
17 processes than just the appearance or disappearance of species.

18 Floristic distances among streets and wharves during both periods revealed high communities'
19 differentiation over more than one century (fig 3 and 4). The flora in each space resembles
20 that in other spaces of its time, much more than does its own flora at 136-year intervals.

21 The most obvious sign of the current flora's adaptation to a warmer climate is the increase in
22 mean EIV values concerning plant communities' temperatures (Fig 2). Some species, because
23 they were well-suited to cold climates, disappeared from Paris (e.g., *Carum carvi* or *Turritis*
24 *glabra*) or displayed reduced abundance. For example, *Centaurea jacea* and *Fallopia*
25 *convolvulus* decreased their occurrences by 2.5 between 1884 and 2021 (3,9 [to 1.6])

1 Conversely, species well-suited to hot climates arrived recently from warmer regions in the
2 world (e.g., *Senecio inaequidens*, from South Africa) or their abundance increased (e.g.,
3 *Hordeum murinum* multiplied its occurrences by 2.5 between these two periods).

4 Changes in city planning also impacted street plant communities. For example, communities'
5 better tolerance to drier soils is observed, though annual rainfall amounts increased very
6 slightly over one century, from 605 to 615 mm/year on average, between 1884 and 2021,
7 according to MétéoFrance (i.e. the French organism for climate survey
8 <https://meteofrance.com/>) data. However, maximum rainfall also increased (from 29.33 to
9 40.63 mm in 136 years, according to MétéoFrance), as did soil impermeability, with paved
10 areas being transformed into asphalted areas. This transformation mainly followed the May
11 1968 demonstrations, during which students used paving stones as projectiles to attack law
12 enforcement officers. As a result, more rainwater ran off into sewer systems and rivers,
13 probably leaving little water available for soil and plant hydration. Wharves have also been
14 channelized, leading to the disappearance of more or less aquatic species, like *Rorippa x*
15 *anceps* and *amphibia*, or *Rumex hydrolapathum*. Recently, *Portulaca oleracea* and *Senecio*
16 *inaequidens*, plants particularly adapted to dried soils, have invaded the French capital's
17 pavements.

18 In the same way, the massive use of construction materials has caused the alkalization of
19 (cement, concrete, etc...) soils. Plant communities seem to be more and more adapted to basic
20 soils. Among the plants specializing in acid soils, *Scleranthus annuus* disappeared from Paris
21 more than a century ago, and *Rumex acetosella* is now very rare in streets.

22 In contrast to what authors had found in the other cities (Godefroid 2001), (Chocholoušková
23 and Pyšek 2003) and (Salinitro *et al.*, 2019b) we did not find any changes in communities'
24 tolerance for light. It should be noted that buildings did not change in size and streets kept the
25 same width from the end of 19th century onwards. Trees had also been planted before Vallot's

1 inventories. Light has therefore probably not changed since then. However, as in other cities,
2 no differences were found concerning communities' sensitivity to atmospheric humidity and
3 salinity (Godefroid 2001).

4 Community characterization in terms of pollinator dependence revealed a significant
5 difference between the two periods (around 40% on average). According to (Ropars et al.
6 2019), cities harbor diverse plant species that flourish all year long because they apply low
7 pesticide policies and are subject to heat island effect. Hence, they provide resources for
8 pollinators throughout the year and create favorable conditions for the maintenance of diverse
9 pollinator communities (Harrison and Winfree 2015). We wish we could have compared these
10 rates with other data for other cities or other environment types, but it seems they do not yet
11 exist in the literature.

12 Here are the three specific human activities supposed to have modified public spaces' plant
13 communities: means of transport; food trade; and weed control. In more than one century,
14 horse-drawn vehicles were replaced by combustion engine automobiles. On the one hand,
15 with the disappearance of horses from Parisian streets, their fodder and dung no longer
16 provide abundant natural fertilizers. On the other hand, current car emissions tend to enrich
17 the soil with nitrates. Thus, preference for soil richness does not seem to have changed over
18 136 years. Nevertheless, plants from horse forage seeds, such as *Avena sativa* and *Phleum*
19 *pratense*, are no longer observed in Paris.

20 This species turnover detected in Paris over 136 years may also reflect the changes in food
21 trade experienced as of the early 20th century. It is estimated that 95% of the fruit and
22 vegetables consumed in the capital were produced more or less locally. Indeed, the market
23 garden production areas constituted a green belt, justified by the size of the city's population;
24 hence of the Parisian market; and also by the use of the city's waste as fertilizer (Moreau and
25 Daverne 1845). The streets adjacent to these gardens probably received the seeds produced by

1 crops there, thus extending their range to the most urbanized areas. These discharges from
2 gardens may also have occurred during the transport of foodstuffs or seeds to points of sale
3 such as markets or traders' carts. Thus, Vallot was, in his inventory, able to list lentils (*Lens*
4 *culinaris*) or asparagus (*Asparagus officinalis*) in some of the streets – vegetables that we did
5 not find during our inventorying.

6 Weeding and street cleaning are probably the practices with the greatest impact on
7 biodiversity changes in Paris as of the 19th century. Over paved streets, sweeping machines
8 were used when possible, otherwise they were cleaned by hand, by teams of men and women
9 equipped with efficient brushes or birch brooms, on an everyday basis (Vallot 1884)
10 (Hausmann 1890). Since then, pavements have gradually been covered with bitumen or tar
11 and the local authority has continued to weed streets and pavements with more or less drastic
12 techniques, depending on financial and human resources, as well as inhabitants' and borough
13 councils' sensitivities. Herbicides have been increasingly implemented. On 1 January 2017,
14 the adoption of the Labbé law, which aims to better control the use of phytosanitary products
15 on the French territory, greatly modified public spaces maintenance. Discontinuing the use of
16 chemical products has led Paris to manage its spaces differently. According to Parisian public
17 spaces' managers, weeding is more seasonal and therefore much less frequent (2 or 3 times a
18 year on average) than in the 19th century. It has become mainly mechanical, with the help of
19 tools capable of tearing and brushing the pavement.

20 Thus, our results showed that Parisian pavements and wharves hosted a richer flora in the 21st
21 century than they did in the 19th, with more species depending on pollinators. We should
22 specify, however, that inventories were carried out during the COVID-19 pandemic crisis,
23 when street use and management may have had a specific effect on biodiversity's quality.
24 Nevertheless, since the abandonment of herbicides on cities' public spaces, the flora has
25 already been found to be enriched.

1 **Conclusion**

2 This study is documenting changes in plant communities over such a long period. The
3 existence of comparable lists of data are scarce because former naturalist rarely studied urban
4 areas and seldom used or outlined reproducible protocols in detail. In our case, the protocol
5 was strictly reproducible and sufficiently precise to allow comparison.

6 A welcome surprise: the present Parisian communities were richer than they seemed to be in
7 the late 19th century. It is probably due to changes in weeding intensity between these two
8 periods. It is only recently that laws beneficial to biodiversity enacted against using pesticides
9 in urban public territories have had an impact. As a result, the Parisian flora did not
10 depauperate over time, which is heart-warming compared to what is observed in all
11 ecosystems worldwide. The arrival of many exotic species offsets the loss of local ones (if
12 they did not cause it) and enabled biodiversity to adapt to warming and other global changes
13 experienced by our cities. Clearly, it enabled insect communities to endure and ensure
14 pollination services.

15 We hope that other policy changes and a better integration of biodiversity in development
16 projects will lead to even better biodiversity quality and better ecosystem functioning, thereby
17 fostering future city dwellers' well-being.

18 **Funding:** The research was funded by the National Museum of Natural History and the
19 URBANATURE program.

20 **Acknowledgement:** We thank Juliette Amara and Alice Amoureux who initiated the study in
21 2020, during their internship at the National Museum of Natural History in Paris. We are also
22 grateful to the specialist language editing service of Dominique Macabies for the language
23 revisions.

24 **Author contributions:** Conceptualization : N Machon, Inventory (2021) : M Gallant-
25 Canguilhem, Data analyses : M Gallant-Canguilhem, L Ropars and J Vallet, writing original

1 draft : N Machon, writing-review : M Gallant-Canguilhem, L Ropars and J Vallet, supervision
2 : N Machon, J Vallet.

3 **Bibliography**

- 4 Aerts R, Honnay O, and Van Nieuwenhuysse A. 2018. Biodiversity and Human Health:
5 Mechanisms and Evidence of the Positive Health Effects of Diversity in Nature and Green
6 Spaces.” *British Medical Bulletin* 127 (1): 5–22. <https://doi.org/10.1093/bmb/ldy021>.
- 7 Aronson MFJ, Nilon CH, Lepczyk CA, Parker TS, Warren PS, Cilliers SS, Goddard MA, *et*
8 *al.* 2016. Hierarchical Filters Determine Community Assembly of Urban Species Pools.”
9 *Ecology* 97 (11): 2952–63. <https://doi.org/10.1002/ecy.1535>.
- 10 Bocci G. 2015. TR8: An R Package for Easily Retrieving Plant Species Traits. Edited by
11 Richard Fitzjohn. *Methods in Ecology and Evolution* 6 (3): 347–50.
12 <https://doi.org/10.1111/2041-210X.12327>.
- 13 Cadotte MW, Carscadden K, Mirotchnick N. 2011. Beyond Species: Functional Diversity and
14 the Maintenance of Ecological Processes and Services: Functional Diversity in Ecology and
15 Conservation. *Journal of Applied Ecology* 48 (5): 1079–87. <https://doi.org/10.1111/j.1365-2664.2011.02048.x>.
- 17 Chocholoušková Z, Pyšek P. 2003. Changes in Composition and Structure of Urban Flora
18 over 120 Years: A Case Study of the City of Plzeň. *Flora - Morphology, Distribution,*
19 *Functional Ecology of Plants* 198 (5): 366–76. <https://doi.org/10.1078/0367-2530-00109>.
- 20 Lee DR. 1945. Measures of the Amount of Ecologic Association Between Species. *Ecology*
21 26 (3): 297–302. <https://doi.org/10.2307/1932409>.
- 22 Diekmann M. 2003. Species Indicator Values as an Important Tool in Applied Plant Ecology
23 – a Review. *Basic and Applied Ecology* 4 (6): 493–506. <https://doi.org/10.1078/1439-1791-00185>.
- 24 00185.

1 Duchenne F, Gabrielle Martin G, and Porcher E. 2021. European Plants Lagging behind
2 Climate Change Pay a Climatic Debt in the North, but Are Favoured in the South. Edited by
3 Regan Early. *Ecology Letters* 24 (6): 1178–86. <https://doi.org/10.1111/ele.13730>.

4 Ellenberg H, ed. 1991. *Biological Monitoring: Signals from the Environment*. Braunschweig:
5 Vieweg.

6 Garnier E, Cortez J, Billès G, Navas ML, Roumet C, Debussche M, Laurent G, *et al.* 2004.
7 Plant functional markers capture ecosystem properties during secondary succession. *Ecology*
8 85 (9): 2630–37. <https://doi.org/10.1890/03-0799>.

9 Godefroid, S. 2001. Temporal Analysis of the Brussels Flora as Indicator for Changing
10 Environmental Quality. *Landscape and Urban Planning* 52 (4): 203–24.
11 [https://doi.org/10.1016/S0169-2046\(00\)00117-1](https://doi.org/10.1016/S0169-2046(00)00117-1).

12 Grimm NB., Faeth SH, Golubiewski NE, Redman CL, Wu J, Bai X, Briggs JM. 2008. Global
13 Change and the Ecology of Cities. *Science* 319 (5864): 756–60.
14 <https://doi.org/10.1126/science.1150195>.

15 Harrison T, Winfree R. 2015. Urban Drivers of Plant- pollinator Interactions. Edited by Karl
16 Evans. *Functional Ecology* 29 (7): 879–88. <https://doi.org/10.1111/1365-2435.12486>.

17 Haussmann GE. 1890. *Mémoires Du Baron Haussmann I Avant l'Hotel de Ville*. Victor
18 Havard Editeur. Paris.

19 Julve P. 1998. Baseflor. Index Botanique, Écologique et Chorologique de La Flore de France.
20 Lille, France: Institut Catholique de Lille.

21 Landolt E. 2000. Some Results of a Floristic Inventory within the City of Zürich. *Preslia*
22 72,441-455.

23 Lundholm JT, Marlin A. 2006. Habitat Origins and Microhabitat Preferences of Urban Plant
24 Species. *Urban Ecosystems* 9 (3): 139–59. <https://doi.org/10.1007/s11252-006-8587-4>.

1 Maxwell SL., Fuller RA, Brooks TM, Watson JEM. 2016. Biodiversity: The Ravages of
2 Guns, Nets and Bulldozers. *Nature* 536 (7615): 143–45. <https://doi.org/10.1038/536143a>.

3 McGill B, Enquist B, Weiher E, Westoby M. 2006. Rebuilding Community Ecology from
4 Functional Traits. *Trends in Ecology & Evolution* 21 (4): 178–85.
5 <https://doi.org/10.1016/j.tree.2006.02.002>.

6 MNHN & OFB (Ed). 2003. Inventaire National Du Patrimoine Naturel (INPN). inpn.mnhn.fr.

7 Moreau JG, and Daverne JJ. 1845. *Manuel Pratique de La Culture Maraichère de Paris*. V.
8 Bouchard-Huzard. <https://books.google.fr/books?id=YclBAAAAIAAJ>.

9 Omar M, Al Sayed N, Barré K, Halwani J, Machon N. 2018. Drivers of the Distribution of
10 Spontaneous Plant Communities and Species within Urban Tree Bases. *Urban Forestry &*
11 *Urban Greening* 35 (October): 174–91. <https://doi.org/10.1016/j.ufug.2018.08.018>.

12 Pellegrini P, Baudry S. 2014. Streets as New Places to Bring Together Both Humans and
13 Plants: Examples from Paris and Montpellier (France). *Social & Cultural Geography* 15 (8):
14 871–900. <https://doi.org/10.1080/14649365.2014.974067>.

15 Ropars L, Dajoz I, Fontaine C, Muratet A, and Geslin B. 2019. Wild Pollinator Activity
16 Negatively Related to Honey Bee Colony Densities in Urban Context. Edited by Wolfgang
17 Blenau. *PLOS ONE* 14 (9): e0222316. <https://doi.org/10.1371/journal.pone.0222316>.

18 Salinitro M, Alessandrini A, Zappi A, Tassoni A. 2019. Impact of Climate Change and Urban
19 Development on the Flora of a Southern European City: Analysis of Biodiversity Change over
20 a 120-Year Period. *Scientific Reports* 9 (1): 9464. [https://doi.org/10.1038/s41598-019-46005-](https://doi.org/10.1038/s41598-019-46005-1)
21 1.

22 Schmidt KJ, Poppendieck HH, Jensen K. 2014. Effects of Urban Structure on Plant Species
23 Richness in a Large European City. *Urban Ecosystems* 17 (2): 427–44.
24 <https://doi.org/10.1007/s11252-013-0319-y>.

1 Serret H, Raymond R, Foltête JC, Clergeau P, Simon L, Machon N. 2014. Potential
2 Contributions of Green Spaces at Business Sites to the Ecological Network in an Urban
3 Agglomeration: The Case of the Ile-de-France Region, France. *Landscape and Urban*
4 *Planning* 131: 27–35. <https://doi.org/10.1016/j.landurbplan.2014.07.003>.

5 Shwartz A, Muratet A, Simon L, Julliard R. 2013. Local and Management Variables
6 Outweigh Landscape Effects in Enhancing the Diversity of Different Taxa in a Big
7 Metropolis. *Biological Conservation* 157 (January): 285–92.
8 <https://doi.org/10.1016/j.biocon.2012.09.009>.

9 Silvertown J, McConway K, Gowing D, Dodd M, Fay MF, Joseph JA, Dolphin K. 2006.
10 Absence of Phylogenetic Signal in the Niche Structure of Meadow Plant Communities.
11 *Proceedings of the Royal Society B: Biological Sciences* 273 (1582): 39–44.
12 <https://doi.org/10.1098/rspb.2005.3288>.

13 Vallot J. 1884. *Essai Sur La Flore Du Pavé de Paris Limité Aux Boulevards Extérieurs, Ou*
14 *Catalogue Des Plantes: Qui Croissent Spontanément Dans Les Rues et Sur Les Quais &*
15 *Florule Des Ruines Du Conseil d'État*. Editions Jacques Lechevalier. Paris.

16 Williams GJ, Graham NAJ, Jouffray JB, Norström AV, Nyström M, Gove JM, Heenan A,
17 and Wedding LM. 2019. Coral Reef Ecology in the Anthropocene. Edited by Charles Fox.
18 *Functional Ecology* 33 (6): 1014–22. <https://doi.org/10.1111/1365-2435.13290>.

19
20

1 **List of figures**

2 Figure 1: Map of the 52 Paris's public spaces inventoried both in 1884 and 2021 (in red)

3 Figure 2: Number of species inventoried by Vallot in 1884 in 52 public spaces in Paris, and
4 number of species inventoried by the authors in 2021 in the same spaces

5 Figure 3: Plots of mean dependency to pollinators (PD) and mean Ellenberg indicator values
6 per public space for each trait: L light; T temperature; Ha atmospheric humidity; Hs soil
7 humidity; pH soil pH; N nitrate; and Sal salinity. Ns means not significant, **** p-value >
8 0.001, * 0.01 < p-value < 0.05

9 Figure 4: Dendrogram of the floristic Jaccard's distances among streets in 1884 (in green) and
10 in 2021 (in black)

11 Figure 5: Dendrogram of the floristic Jaccard's distances along the Seine wharves in 1884 (in
12 green) and in 2021 (in black) 19th century

13

1 **Table 1:** Wilcoxon's paired series median comparison tests: PD is the pollinator dependency;
2 L light; T temperature; Ha atmospheric humidity; Hs soil humidity; pH soil pH; N nitrate and
3 Sal salinity Ellenberg indicator values.

4

	1884	2021	u-stat	p-value
PD (%)	36.39±20.6	45.13±4.56	434.0	0.02
L	7.13±0.4	7.15±0.21	558.0	0.591
T	5.21±0.27	5.69±0.16	34.0	<0.001
Ha	4.68±0.55	4.65±0.18	756.0	0.545
Hs	4.75±0.69	4.73±0.16	929.5	0.013
pH	5.82±0.54	6.24±0.17	186.0	<0.001
N	6.73±1.26	6.76±0.24	782.0	0.4
Sal	0.23±0.27	0.16±0.08	698.5	0.743

5

6

7