



HAL
open science

Editorial: African vertebrates through times

Nour-Eddine Jalil, Emmanuel Gheerbrant, Martin Rücklin

► **To cite this version:**

Nour-Eddine Jalil, Emmanuel Gheerbrant, Martin Rücklin. Editorial: African vertebrates through times. *Frontiers in Earth Science*, 2023, 10, <10.3389/feart.2022.1111339>. <mnhn-04235456>

HAL Id: mnhn-04235456

<https://mnhn.hal.science/mnhn-04235456v1>

Submitted on 10 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



OPEN ACCESS

EDITED AND REVIEWED BY
Bruce S. Lieberman,
University of Kansas, United States

*CORRESPONDENCE
Nour-Eddine Jalil,
✉ nour-eddine.jalil@mnhn.fr

SPECIALTY SECTION
This article was submitted to Paleontology,
a section of the journal
Frontiers in Earth Science

RECEIVED 29 November 2022
ACCEPTED 16 December 2022
PUBLISHED 09 January 2023

CITATION
Jalil N-E, Gheerbrant E and Rücklin M
(2023), Editorial: African vertebrates
through times.
Front. Earth Sci. 10:1111339.
doi: 10.3389/feart.2022.1111339

COPYRIGHT
© 2023 Jalil, Gheerbrant and Rücklin. This
is an open-access article distributed under
the terms of the [Creative Commons
Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use,
distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in
this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Editorial: African vertebrates through times

Nour-Eddine Jalil^{1,2*}, Emmanuel Gheerbrant¹ and Martin Rücklin³

¹UMR7207 Centre de Recherche en paléontologie- Paris (CR2P) National Museum of Natural History, Paris, France, ²Museum of Natural History of Marrakech, Department of Geology FSSM, Cadi Ayyad University, Marrakech, Morocco, ³Naturalis Biodiversity Center, Sylvius Laboratory, Leiden University, Leiden, Netherlands

KEYWORDS

Africa, vertebrates, fossiles, anatomy, phylogeny, biogeography

Editorial on the Research Topic African vertebrates through times

The first record of vertebrate fossils from Africa can be traced back to the beginning of the nineteenth century, almost two centuries of paleontological discoveries. Today Africa is an important source of knowledge in regard to the evolution of vertebrates and plays a pivotal role in understanding the impact of geodynamic events in macroevolution (extinctions and succeeding recovery phases). It is a key continental theater for the study of endemic evolutionary radiations. This volume despite the small number of its items, covers the three geological eras since it presents the results of new research on Paleozoic, Mesozoic and Cenozoic vertebrates from Africa.

The placoderms also called “Armored fishes”, because of the heavily mineralized plates that protect their heads and thorax, are extinct early jawed vertebrates that lived from the Silurian to the end of the Devonian period. They were the dominant marine vertebrates during the Devonian being abundant, diversified and adapted to various environments and food strategies. As one of the first jawed vertebrates (or gnathostomes), their study is crucial to understand the origin and evolution of the gnathostome body plan. [Jobbins et al.](#) described an exquisitely well preserved articulated new selenosteoid placoderm from the Devonian of Morocco. They add to the biodiversity of Devonian placoderms from Morocco and thanks to the exceptional preservation of the body outline, they shed new lights on the body shape and ecomorphology of the selenosteoids.

During the last decades new paleontological findings have revealed exceptional mosasaur diversity from the Northern part of Africa. The mosasaurs were a specialized group of squamates that adapted to marine life in the mid-Cretaceous, before becoming the dominant marine predators in the latest Cretaceous. They were the most common reptiles and the dominant marine reptiles of the Late Cretaceous. [Woolley et al.](#) redescribed mosasaur remains collected at the beginning of the last century from the latest Cretaceous of South Africa. By the identification of at least three taxa, this study revealed an unsuspected mosasaur diversity in South Africa. It also underlines the potential of the Cretaceous outcrops in the east coast of South Africa for new finds and in the understanding of the diversity and dispersal of mosasaurs in and around Southern Africa.

The Cretaceous “Continental Intercalaire”, which extends across North Africa (the Kem beds of Morocco, Gara Samani in Algeria, Tataouine region in Tunisia, Mizdah Formation in Libya), and its contemporary Baharyia Formation of Egypt and Echkar Formation in Central Niger provides abundant data on vertebrate paleobiodiversity and evolutionary history from the Early to Late Cretaceous, and shows how dinosaurs evolved in the context of the continental

breakup. These localities raised particular interest among scientists aiming to get a good look at dinosaur life when the African continent drifted into isolated Isle. Noteworthy is the fact that the faunas from these formations are unbalanced by a predominance in number and diversity of the large bodied carnivorous over supplanting that of the herbivorous. A terrestrial ecosystem unparalleled in today's nature. In [Benyoucef et al.](#) the geological context of the "Continental Intercalaire" deposits occurring in the Gara Samani area, southern Algeria is provided. The faunal list yielded by this formation was supplemented thanks to new findings in this area. This study adds to our knowledge on the late Albian-Cenomanian terrestrial vertebrate biodiversity of Algeria.

In the last paper, [Murray](#) described a new species of cichlid « fishes » from Oligocene deposits of Somalia. Cichlids are an important component of the African freshwater ecosystem. This study offers the opportunity to discuss the biogeographic history of the group. Answers on the question if their modern distributions the result of a dispersal event or are they the result of a vicariant event after the break-up of the two continental masses in the mid-Cretaceous are possible. This sheds new lights on the early history of cichlids in Africa.

Author contributions

Original draft by N-EJ; EG and MR review and edit the original draft. All authors approved it for publication.

Conflict of interest

The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.